

Euler Characteristics of Teichmüller Curves in Genus Two

Matt Bainbridge

February 2, 2008

Contents

1	Introduction	4
2	Abelian surfaces and real multiplication	15
2.1	Abelian surfaces	15
2.2	Real multiplication	16
2.3	Hilbert modular surfaces	17
3	Prototypes	32
3.1	P_D , W_D , and Y_D -prototypes	32
3.2	Quasi-invertible $\mathcal{O}_D$ -modules	34
4	Abelian differentials	40
4.1	Flat geometry of Abelian differentials	40
4.2	Moduli of Abelian differentials	44
4.3	$\mathrm{SL}_2\mathbb{R}$ orbits in genus two.	48
5	Deligne-Mumford compactification of moduli space	54
5.1	Stable Riemann surfaces	54
5.2	Deligne-Mumford compactification	56
5.3	Degenerating Abelian differentials	60
6	Local coordinates for strata in $\Omega\overline{\mathcal{M}}_2$	68
6.1	Strata in $\Omega\overline{\mathcal{M}}_2$	68
6.2	The strata $\Omega\mathcal{M}_2(2)$ and $\Omega\mathcal{M}_2(T_{0,1})$	71
6.3	The strata $\Omega\mathcal{M}_2^0(T_{1,0}; 1, 1)$ and $\Omega\mathcal{M}_2^0(T_{1,0}; 2)$	73
6.4	The stratum $\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$	76
6.5	The stratum $\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$	78
6.6	The strata $\Omega\mathcal{M}_2^0(T_{2,0}; 1, 1)$, $\Omega\mathcal{M}_2^0(T_{2,0}; 2)$, and $\Omega\mathcal{M}_2^0(T_{2,1})$	79
6.7	The stratum $\Omega\mathcal{M}_2^0(T_{3,0})$	82
6.8	The stratum $\Omega\mathcal{M}_2^0(T_{3,0}, T_{3,0}^3)$	83

7	Limits of eigenforms	85
7.1	Introduction	85
7.2	Empty strata	87
7.3	The strata $X_D(T_{1,0}; 1, 1)$ and $X_D(T_{1,0}, 2)$	88
7.4	The stratum $X_D(T_{1,0}, T_{1,0}^1)$	92
7.5	The stratum $X_D(T_{2,0}, T_{2,0}^2)$	95
7.6	The strata $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, and $X_D(T_{2,1})$	97
7.7	The strata $X_D(T_{3,0})$ and $X_D(T_{3,0}, T_{3,0}^3)$	106
8	Geometric compactification of X_D	112
8.1	Geometry of Y_D	112
8.2	Y_D maps to $\widehat{X}_D$	120
8.3	Involution of Y_D	124
9	Line bundles over Y_D	128
9.1	Orbifold vector bundles	128
9.2	Good metrics	129
9.3	Extension of the bundle ΩX_D to Y_D	132
9.4	Extension of the foliation $\mathcal{A}_D$ to Y_D	136
10	Euler characteristic of W_D	141
11	Fundamental class of $\overline{W}_D$	145
12	Normal Bundles	149
13	Euler characteristic of W_D^ϵ	152
14	Siegel-Veech Constants	158
15	Lyapunov Exponents	162
A	Normal varieties	170

Abstract

For any integer $D \equiv 0$ or $1 \pmod{4}$, there is a Teichmüller curve W_D immersed in the moduli space $\mathcal{M}_2$ of genus two Riemann surfaces. The curve W_D is naturally embedded in a Hilbert modular surface X_D . Our main result is that the Euler characteristic of W_D is proportional to the Euler characteristic of X_D . More precisely,

$$\chi(W_D) = -\frac{9}{2}\chi(X_D),$$

when D is not square.

When $D \equiv 1 \pmod{8}$, the curve W_D has two connected components, W_D^0 and W_D^1 . We also calculate the Euler characteristics of these components. When D is not square, we show that

$$\chi(W_D^0) = \chi(W_D^1).$$

When D is square, we show that

$$\begin{aligned}\chi(W_{d^2}^0) &= -\frac{1}{32}d^2(d-1) \sum_{r|d} \frac{\mu(r)}{r^2}, \quad \text{and} \\ \chi(W_{d^2}^1) &= -\frac{1}{32}d^2(d-3) \sum_{r|d} \frac{\mu(r)}{r^2}.\end{aligned}$$

The idea of the calculation of $\chi(W_D)$ is to use techniques from algebraic geometry to compute the fundamental class of the closure W_D in a compactification of X_D . We define a compactification Y_D of X_D which maps to the Deligne-Mumford compactification of $\mathcal{M}_2$ by a finite morphism. We then exhibit W_D as the zero locus of a meromorphic section of a line bundle over Y_D , which allows us to calculate the fundamental class of W_D .

To calculate the Euler characteristics of the connected components W_D^ϵ , we find several relations involving the fundamental class of the closure of W_D^ϵ which allow us to solve for $\chi(W_D^\epsilon)$. For example, we calculate the self-intersection numbers of the W_D^ϵ in terms of $\chi(W_D^\epsilon)$.

We apply these results to calculate the Siegel-Veech constants for counting closed billiards paths in certain L-shaped polygons. We then calculate the Lyapunov exponents of the Kontsevich-Zorich cocycle for any ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measure on the moduli space $\Omega_1\mathcal{M}_2$ of Abelian differentials in genus two (previously calculated in unpublished work of Kontsevich and Zorich).

1 Introduction

Let $\mathcal{M}_2$ be the moduli space of genus two Riemann surfaces. Given any $D > 0$ with $D \equiv 0$ or $1 \pmod{4}$, let $W'_D \subset \mathcal{M}_2$ be the locus of Riemann surfaces X such that:

- The Jacobian $\text{Jac}(X)$ has real multiplication by $\mathcal{O}_D$, the unique real quadratic order of discriminant D , and
- There is an Abelian differential ω on X which is an eigenform for real multiplication and has a double zero.

Let W_D be the normalization of W'_D , a possibly disconnected curve.

The *Hilbert modular surface*,

$$X_D = \mathbb{H} \times \mathbb{H} / \text{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee),$$

is the moduli space for Abelian surfaces with real multiplication by $\mathcal{O}_D$. In X_D , let P_D be the Shimura curve consisting of those Abelian surfaces in X_D which are polarized products of elliptic curves. There is a commutative diagram,

$$\begin{array}{ccc} W_D & \hookrightarrow & X_D \setminus P_D \\ & \searrow & \downarrow \\ & & \mathcal{M}_2 \end{array}$$

where the top map is an embedding, and the vertical map is a two-to-one map sending an Abelian surface A to the unique Riemann surface $X \in \mathcal{M}_2$ such that $\text{Jac}(X) \cong A$.

The curve W_D is not a Shimura curve on X_D ; it is, however, a *Teichmüller curve*, a curve which is isometrically immersed in the moduli space $\mathcal{M}_2$ with respect to the Teichmüller metric. In fact, the curves W_D with D nonsquare are all but one of the *primitive* Teichmüller curves in $\mathcal{M}_2$, where a Teichmüller curve is said to be primitive if it does not arise from a Teichmüller curve of lower genus by a certain branched covering construction. The curves W_D arise from the study of billiards in certain L -shaped polygons, and the study of these curves has applications to the dynamics of billiards in these polygons.

Euler characteristics. The main object of this paper is to calculate the Euler characteristics of the curves W_D . The idea is to relate $\chi(W_D)$ to $\chi(X_D)$ and $\chi(P_D)$. It turns out that these Euler characteristics are all proportional if D is not square. The following is our main result.

Theorem 1.1. *If D is not square, then*

$$\chi(W_D) = -\frac{9}{2}\chi(X_D). \tag{1.1}$$

Remark. Zagier conjectured formula (1.1) from numerical evidence. In [Sie36], Siegel calculated the volume of X_D when D is a fundamental discriminant (that is D is not of the form $D = f^2 E$ for some $E \equiv 0$ or $1 \pmod{4}$ and $f > 1$). This yields formulas for $\chi(X_D)$ for all D , and for $D \neq 1$ a fundamental discriminant, we obtain

$$\chi(W_{f^2 D}) = -9\zeta_{\mathbb{Q}(\sqrt{D})}(-1)f^3 \sum_{r|f} \left(\frac{D}{r}\right) \frac{\mu(r)}{r^2},$$

where μ is the Möbius function, and $\left(\frac{D}{r}\right)$ is the Kronecker symbol. When $f = 1$, this reduces to

$$\chi(W_D) = -9\zeta_{\mathbb{Q}(\sqrt{D})}(-1).$$

The Euler characteristics $\chi(W_D)$ when D is not square are given by the Fourier coefficients of a modular form. More precisely, there is a function $H(2, D)$ such that for $D \neq 1$ a fundamental discriminant and $f \in \mathbb{N}$,

$$\sum_{r|f} \chi(W_{r^2 D}) = \frac{3}{4} H(2, f^2 D), \quad \text{and}$$

$$\sum_{\substack{D \equiv 0,1 \pmod{4} \\ D \geq 0}} H(2, D) q^D$$

is a modular form studied by Cohen in [Coh75]. See §2.3 for more about this form.

When $D = d^2$, Eskin, Masur, and Schmoll calculated $\chi(W_{d^2})$ in [EMS03]. Our methods give a new proof of their formula. In this case, it is no longer true that $\chi(W_{d^2})$ is proportional to $\chi(X_{d^2})$.

Theorem 1.2. *For any $d > 1$,*

$$\chi(W_{d^2}) = -\frac{1}{16} d^2 (d-2) \sum_{r|d} \frac{\mu(r)}{r^2}. \quad (1.2)$$

Connected components of W_D . It is known that when $D \neq 9$ and $D \equiv 1 \pmod{8}$, W_D has two connected components, W_D^0 and W_D^1 ; otherwise W_D is connected [McM05a].

We will also show that when D is not square, the connected components of W_D have the same Euler characteristic. It should be true that these components are actually homeomorphic, but we have not proved this. It is not true that the connected components of W_D are isomorphic as curves over $\mathbb{C}$.

Theorem 1.3. *If $D \equiv 1 \pmod{8}$ is not square, then*

$$\chi(W_D^0) = \chi(W_D^1).$$

There is a canonical involution $\tau: X_D \rightarrow X_D$ which is covered by the involution $(z, w) \mapsto (w, z)$ of $\mathbb{H} \times \mathbb{H}$. This involution replaces an Abelian surface A together with a choice of real multiplication,

$$\rho: \mathcal{O}_D \rightarrow \text{End } A,$$

with the Galois conjugate real multiplication ρ' obtained by precomposing ρ with the Galois automorphism of $\mathcal{O}_D$. It is not true that τ permutes the connected components of W_D (otherwise Theorem 1.3 would be trivial). In fact, W_D is not even invariant under τ .

We also calculate $\chi(W_{d^2}^\epsilon)$, but it is no longer true in the square discriminant case that the connected components of W_{d^2} have the same Euler characteristic.

Theorem 1.4. *If $d > 0$ and $d^2 \equiv 1 \pmod{8}$, then*

$$\begin{aligned} \chi(W_{d^2}^0) &= -\frac{1}{32}d^2(d-1) \sum_{r|d} \frac{\mu(r)}{r^2}, \quad \text{and} \\ \chi(W_{d^2}^1) &= -\frac{1}{32}d^2(d-3) \sum_{r|d} \frac{\mu(r)}{r^2}. \end{aligned} \tag{1.3}$$

Remark. These formulas were established independently by Lelièvre and Royer in [LR06].

Siegel-Veech constants. Given a positive, nonsquare integer $D \equiv 0$ or $1 \pmod{4}$, let $P(D)$ be the L-shaped polygon as in Figure 1, where the side lengths are given by

$$a = b = \frac{1 + \sqrt{D}}{2}$$

if D is odd, and

$$a = \frac{\sqrt{D}}{2} \quad \text{and} \quad b = 1 + \frac{\sqrt{D}}{2}$$

if D is even. By a standard unfolding construction, a L -shaped polygon determines a genus two Riemann surface X equipped with an Abelian differential ω , and the (X, ω) determined by $P(D)$ lies on ΩW_D .

A billiards path on $P(D)$ is a path which is geodesic on the interior of $P(D)$ and bounces off the walls as a physical billiards ball would (angle of incidence equals angle of reflection). We allow billiards paths to pass through the corners of angle $\pi/2$ but not through the one of angle $3\pi/2$. Closed billiards paths occur in parallel families which are bounded by two paths which each start and end at the corner of angle $3\pi/2$. Let

$$N(P(D), L) = \#\{\text{families of simple closed billiards paths on } P(D) \\ \text{of length at most } L\}.$$




Figure 1: $P(D)$

Note that we do not count paths which go around a single closed path several times. Also paths are not oriented, so we do not count a path going once in each direction. Since the unfolding lies on a Teichmüller curve, we have by [Vee89]

$$N(P(D), L) \sim c(D) \frac{\pi}{4 \text{Area}(P(D))} L^2.$$

for some constants $c(D)$ called Siegel-Veech constants. Evaluating the constants $c(D)$ amounts to calculating the Euler characteristics of the W_D together with the volumes of certain neighborhoods of the cusps of W_D . In §14, we calculate these constants. For example, we obtain:

Theorem 1.5. *For small values of D , the Siegel-Veech constants $c(D)$ are as given in Table 1.*

D	5	8	12	13	17	20	21	24	28	29
$c(D)$	$\frac{25}{3}$	$\frac{28}{3}$	$\frac{26}{3}$	$\frac{91}{9}$	$\frac{221}{24} + \frac{1}{8}\sqrt{17}$	$\frac{31}{3}$	$\frac{133}{15}$	$\frac{148}{15}$	$\frac{82}{9}$	$\frac{377}{35}$

Table 1: Siegel-Veech constants for $P(D)$

Note that there are other L-shaped polygons, classified in [McM03a], whose unfoldings lie on the Teichmüller curves W_D and our results apply equally well to those. The Siegel-Veech constants only depend on the connected component of W_D on which the unfolding lies. We also show in §14:

Theorem 1.6. *If W_D has two connected components, then the associated Siegel-Veech constants are Galois-conjugate elements of $\mathbb{Q}(\sqrt{D})$. If W_D is connected, then the Siegel-Veech constants are rational.*

The Kontsevich-Zorich cocycle. There is a well-known action of $\mathrm{SL}_2\mathbb{R}$ on $\Omega\mathcal{M}_g$, the moduli space of Abelian differentials, which preserves the subspace $\Omega_1\mathcal{M}_g$ of Abelian differentials of norm one with respect to the norm,

$$\|\omega\| = \left(\int_X |\omega|^2 \right)^{1/2},$$

on $\Omega(X)$. The diagonal subgroup $A \subset \mathrm{SL}_2\mathbb{R}$ induces a flow g_t on $\Omega\mathcal{M}_g$, the *Teichmüller geodesic flow*. Kontsevich and Zorich [Kon97, Zor99] introduced a linear cocycle over this flow which is closely related to the dynamics of flows on surfaces. There is a bundle $\mathcal{H}_1(\mathbb{R}) \rightarrow \Omega_1\mathcal{M}_g$ whose fiber over an Abelian differential (X, ω) is $H_1(X; \mathbb{R}) / \mathrm{Aut}(X, \omega)$. This bundle has a natural flat connection, the *Gauss-Manin connection*. By coupling the geodesic flow g_t with the Gauss-Manin connection, we get a flow $\tilde{g}_t$, the *Kontsevich-Zorich cocycle*, on $\mathcal{H}_1(\mathbb{R})$ which is linear on the fibers and covers the flow g_t on $\Omega_1\mathcal{M}_g$.

Consider the closure $S \subset \Omega_1\mathcal{M}_2$ of an $\mathrm{SL}_2\mathbb{R}$ orbit. By results of [McMa], S is a suborbifold of $\Omega_1\mathcal{M}_2$ which comes equipped with a unique $\mathrm{SL}_2\mathbb{R}$ -invariant, ergodic, absolutely continuous, probability measure, the *period measure*. In the case of the Teichmüller curves, period measure is just the normalized hyperbolic area measure (because Haar measure is unique).

Associated to any ergodic, g_t -invariant measure μ are the *2g Lyapunov exponents*,

$$1 = \lambda_1(\mu) > \cdots > \lambda_g(\mu) > -\lambda_g(\mu) > \cdots > -\lambda_1(\mu).$$

Kontsevich gave a formula in [Kon97] which calculates $\sum \lambda_i(\mu)$ as a certain integral over moduli space when μ is $\mathrm{SL}_2\mathbb{R}$ invariant. The same methods which we use to calculate $\chi(W_D)$ allow us to evaluate these integrals and to calculate $\lambda_2(\mu)$ for all ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measures on $\Omega_1\mathcal{M}_2$. Let $\Omega_1\mathcal{M}_2(2) \subset \Omega_1\mathcal{M}_2$ be the locus of forms with a double zero.

Theorem 1.7. *If μ is any finite, ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measure on $\Omega_1\mathcal{M}_2$, then*

$$\lambda_2(\mu) = \begin{cases} 1/3, & \text{if } \mu \text{ is supported on } \Omega_1\mathcal{M}_2(2); \\ 1/2, & \text{if } \mu \text{ is not supported on } \Omega_1\mathcal{M}_2(2). \end{cases}$$

Remark. Theorem 1.7 is an unpublished result of Kontsevich and Zorich which is mentioned in [Zor06].

Outline of proof of Theorem 1.1. The strategy for computing the Euler characteristic of W_D is to study the relationship between W_D and the Hilbert modular surface X_D on which it lies. More precisely, we will define a compactification Y_D of X_D and compute the fundamental class of $\overline{W}_D$ in Y_D by expressing it as the zero locus of a meromorphic section of a line bundle over Y_D .

Here is a sketch of the calculation of the Euler characteristic $\chi(W_D)$ in the case when D is nonsquare. When D is a square, the calculation is more complicated because there are some extra curves in $Y_{d^2} \setminus X_{d^2}$ which have to be taken into account.

1. Given a nonsquare $D \in \mathbb{N}$ with $D \equiv 0$ or $1 \pmod{4}$, let

$$\mathcal{O}_D = \mathbb{Z}[T]/(T^2 + bT + c)$$

for integers $b, c \in \mathbb{Z}$ such that $b^2 - 4c = D$. The discriminant D determines $\mathcal{O}_D$ up to isomorphism, and $\mathcal{O}_D$ is naturally embedded in its field of fractions K_D , which is isomorphic to $\mathbb{Q}(\sqrt{D})$. Let $\mathcal{O}_D^\vee \subset K_D$ be the inverse different of $\mathcal{O}_D$, and let $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee) \subset \mathrm{SL}_2 K_D$ be the subgroup which preserves $\mathcal{O}_D \oplus \mathcal{O}_D^\vee$.

There are two closed two-forms ω_i on

$$X_D = \mathbb{H} \times \mathbb{H} / \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$$

covered by the forms

$$\frac{1}{2\pi} \frac{dx_i \wedge dy_i}{y_i^2}. \quad (1.4)$$

on $\mathbb{H} \times \mathbb{H}$. The inverse image of W_D in $\mathbb{H} \times \mathbb{H}$ is the union of the graphs of countably many holomorphic functions $\mathbb{H} \rightarrow \mathbb{H}$. It follows that the restriction of ω_1 to W_D is $1/2\pi$ times the hyperbolic volume form on W_D . Gauss-Bonnet then tells us that

$$\chi(W_D) = - \int_{W_D} \omega_1.$$

2. There is a Shimura curve $P_D \subset X_D$ which parameterizes Abelian surfaces which are polarized products of elliptic curves. It is covered in $\tilde{X}_D$ by a union of graphs of Möbius transformations $\mathbb{H} \rightarrow \mathbb{H}$. We introduce these curves in §2.3 and show that

$$\chi(P_D) = -\frac{5}{2}\chi(X_D). \quad (1.5)$$

3. The bundle $\Omega\mathcal{M}_2$ over $\mathcal{M}_2$ whose fiber over a Riemann surface X is the space $\Omega(X) \setminus \{0\}$ of nonzero Abelian differentials on X extends to a bundle $\Omega\overline{\mathcal{M}}_2$ over the Deligne-Mumford compactification which we discuss in §5. In §7 and §8, we construct a compactification Y_D of X_D , a complex projective orbifold which is obtained by taking the normalization of the closure of the image of an embedding $X_D \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$. We call this the *geometric compactification* of X_D . The complement $\partial X_D = Y_D \setminus X_D$ consists of finitely many rational curves, each labeled by a discrete invariant which we call a Y_D -prototype. We discuss these invariants in §3. For each Y_D -prototype P , there is one rational curve $C_P \subset \partial X_D$.
4. Associated to each point $p \in Y_D$, there is a stable Riemann surface $X \in \overline{\mathcal{M}}_2$ together with an action of $\mathcal{O}_D$ on $\mathrm{Jac}(X)$ by real multiplication. This determines a splitting of the space $\Omega(X)$ of stable Abelian differentials on X into two eigenspaces for real multiplication,

$$\Omega(X) = \Omega^1(X) \oplus \Omega^2(X).$$

(See §5 for information about stable Riemann surfaces and stable Abelian differentials.) A choice of an embedding $\mathcal{O}_D \rightarrow \mathbb{R}$ determines one of these two eigenspaces, so the point $p \in Y_D$ determines an eigenform in $\Omega(X)$ up to scale if we fix such an embedding. There is a holomorphic line bundle ΩY_D whose fiber over $p \in Y_D$ is the eigenspace $\Omega^1(X)$.

In §9, we study a holomorphic foliation $\mathcal{A}_D$ of Y_D along whose leaves the absolute periods of eigenforms in ΩY_D are locally constant. Each rational curve $C_P \subset \partial X_D$ is a leaf of $\mathcal{A}_D$, and $\mathcal{A}_D$ has isolated singularities at the intersection of two curves C_P and C_Q . A leaf of $\mathcal{A}_D$ can be locally parameterized away from W_D and P_D by the *relative periods* of (X, ω) , the integrals along paths joining the two zeros ω . Following [McMb], we use this parameterization to define a quadratic differential on each leaf of $\mathcal{A}_D$. These quadratic differentials can be pieced together to define a section q of the line bundle

$$\mathcal{L} = (\Omega Y_D)^{-2} \otimes (T^* \mathcal{A}_D)^2$$

over Y_D . This section q vanishes along $\overline{W}_D$, has a simple pole along $\overline{P}_D$, and is elsewhere finite and nonzero. This gives a relation between the fundamental classes of $\overline{W}_D$, $\overline{P}_D$, and the Chern class of $\mathcal{L}$, resulting in the formula,

$$[\overline{W}_D] = [\overline{P}_D] + c_1(\mathcal{L}). \quad (1.6)$$

in $H^2(Y_D; \mathbb{Q})$. We derive this formula in §10. This is the main idea of the calculation of $\chi(W_D)$, and the reader who only wants the gist of the argument might find it useful to skip directly to this section.

5. In §9, we calculate $c_1(\mathcal{L})$. We define a Hermitian metric h on ΩY_D which is singular along ∂X_D . We show that h is a *good metric* in the sense of Mumford [Mum77]. This implies that the Chern form $c_1(\Omega Y_D, h) = \omega_1/2$ is a closed current on Y_D , and

$$c_1(\Omega Y_D) = \frac{1}{2}[\omega_1] \quad (1.7)$$

in $H^2(Y_D; \mathbb{Q})$.

The canonical involution τ of X_D extends to an involution of Y_D , which we continue to call τ . We also show that,

$$\tau^*(\Omega Y_D)^2 = T^* \mathcal{A}_D,$$

which implies,

$$c_1(T^* \mathcal{A}_D) = [\omega_2]. \quad (1.8)$$

Putting together (1.7) and (1.8), we obtain

$$c_1(\mathcal{L}) = -[\omega_1] + 2[\omega_2]. \quad (1.9)$$

6. Combining (1.6) and (1.9), we obtain

$$[\overline{W}_D] = [\overline{P}_D] - [\omega_1] + 2[\omega_2]. \quad (1.10)$$

By pairing $-\omega_1$ with both sides and applying the Gauss-Bonnet Theorem, we obtain

$$\chi(W_D) = \chi(P_D) - 2\chi(X_D). \quad (1.11)$$

Putting (1.5) in (1.11), we obtain (1.1).

The proof of Theorem 1.1 uses very little about the compactification Y_D . Equation (1.10) is true, when interpreted as an equation of cohomology classes of closed currents on X_D ; however, we are not allowed to pair ω_1 with this equation because ω_1 is not compactly supported. Working in the compactification Y_D allows us to justify this pairing. The proof of Theorem 1.3 uses the compactification Y_D in a more essential way.

When $D = d^2$, there are some extra curves $S_{d^2}^1$ and $S_{d^2}^2$ in $Y_{d^2} \setminus X_{d^2}$, which complicates the calculation of $\chi(W_{d^2})$. One difference from the square case is that the section q of $\mathcal{L}$ vanishes on $\overline{S}_{d^2}^2$ as well as $\overline{P}_{d^2}$. Also, (1.8) becomes

$$c_1(T^*\mathcal{A}_D) = [\omega_2] - [\overline{S}_{d^2}^2].$$

Instead of (1.11), we get

$$\chi(W_{d^2}) = \chi(P_{d^2}) - 2\chi(X_{d^2}) - \chi(S_{d^2}^2).$$

In this case, $\chi(P_{d^2})$ and $\chi(S_{d^2}^2)$ are not proportional to $\chi(X_{d^2})$, so neither is $\chi(W_{d^2})$.

Outline of proof of Theorem 1.3. The idea of the proof of Theorem 1.3 is to find as many equations as we can involving the fundamental classes $[\overline{W}_D^0]$ and $[\overline{W}_D^1]$. Once we have enough equations, we are able to solve for the pairing $[\omega_1] \cdot [\overline{W}_D^e] = -\chi(W_D^e)$.

1. In §7 and §8, we will prove the following properties of Y_D :

- (a) The closures $\overline{W}_D$ and $\overline{P}_D$ are disjoint suborbifolds of Y_D .
- (b) Each rational curve $C_P \subset Y_D \setminus X_D$ meets $\overline{W}_D$ and $\overline{P}_D$ transversely and meets each in the same number of points.
- (c) The canonical involution τ of Y_D has the property that for each rational curve $C_P \subset \partial X_D$,

$$\overline{W}_D^0 \cdot C_P = \overline{W}_D^1 \cdot \tau(C_P).$$

(d) The rational cohomology of Y_D is an orthogonal direct sum:

$$H^2(Y_D; \mathbb{Q}) \cong B \oplus \langle [\omega_1], [\omega_2] \rangle \oplus J,$$

where B is the subspace generated by the rational curves $C \subset \partial X_D$, and J is the orthogonal complement of the other two terms. J contains all of $H^{2,0}(Y_D; \mathbb{Q})$ and $H^{0,2}(Y_D; \mathbb{Q})$ and most of $H^{1,1}(Y_D; \mathbb{Q})$.

2. There is a well-known formula relating the first Chern class of the bundle $\Omega \overline{\mathcal{M}}_2$ over $\overline{\mathcal{M}}_2$ to the fundamental classes of divisors in $\overline{\mathcal{M}}_2 \setminus \mathcal{M}_2$. Pulling back this formula by the natural map $Y_D \rightarrow \overline{\mathcal{M}}_2$, we obtain a formula for the fundamental class of $[\overline{P}_D]$:

$$[\overline{P}_D] = \frac{5}{2}[\omega_1] + \frac{5}{2}[\omega_2] + \pi_B[\overline{P}_D], \quad (1.12)$$

where π_B is the orthogonal projection of $H^2(Y_D; \mathbb{Q})$ to B . Combining this with (1.10), we obtain

$$[\overline{W}_D] = \frac{3}{2}[\omega_1] + \frac{9}{2}[\omega_2] + \pi_B[\overline{W}_D].$$

It follows from (1.10) that $\pi_B[\overline{P}_D] = \pi_B[\overline{W}_D]$. We derive these formulas in §11.

3. In §12, we study the normal bundles of $\overline{W}_D$ and $\overline{P}_D$ in Y_D . The restriction of ΩY_D to $\overline{W}_D^\epsilon$ is a line bundle $\Omega \overline{W}_D^\epsilon$. The degree of normal bundle $N(\overline{W}_D^\epsilon)$ can be calculated in terms of $\Omega \overline{W}_D^\epsilon$. We obtain the following formula for the self intersection number of $\overline{W}_D^\epsilon$:

$$[\overline{W}_D^\epsilon]^2 = \deg N(\overline{W}_D^\epsilon) = -\frac{2}{3} \deg \Omega \overline{W}_D^\epsilon = \frac{1}{3} \chi(W_D^\epsilon). \quad (1.13)$$

The proof of the middle equality involves defining a three-to-one map from a tubular neighborhood $\overline{W}_D^\epsilon$ to $(\Omega \overline{W}_D^\epsilon)^{-2}$ via an operation called “collapsing a saddle connection,” which we introduce in §4.1. This operation replaces an Abelian differential together with a saddle connection joining a pair of simple zeros with an Abelian differential with a double zero.

Since $\overline{W}_D^\epsilon$ is transverse to the foliation $\mathcal{A}_D$ of Y_D , we also have the relation,

$$N(\overline{W}_D^\epsilon) \cong T\mathcal{A}_D,$$

which together with (1.8) implies

$$[\overline{W}_D^\epsilon]^2 = \int_{W_D^\epsilon} \omega_2. \quad (1.14)$$

From (1.13) and (1.14), we obtain

$$\int_{W_D^\epsilon} \omega_2 = \frac{1}{3} \int_{W_D^\epsilon} \omega_1. \quad (1.15)$$

4. We show in §13 that the fundamental classes $[\overline{W}_D^\epsilon]$ are given by

$$[\overline{W}_D^\epsilon] = \frac{3}{4}[\omega_1] + \frac{9}{4}[\omega_2] + \pi_B[W_D^\epsilon] \pm j, \quad (1.16)$$

for some $j \in J$. The plan is to leave the coefficient of $[\omega_1]$ as an unknown a and then to use our knowledge of Y_D to write down equations involving the class $[\overline{W}_D^\epsilon]$ that allow us to solve for a . The following equations follow from Properties (b) and (c) of Y_D :

$$\begin{aligned} [\overline{W}_D^0] \cdot [\overline{P}_D] &= 0, \\ [\overline{W}_D^0] \cdot [\overline{W}_D^0] &= [\overline{W}_D^1] \cdot [\overline{W}_D^1], \end{aligned}$$

These together with (1.15) give us enough equations to solve for a . Theorem 1.3 follows from (1.16) by pairing $-\omega_1$ with both sides as in the proof of Theorem 1.1.

Notes and references. The orbits of the $\mathrm{SL}_2\mathbb{R}$ action on $\Omega_1\mathcal{M}_g$ project to immersions $\mathbb{H} \rightarrow \mathcal{M}_g$ of the hyperbolic plane into $\mathcal{M}_g$ which are isometric and totally geodesic with respect to the Teichmüller metric on $\mathcal{M}_g$. A totally geodesic immersion $\mathbb{H} \rightarrow \mathcal{M}_g$ is called a *Teichmüller disk*. It sometimes happens that a Teichmüller disk covers an algebraic curve C in $\mathcal{M}_g$. In that case, the normalization of C is called a *Teichmüller curve*. The curves W_D are examples of Teichmüller curves in genus two. This action of $\mathrm{SL}_2\mathbb{R}$ is also closely related to the study of billiards in rational angled polygons, as well as the study of interval exchange maps. For information about Teichmüller curves, the action of $\mathrm{SL}_2\mathbb{R}$ on $\Omega\mathcal{M}_g$, and its relation with rational billiards and interval exchange maps, see for example [KMS86], [Mas82], [MT02], [Vee86], [Vee89], [Vee90], [Vee92], [KZ], [Zor99], [Möl06], or [BM].

By analogy with the work of Ratner [Rat94] on the dynamics of actions of groups generated by unipotent elements on homogeneous spaces, it is conjectured that the closure of every Teichmüller disk in $\mathcal{M}_g$ is an algebraic sub-orbifold of $\mathcal{M}_g$. McMullen's work establishes this conjecture in genus two for Teichmüller disks generated by Abelian differentials. McMullen analyzed the dynamics of $\mathrm{SL}_2\mathbb{R}$ on $\Omega_1\mathcal{M}_2$ in the series of papers, [McM03a], [McM03b], [McMa], [McM05a], [McM05b], and [McM06]. In these papers, he classified the closures of $\mathrm{SL}_2\mathbb{R}$ orbits in $\Omega_1\mathcal{M}_2$, and classified Teichmüller curves in genus two. He also introduced the curves W_D in these papers and showed that they are all of the Teichmüller curves which are generated by an Abelian differential with a double zero. According to [McM03a], the Teichmüller curve W_D is primitive exactly when D is nonsquare. The form generated by identifying opposite sides of a regular decagon has two simple zeros and generates a primitive Teichmüller curve D_{10} in $\mathcal{M}_2$. In [McM06], McMullen showed that D_{10} and the W_D for D nonsquare are in fact all of the primitive Teichmüller curves in $\mathcal{M}_2$ generated by an Abelian differential.

Teichmüller curves in genus two were also studied by Calta in [Cal04], using the Kenyon-Smillie invariant introduced in [KS00].

Associated to every known $\mathrm{SL}_2\mathbb{R}$ orbit closure in $\Omega_1\mathcal{M}_g$ is a canonical finite, ergodic measure μ . The volume of μ is an interesting quantity, which according to Veech [Vee98] and Eskin-Masur [EM01] gives information about the dynamics of the geodesic flow of a generic Abelian differential $(X, \omega) \in S$ with respect to the canonical flat metric on X determined by ω . Volumes of orbit closures and their applications to dynamics are studied in [EO01], [EOP], [EMS03], [EMZ03], [HL06], [LR06], and [Lel06].

A closed $\mathrm{SL}_2\mathbb{R}$ -orbit S lies over a Teichmüller curve C . In this case, the measure μ descends to a multiple of the hyperbolic area measure on C . These means that calculating the volume of μ is equivalent to calculating $\chi(C)$. In genus two, the Euler characteristics of the Teichmüller curves W_{d^2} (as well as the volume of a related $\mathrm{SL}_2\mathbb{R}$ -invariant measure on X_{d^2}) were studied in [EMS03]. They established (1.2) by counting certain special Abelian differentials, called square-tiled surfaces. The connected components of W_D were classified in [HL06] when D is the square of a prime, and in [McM05a] for arbitrary D . Theorem 1.4 calculating $\chi(W_{d^2}^\epsilon)$ was established in [HL06] when d is prime and was conjectured for arbitrary d . In [LR06] Lelièvre and Royer established Theorem 1.4 independently by counting square-tiled surfaces using the theory of quasimodular forms.

With the calculation of $\chi(W_D)$, we now know the Euler characteristics of all of the Teichmüller curves in $\mathcal{M}_2$ which are generated by Abelian differentials with a double zero. The number of cusps of W_D was calculated by McMullen in [McM05a]. We can also calculate the number of elliptic points on W_D using known formulas for the numbers of elliptic points on X_D and P_D . Thus we can calculate the genus of W_D for any D . At this point we don't know the number of elliptic points on the components W_D^ϵ , so we can't calculate the genus of these components, but we would conjecture that the two components have the same number of elliptic points and genera if D is not square. The curve W_D is defined over $\mathbb{Q}$, so the Galois automorphism σ of $\mathbb{Q}(\sqrt{D})$ acts on the $\overline{\mathbb{Q}}$ points of W_D . An alternative approach to showing that $\chi(W_D^0) = \chi(W_D^1)$, as well as showing that the components have the same number of elliptic points and genera, would be to show that σ permutes the two components of W_D .

Acknowledgments. This paper is an expanded version of my Ph.D. thesis at Harvard. I would like to thank my advisor, Curt McMullen, for his invaluable help with every stage of this work. I would also like to thank Sabin Cautis, Alex Eskin, Florian Herzig, and Howie Masur for useful conversations, and I would like to thank Joe Harris and Yum-Tong Siu for reading an earlier version of this paper.

2 Abelian surfaces and real multiplication

We discuss in this section the necessary preliminaries involving Abelian surfaces. In §2.1, we discuss the Siegel modular varieties which parameterize Abelian varieties. In §2.2, we introduce real multiplication, and in §2.3, we discuss Hilbert modular surfaces.

2.1 Abelian surfaces

Abelian varieties. A *complex torus* is a quotient $A = V/\Lambda$, where Λ is a lattice in a finite dimensional complex vector space V . A *principal polarization* on A is a Hermitian form H on V such that $\text{Im } H$ takes integral values on $\Lambda \times \Lambda$, and the pairing

$$\text{Im } H : \Lambda \times \Lambda \rightarrow \mathbb{Z}$$

is unimodular. A *principally polarized Abelian variety* is a complex torus equipped with a principal polarization.

Siegel modular varieties. The Siegel upper half space is

$$\mathcal{H}_g = \{Z \in M_g(\mathbb{C}) : Z^t = Z, \text{Im } Z > 0\},$$

an open subset in the $g(g+1)/2$ dimensional space of symmetric matrices in $M_g(\mathbb{C})$. The group $\text{Sp}_{2g}\mathbb{R}$ acts on $\mathcal{H}_g$ as a group of biholomorphic transformations by

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \cdot Z = (\alpha Z + \beta)(\gamma Z + \delta)^{-1}.$$

Equip $\Lambda = \mathbb{Z}^{2g}$ with the usual symplectic form defined by the matrix,

$$\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}. \quad (2.1)$$

Following [BL04], we define for each $Z \in \mathcal{H}_g$ a principally polarized Abelian surface X_Z together with a symplectic isomorphism $\Lambda \rightarrow H_1(X_Z; \mathbb{Z})$. For $Z \in \mathcal{H}_g$, let

$$\Lambda_Z = (Z, I)\Lambda,$$

a lattice in $\mathbb{C}^g$. Let $X_Z = \mathbb{C}^g/\Lambda_Z$, and give X_Z the polarization coming from the Hermitian form on $\mathbb{C}^g$ defined by the matrix,

$$(\text{Im } Z)^{-1}.$$

It can be shown that two given points $Z, Z' \in \mathcal{H}_g$, the polarized Abelian surfaces X_Z and $X_{Z'}$ are isomorphic if and only if Z and Z' are equivalent under the action of $\text{Sp}_{2g}\mathbb{Z}$.

The Siegel modular variety is $\mathcal{A}_g = \mathcal{H}_g/\text{Sp}_{2g}\mathbb{Z}$. From the previous paragraph, we obtain:

Theorem 2.1. *The normal analytic space $\mathcal{A}_g$ is the moduli space of principally polarized Abelian varieties of dimension g .*

Satake compactification. Following [vdG88], we briefly describe the Satake compactification $\widehat{\mathcal{A}}_g$ of $\mathcal{A}_g$, introduced by Satake in [Sat56b].

The transformation,

$$Z \mapsto (1 + iZ)(1 - iZ)^{-1},$$

maps $\mathcal{H}_g$ biholomorphically onto the bounded symmetric domain,

$$\mathcal{D}_g = \{Z \in M_g(\mathbb{C}) : Z = Z^t, 1_g - Z\overline{Z} > 0\}.$$

For $r \leq n$, we map $\mathcal{D}_r$ into $\overline{\mathcal{D}}_g$ by

$$Z \mapsto \begin{pmatrix} Z & 0 \\ 0 & 1_{n-r} \end{pmatrix},$$

and let

$$\mathcal{D}_g^* = \bigcup_{0 \leq r \leq g} \bigcup_{h \in \mathrm{Sp}_{2g}\mathbb{Q}} h\mathcal{D}_r.$$

Satake defined a natural topology on $\mathcal{D}_g^*$ for which the action of $\mathrm{Sp}_{2g}\mathbb{Q}$ extends continuously to an action on $\mathcal{D}_g^*$. With this topology, the Satake showed that the quotient $\widehat{\mathcal{A}}_g = \mathcal{D}_g^*/\mathrm{Sp}_{2g}\mathbb{Z}$ is a compact, normal complex analytic space which contains one copy of $\mathcal{A}_r$ for each $r \leq g$. Baily showed in [Bai58] that $\widehat{\mathcal{A}}_g$ has the structure of a normal projective variety.

2.2 Real multiplication

Quadratic orders. Let K be a quadratic field or $\mathbb{Q} \oplus \mathbb{Q}$. A *quadratic order* is a subring $\mathcal{O}$ of K such that $1 \in \mathcal{O}$ and $\mathcal{O} \otimes \mathbb{Q} = K$. Any quadratic order is isomorphic to one of

$$\mathcal{O}_D = \mathbb{Z}[T]/(T^2 + bT + c),$$

where $b, c \in \mathbb{Z}$ and $b^2 - 4c = D$. The isomorphism class of $\mathcal{O}_D$ only depends on D , so this defines a unique quadratic order for every nonzero integer $D \equiv 0$ or $1 \pmod{4}$. This integer D is called the *discriminant* of $\mathcal{O}_D$, and D is a *fundamental discriminant* if D is not of the form f^2E for some integers f and E with $E \equiv 0$ or $1 \pmod{4}$ and $f > 1$.

If D is not square, then $\mathcal{O}_D$ is a subring of $\mathbb{Q}(\sqrt{D})$, and $\mathcal{O}_D$ is the ring of integers of $\mathbb{Q}(\sqrt{D})$ if and only if D is a fundamental discriminant; otherwise, $\mathcal{O}_D$ is a subring of the ring of integers. If D is square, then $\mathcal{O}_D$ is a subring of $\mathbb{Q} \oplus \mathbb{Q}$. In that case,

$$\mathcal{O}_{d^2} = \{(x, y) \in \mathbb{Z} \times \mathbb{Z} : x \equiv y \pmod{d}\}. \quad (2.2)$$

We will regard $\mathbb{Q} \oplus \mathbb{Q}$ as an extension of $\mathbb{Q}$ by the diagonal map $\mathbb{Q} \rightarrow \mathbb{Q} \oplus \mathbb{Q}$. The Galois automorphism of $\mathbb{Q} \oplus \mathbb{Q}$ is $(x, y)' = (y, x)$, and we can use this to define norm and trace on $\mathbb{Q} \oplus \mathbb{Q}$ as for a field.

For the rest of this paper, fix two nonzero homomorphisms $\iota_i : \mathcal{O}_D \rightarrow \mathbb{R}$. If D is not square, then ι_i is an embedding, and if D is square, then ι_i is induced

by one of the two projections $\mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z}$. We will often abuse terminology and call ι_i an embedding even when D is square. We will also use the notation $\lambda^{(i)} = \iota_i(\lambda)$.

Inverse different. Given a quadratic order $\mathcal{O}_D$, the inverse different is the fractional ideal,

$$\mathcal{O}_D^\vee = \{x \in K_D : \text{Tr}(xy) \in \mathbb{Z}, \forall y \in \mathcal{O}_D\}.$$

More concretely,

$$\mathcal{O}_D^\vee = \frac{1}{\sqrt{D}} \mathcal{O}_D,$$

where if $D = d^2$, we interpret $\sqrt{D}$ to be $(d, -d)$.

We equip $\mathcal{O}_D \oplus \mathcal{O}_D^\vee$ with the unimodular symplectic pairing,

$$\langle (x_1, y_1), (x_2, y_2) \rangle = \text{Tr}(x_1 y_2 - x_2 y_1).$$

Real multiplication. Consider an Abelian surface $A = V/\Lambda$. *Real multiplication by $\mathcal{O}_D$* on A is a monomorphism $\rho: \mathcal{O}_D \rightarrow \text{End}(A)$ (where $\text{End}(A)$ is the ring of holomorphic endomorphisms of A) with the following properties:

- For each $\lambda \in \mathcal{O}_D$, the lift $\tilde{\rho}(\lambda): V \rightarrow V$ is self-adjoint with respect to the Hermitian form on V given by the polarization of A .
- ρ is *proper* in the sense that it doesn't extend to a monomorphism

$$\rho': \mathcal{O}_E \rightarrow \text{End}(A)$$

for some $\mathcal{O}_E \supset \mathcal{O}_D$ – that is, for some $\mathcal{O}_E$ with $E = f^2 D$ for some $f > 1$.

Let $\Omega(A) \cong V^*$ be the space of holomorphic one-forms on A . Since the real multiplication is self-adjoint, there is an eigenspace decomposition

$$\Omega(A) = \Omega^1(A) \oplus \Omega^2(A).$$

We order the eigenspaces so that $\rho(\lambda) \cdot \omega = \lambda^{(i)} \omega$ for $\omega \in \Omega^i(A)$.

2.3 Hilbert modular surfaces

For any quadratic discriminant D , let $K_D = \mathcal{O}_D \otimes \mathbb{Q}$, which is $\mathbb{Q}(\sqrt{D})$ if D is not square and is $\mathbb{Q} \oplus \mathbb{Q}$ if D is square. Define the group,

$$\text{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2 K_D : a \in \mathcal{O}_D, b \in (\mathcal{O}_D^\vee)^{-1}, c \in \mathcal{O}_D^\vee, d \in \mathcal{O}_D \right\}.$$

$\text{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ has two embeddings in $\text{SL}_2 \mathbb{R}$ induced by the two embeddings $\iota_i: K_D \rightarrow \mathbb{R}$.

Definition. The *Hilbert modular surface* of discriminant D is the quotient,

$$X_D = \mathbb{H} \times \mathbb{H} / \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee),$$

where $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ acts on $\mathbb{H} \times \mathbb{H}$ by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (z_1, z_2) = \left(\frac{a^{(1)}z_1 + b^{(1)}}{c^{(1)}z_1 + d^{(1)}}, \frac{a^{(2)}z_2 + b^{(2)}}{c^{(2)}z_2 + d^{(2)}} \right).$$

There is an isomorphism $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee) \rightarrow \mathrm{SL}_2\mathcal{O}_D$ defined by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} a & \frac{1}{\sqrt{D}}b \\ \sqrt{D}c & d \end{pmatrix},$$

and the map $T: \mathbb{H} \times \mathbb{H} \rightarrow \mathbb{H} \times (-\mathbb{H})$ induces an isomorphism,

$$X_D \cong \mathbb{H} \times (-\mathbb{H}) / \mathrm{SL}_2\mathcal{O}_D.$$

Theorem 2.2. *The Hilbert modular surface X_D is the moduli space of all pairs (A, ρ) , where A is a principally polarized Abelian surface, and $\rho: \mathcal{O}_D \rightarrow \mathrm{End}(A)$ is a choice of real multiplication on A .*

Sketch of proof (following [McMb]). Given $\tau = (\tau_1, \tau_2) \in \mathbb{H} \times \mathbb{H}$, let

$$\phi_\tau: \mathcal{O}_D \oplus \mathcal{O}_D^\vee$$

be the embedding,

$$\phi_\tau(x, y) = (x^{(1)} + y^{(1)}\tau_1, x^{(2)} + y^{(2)}\tau_2).$$

Let $A_\tau = \mathbb{C}^2 / \phi_\tau(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ with the principal polarization induced by the symplectic pairing on $\mathcal{O}_D \oplus \mathcal{O}_D^\vee$. This polarization is also given by the Hermitian form,

$$H_\tau(z, w) = \frac{1}{\mathrm{Im} \tau_1} z_1 \bar{w}_1 + \frac{1}{\mathrm{Im} \tau_2} z_2 \bar{w}_2. \quad (2.3)$$

Define real multiplication on A_τ by

$$\lambda \cdot (z_1, z_2) = (\lambda^{(1)}z_1, \lambda^{(2)}z_2).$$

We thus get a map from $\mathbb{H} \times \mathbb{H}$ to the set of all triples (A, ρ, ϕ) , where (A, ρ) is a principally polarized Abelian surface with real multiplication by $\mathcal{O}_D$, and ϕ is a choice of an $\mathcal{O}_D$ -linear, symplectic isomorphism $\phi: \mathcal{O}_D \oplus \mathcal{O}_D^\vee \rightarrow H_1(A; \mathbb{Z})$.

Given

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2 K_D,$$

let

$$g^* = \begin{pmatrix} a & -b \\ -c & d \end{pmatrix},$$

and define

$$\chi(g, \tau) = \begin{pmatrix} (c^{(1)}\tau_1 + d^{(1)})^{-1} & 0 \\ 0 & (c^{(2)}\tau_2 + d^{(2)})^{-1} \end{pmatrix}. \quad (2.4)$$

We have the following commutative diagram.

$$\begin{array}{ccc} \mathcal{O}_D \oplus \mathcal{O}_D^\vee & \xrightarrow{\phi_\tau} & \mathbb{C}^2 \\ g^* \downarrow & & \downarrow \chi(g, \tau) \\ \mathcal{O}_D \oplus \mathcal{O}_D^\vee & \xrightarrow{\phi_{g \cdot \tau}} & \mathbb{C}^2 \end{array}$$

Thus $\chi(g, \tau)$ induces an isomorphism between A_τ and $A_{g \cdot \tau}$ which preserves the polarizations and commutes with the action of real multiplication. We then get a map from X_D to the set of all principally polarized Abelian surfaces with a choice of real multiplication, which can be shown to be a bijection. $\blacksquare$

Replacing each pair (A, ρ) with (A, ρ') , where ρ' is the composition of ρ with the Galois automorphism of $\mathcal{O}_D$, induces an involution τ of X_D . The lift of τ to the universal cover $\mathbb{H} \times \mathbb{H}$ of X_D is given by $\tilde{\tau}(z_1, z_2) = (z_2, z_1)$.

There is a natural map $j: X_D \rightarrow \mathcal{A}_2$ which forgets the choice of real multiplication. This map j is generically two to one and is equivariant with respect to τ .

Baily-Borel compactification. We can regard the boundary $\partial\mathbb{H} \subset \mathbb{P}^1(\mathbb{C})$ as $\mathbb{P}^1(\mathbb{R}) \cong \mathbb{R} \cup \{\infty\}$. Given a real quadratic field $K \subset \mathbb{R}$, define an embedding $\mathbb{P}^1(K) \rightarrow \mathbb{P}^1(\mathbb{R})^2$ by

$$[x : y] \mapsto ([x : y], [x' : y']).$$

When D is not square, define via this embedding

$$(\mathbb{H} \times \mathbb{H})_D = (\mathbb{H} \times \mathbb{H}) \cup \mathbb{P}^1(K_D).$$

When D is square, define

$$(\mathbb{H} \times \mathbb{H})_D = (\mathbb{H} \cup \mathbb{P}^1(\mathbb{Q})) \times (\mathbb{H} \cup \mathbb{P}^1(\mathbb{Q})).$$

We give $\mathbb{H} \cup \mathbb{P}^1(\mathbb{Q})$ the usual topology where if $r \in \mathbb{P}^1(\mathbb{Q})$, then a basis of open neighborhoods of r is given by sets of the form $U \cup \{r\}$, where $U \subset \mathbb{H}$ is an open horoball resting on r . We then give $(\mathbb{H} \times \mathbb{H})_{d^2}$ the product topology. When D is not square, there is a similar natural topology on $(\mathbb{H} \times \mathbb{H})_D$, which is described in [vdG88].

The action of $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ on $\mathbb{H} \times \mathbb{H}$ extends continuously to $(\mathbb{H} \times \mathbb{H})_D$. The quotient,

$$\widehat{X}_D = (\mathbb{H} \times \mathbb{H})_D / \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee),$$

is compact and Hausdorff. The space $\widehat{X}_D$ is the Baily-Borel compactification of X_D .

Theorem 2.3 ([Bai58]). *The compactification $\widehat{X}_D$ is a normal, projective, algebraic variety.*

When D is not square, $\widehat{X}_D \setminus X_D$ consists of finitely many points, which we will call the cusps of X_D . When $D = d^2$, the image of $\mathbb{P}^1(\mathbb{Q}) \times \mathbb{P}^1(\mathbb{Q})$ in $\widehat{X}_{d^2}$ also consists of finitely many points which we will call the cusps of $\widehat{X}_{d^2}$. In $\widehat{X}_{d^2}$, let

$$R_{d^2}^1 = \pi \left(\bigcup_{r \in \mathbb{P}^1(\mathbb{Q})} \mathbb{H} \times \{r\} \right),$$

and let

$$R_{d^2}^2 = \pi \left(\bigcup_{r \in \mathbb{P}^1(\mathbb{Q})} \{r\} \times \mathbb{H} \right),$$

where $\pi: (\mathbb{H} \times \mathbb{H})_D \rightarrow \widehat{X}_D$ is the natural quotient map. Then we have a disjoint union,

$$\widehat{X}_{d^2} = X_{d^2} \amalg R_{d^2}^1 \amalg R_{d^2}^2 \amalg C,$$

where C is the set of cusps of $\widehat{X}_{d^2}$.

For $d \in \mathbb{N}$, define

$$\Gamma_1(d) = \left\{ \begin{pmatrix} a & b \\ c & e \end{pmatrix} \in \mathrm{SL}_2\mathbb{Z} : a \equiv e \equiv 1 \pmod{d}, \text{ and } c \equiv 0 \pmod{d} \right\}.$$

Proposition 2.4. *The curve $R_{d^2}^i$ is irreducible, and*

$$R_{d^2}^i \cong \mathbb{H}/\Gamma_1(d).$$

Proof. By (2.2), we can regard $\mathrm{SL}_2\mathcal{O}_{d^2}$ as

$$\{(A, B) \in (\mathrm{SL}_2\mathbb{Z})^2 : A \equiv B \pmod{d}\}. \quad (2.5)$$

Since $\mathrm{SL}_2\mathbb{Z}$ acts transitively on $\mathbb{P}^1(\mathbb{Q})$, we have

$$R_{d^2}^1 = \mathbb{H} \times \{\infty\} / \mathrm{Stab}_{\mathbb{H} \times \{\infty\}}.$$

The matrices A in the pair $(A, B) \in \mathrm{Stab}_{\mathbb{H} \times \{\infty\}}$ are exactly the matrices which are congruent to an upper triangular matrix mod d . Thus $R_{d^2}^1$ is as claimed. ■

When D is not square, the cusps of $\widehat{X}_D$ are complicated singularities; however, when D is square, they are just orbifold singularities.

Proposition 2.5. *The Baily-Borel compactification $\widehat{X}_{d^2}$ is a compact, complex orbifold with singularities at the elliptic points of X_{d^2} , the elliptic points of $R_{d^2}^i$, and possibly the cusps of $\widehat{X}_{d^2}$.*

Proof. Consider the principal congruence subgroup,

$$\Gamma(d) = \{A \in \mathrm{SL}_2\mathbb{Z} : A \equiv I \pmod{d}\}.$$

The product,

$$\Gamma(d) \times \Gamma(d) \subset \mathrm{SL}_2\mathcal{O}_{d^2},$$

is a finite index, normal subgroup. We have,

$$(\mathbb{H} \times \mathbb{H})_{d^2}/(\Gamma(d) \times \Gamma(d)) = \overline{\mathbb{H}/\Gamma(d)} \times \overline{\mathbb{H}/\Gamma(d)},$$

which is a manifold, so

$$\widehat{X}_{d^2} = (\overline{\mathbb{H}/\Gamma(d)} \times \overline{\mathbb{H}/\Gamma(d)})/G,$$

where $G = \mathrm{SL}_2\mathcal{O}_{d^2}/(\Gamma(d) \times \Gamma(d))$, a finite group. Thus $\widehat{X}_{d^2}$ is compact orbifold with singularities at the fixed points of G . $\blacksquare$

The involution $(z_1, z_2) \mapsto (z_2, z_1)$ of $\mathbb{H} \times \mathbb{H}$ extends continuously to $(\mathbb{H} \times \mathbb{H})_D$, so the involution τ extends to an involution τ of $\widehat{X}_D$, which preserves the cusps of $\widehat{X}_D$. If $D = d^2$, then we also have $\tau(R_{d^2}^1) = R_{d^2}^2$.

Theorem 2.6. *The natural map $j: X_D \rightarrow \mathcal{A}_2$ extends to a finite morphism $j: \widehat{X}_D \rightarrow \widehat{\mathcal{A}}_2$ which sends the cusps of $\widehat{X}_D$ to the point $\mathcal{A}_0 \in \mathcal{A}_2$ and sends the curves R_D^i to $\mathcal{A}_1$.*

This allows us to define $\widehat{X}_D$ alternatively in terms of the image of X_D in $\widehat{\mathcal{A}}_2$. Recall the notion of the normalization of a variety in a finite algebraic extension of its function field discussed in §A.

Corollary 2.7. *The Baily-Borel compactification $\widehat{X}_D$ is the normalization of the closure of $j(X_D)$ in $\mathcal{A}_2$ in the field $K(X_D)$, the function field of X_D .*

Line bundles on X_D . We now discuss some bundles on X_D , which will be used in §9. For $i = 1$ or 2 , define an action of $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ on $(\mathbb{H} \times \mathbb{H}) \times \mathbb{C}$ by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (z_1, z_2, w) = \left(\frac{a^{(1)}z_1 + b^{(1)}}{c^{(1)}z_1 + d^{(1)}}, \frac{a^{(2)}z_2 + b^{(2)}}{c^{(2)}z_2 + d^{(2)}}, (c^{(i)}z_i + d^{(i)})^2 w \right).$$

The quotient is a line bundle over X_D , which we call L_i .

We define a Hermitian metric $\tilde{h}_i$ on $(\mathbb{H} \times \mathbb{H}) \times \mathbb{C}$ by defining on the fiber over (z_1, z_2) ,

$$\tilde{h}_i(w, w) = y_i^2 |w|^2.$$

The metric $\tilde{h}_i$ is $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ -invariant, so it descends to a Hermitian metric h_i on L_i . The Chern form of h_i is,

$$\begin{aligned} c_1(L_i, h_i) &= -\frac{i}{\pi} \partial \bar{\partial} \log(y_i) \\ &= \frac{1}{2\pi} \frac{dx_i \wedge dy_i}{y_i^2}, \end{aligned} \tag{2.6}$$

so the Chern form of h_i is $c_1(L_i, h_i) = \omega_i$, where ω_i is the 2-form on X_D covered by the form (2.6) on $\mathbb{H} \times \mathbb{H}$.

Define line bundles,

$$\Omega^i X_D = \{(A, \omega) : A \in X_D \text{ and } \omega \in \Omega^i(A)\},$$

and

$$Q^i X_D = (\Omega^i X_D)^{\otimes 2},$$

for $i = 1$ or 2 . We will mostly use these bundles when $i = 1$, so we will write ΩX_D or $Q X_D$ for $\Omega^1 X_D$ or $Q^1 X_D$.

For $A \in X_D$, the polarization defines a Hermitian metric on $\Omega(A)$, which restricts to a Hermitian metric on $\Omega^i(A)$. Put this Hermitian metric on each $\Omega^i(A)$ to define a Hermitian metric h_Ω^i on $\Omega^i X_D$. Let h_Q^i be the induced metric on $Q^i X_D$.

Proposition 2.8. *There is an isomorphism $L_i \rightarrow Q^i X_D$ which preserves the Hermitian metrics on these bundles. Thus,*

$$c_1(Q^i X_D, h_Q^i) = \omega_i.$$

Proof. Define a map $f: (\mathbb{H} \times \mathbb{H}) \times \mathbb{C} \rightarrow Q^i X_D$ by

$$(\tau, w) \mapsto (A_\tau, w dz_i^2),$$

where $A_\tau = \mathbb{C}^2 / \phi_\tau(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ is as in the proof of Theorem 2.2. With $\chi(g, \tau)$ as in (2.4), we have

$$\chi(g, \tau)_* dz_i^2 = (c^{(i)} \tau_i + d^{(i)})^2 dz_i^2.$$

This means that f descends to an isomorphism $L_i \rightarrow Q^i X_D$ as claimed. The fact that this isomorphism preserves the Hermitian metrics follows directly from the formula (2.3) for the polarization. $\blacksquare$

The product foliation of $\mathbb{H} \times \mathbb{H}$ with leaves of the form $\{c\} \times \mathbb{H}$ is invariant under $\text{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$. Let $\mathcal{A}_D$ be the induced foliation of X_D , and let

$$T^* \mathcal{A}_D \rightarrow X_D$$

be the line bundle whose fiber over a point p is the cotangent bundle to the leaf of $\mathcal{A}_D$ through p . We give the leaves of $\mathcal{A}_D$ their hyperbolic metric, and give $T^* \mathcal{A}_D$ the induced Hermitian metric $h_{\mathcal{A}_D}$. The following is easy to check.

Proposition 2.9. *There is an isomorphism $L_2 \rightarrow T^* \mathcal{A}_D$ preserving the Hermitian metrics. Thus,*

$$c_1(T^* \mathcal{A}_D, h_{\mathcal{A}_D}) = \omega_2.$$

The following relation between ΩX_D and $T^* \mathcal{A}_D$ will be used in §10.

Proposition 2.10. *For any $p \in X_D$, there is a neighborhood U of p in the leaf L of $\mathcal{A}_D$ through p and a section ω of ΩX_D over U such that the periods of the forms $\omega(z)$ are constant over U .*

Proof. Let

$$(\tau_1, \tau_2) \in \tilde{X}_D = \mathbb{H} \times \mathbb{H}$$

lie over p in the universal cover of X_D . Define a section s of $\Omega \tilde{X}_D$ by

$$s(\tau_1, \tau_2) = dz_1 \in \Omega^1(A_{(\tau_1, \tau_2)}),$$

with A_τ as in the proof of Theorem 2.2. By the definition of A_τ , the periods of $s(\tau_1, \tau_2)$ only depend on τ_1 , so are constant along $\mathcal{A}_D$. ■

Euler characteristic of X_D . We now discuss the Euler characteristic $\chi(X_D)$, which by the generalized Gauss-Bonnet theorem is given by the volume,

$$\chi(X_D) = \int_{X_D} \omega_1 \wedge \omega_2,$$

where ω_1 are the 2-forms on X_D defined by (2.6). The volume of X_D was calculated by Siegel when D is a fundamental discriminant.

For any number field K , the *Dedekind zeta-function* of K is defined by

$$\zeta_K(s) = \sum_{\mathfrak{a}} \frac{1}{N_{\mathbb{Q}}^K(\mathfrak{a})^s},$$

where the sum is over all nonzero ideals $\mathfrak{a} \subset \mathcal{O}$ with $\mathcal{O}$ the ring of integers in K . If $K = \mathbb{Q}$, then ζ_K is just the Riemann zeta-function. The definition of ζ_K also makes sense if $K = \mathbb{Q} \oplus \mathbb{Q}$. In that case,

$$\zeta_{\mathbb{Q} \oplus \mathbb{Q}}(s) = \zeta_{\mathbb{Q}}(s)^2.$$

ζ_K can be analytically continued to a meromorphic function on $\mathbb{C}$ which has a simple pole at $s = 1$.

Theorem 2.11 ([Sie36]). *When D is a fundamental discriminant, and $D \neq 1$,*

$$\chi(X_D) = 2\zeta_{K_D}(-1).$$

We will also want to know $\chi(X_D)$ when D is not a fundamental discriminant. For integers a and b with $b > 0$, let

$$\left(\frac{a}{b}\right)$$

be the Kronecker symbol, defined in [Miy89, p. 82].

Theorem 2.12. *If D is a fundamental discriminant, and $f \in \mathbb{N}$, then*

$$\chi(X_{f^2D}) = \begin{cases} 1/36 & \text{if } D = f = 1; \\ 1/6 & \text{if } D = 1 \text{ and } f = 2. \end{cases}$$

Otherwise,

$$\chi(X_{f^2D}) = 2f^3 \zeta_{K_D}(-1) \sum_{r|f} \left(\frac{D}{r} \right) \frac{\mu(r)}{r^2}. \quad (2.7)$$

Remark. When $D = 1$, (2.7) reduces to

$$\chi(X_{d^2}) = \frac{1}{72} d^3 \sum_{r|d} \frac{\mu(r)}{r^2}.$$

We will derive Theorem 2.12 from Theorem 2.11 in a sequence of lemmas.

Lemma 2.13. *We have,*

$$[\mathrm{PSL}_2\mathbb{Z} \times \mathrm{PSL}_2\mathbb{Z} : \mathrm{PSL}_2\mathcal{O}_4] = 6,$$

and

$$[\mathrm{PSL}_2\mathbb{Z} \times \mathrm{PSL}_2\mathbb{Z} : \mathrm{PSL}_2\mathcal{O}_{d^2}] = \frac{1}{2} d^3 \sum_{r|d} \frac{\mu(r)}{r^2}$$

if $d > 2$.

Proof. We will regard $\mathrm{SL}_2\mathcal{O}_{d^2}$ as the subgroup of $\mathrm{SL}_2\mathbb{Z} \times \mathrm{SL}_2\mathbb{Z}$ given in (2.5). The natural map,

$$\mathrm{SL}_2\mathbb{Z} \rightarrow \mathrm{SL}_2(\mathbb{Z}/d),$$

is surjective by [Miy89, Theorem 4.2.1] and has kernel the principal congruence subgroup $\Gamma(d)$, so we have the exact sequence,

$$0 \rightarrow \Gamma(d) \times \Gamma(d) \rightarrow \mathrm{SL}_2\mathcal{O}_{d^2} \rightarrow \mathrm{SL}_2(\mathbb{Z}/d) \rightarrow 0.$$

Thus,

$$[\mathrm{SL}_2\mathcal{O}_{d^2} : \Gamma(d) \times \Gamma(d)] = |\mathrm{SL}_2(\mathbb{Z}/d)|,$$

and

$$\begin{aligned} [\mathrm{SL}_2\mathbb{Z} \times \mathrm{SL}_2\mathbb{Z} : \mathrm{SL}_2\mathcal{O}_{d^2}] &= \frac{[\mathrm{SL}_2\mathbb{Z} \times \mathrm{SL}_2\mathbb{Z} : \Gamma(d) \times \Gamma(d)]}{[\mathrm{SL}_2\mathcal{O}_{d^2} : \Gamma(d) \times \Gamma(d)]} \\ &= |\mathrm{SL}_2(\mathbb{Z}/d)|. \end{aligned} \quad (2.8)$$

By [Miy89, Theorem 4.2.4],

$$|\mathrm{SL}_2(\mathbb{Z}/d)| = d^3 \sum_{r|d} \frac{\mu(r)}{r^2}.$$

The kernel of $\mathrm{SL}_2\mathbb{Z} \times \mathrm{SL}_2\mathbb{Z} \rightarrow \mathrm{PSL}_2\mathbb{Z} \times \mathrm{PSL}_2\mathbb{Z}$ has order 4, and the kernel of $\mathrm{SL}_2\mathcal{O}_{d^2} \rightarrow \mathrm{PSL}_2\mathcal{O}_{d^2}$ has order 4 if $d = 2$, and order 2 if $d > 2$. The desired formulas follow from this and (2.8). $\blacksquare$

Lemma 2.14. *For any fundamental discriminant $D \neq 1$, we have,*

$$\begin{aligned} [\mathrm{PSL}_2 \mathcal{O}_D : \mathrm{PSL}_2 \mathcal{O}_{f^2 D}] &= f^3 \prod_{p|f} \left(1 - \left(\frac{D}{p} \right) p^{-2} \right) \\ &= f^3 \sum_{r|f} \left(\frac{D}{r} \right) \frac{\mu(r)}{r^2}. \end{aligned}$$

Proof. Since the map $\mathrm{SL}_2 \mathcal{O}_{f^2 D} \rightarrow \mathrm{PSL}_2 \mathcal{O}_{f^2 D}$ has kernel $\pm I$, we have

$$[\mathrm{PSL}_2 \mathcal{O}_D : \mathrm{PSL}_2 \mathcal{O}_{f^2 D}] = [\mathrm{SL}_2 \mathcal{O}_D : \mathrm{SL}_2 \mathcal{O}_{f^2 D}].$$

The natural map,

$$\mathrm{SL}_2 \mathcal{O}_D / \mathrm{SL}_2 \mathcal{O}_{f^2 D} \rightarrow \prod_{\substack{p|f \\ p \text{ prime}}} \mathrm{SL}_2(\mathcal{O}_D/p) / \mathrm{SL}_2(\mathbb{Z}/p),$$

is a bijection. Thus, we need only to show that

$$[\mathrm{SL}_2(\mathcal{O}_D/p) : \mathrm{SL}_2 \mathbb{Z}] = p^3 - \left(\frac{D}{p} \right) p$$

for every prime $p|f$.

First, suppose that $(D/p) = -1$. This means that p remains prime in $\mathcal{O}_D$, so $\mathcal{O}_D/p \cong \mathbb{F}_{p^2}$, where we write $\mathbb{F}_q$ for the unique finite field of order q . It is easy to show that

$$|\mathrm{SL}_2 \mathbb{F}_q| = q^3 - q,$$

for any q . Thus

$$[\mathrm{SL}_2 \mathbb{F}_{p^2} : \mathrm{SL}_2 \mathbb{F}_p] = p^3 + p$$

as desired.

Now suppose that $(D/p) = 1$. This means that p splits in $\mathcal{O}_D$, so $\mathcal{O}_D/p \cong \mathbb{F}_p \oplus \mathbb{F}_p$. Therefore,

$$\begin{aligned} [\mathrm{SL}_2(\mathcal{O}_D/p) : \mathrm{SL}_2(\mathbb{Z}/p)] &= [\mathrm{SL}_2 \mathbb{F}_p \times \mathrm{SL}_2 \mathbb{F}_p : \mathrm{SL}_2 \mathbb{F}_p] \\ &= |\mathrm{SL}_2 \mathbb{F}_p| \\ &= p^3 - p. \end{aligned}$$

Finally, suppose that $(D/p) = 0$. This means that p ramifies in $\mathcal{O}_D$, so $\mathcal{O}_D/p \cong \mathbb{F}_p(\epsilon)/(\epsilon^2)$. We have

$$\begin{pmatrix} a + a'\epsilon & b + b'\epsilon \\ c + c'\epsilon & d + d'\epsilon \end{pmatrix} \in \mathrm{SL}_2(\mathbb{F}_p(\epsilon)/(\epsilon^2))$$

if and only if

$$ad - bc = 1, \quad \text{and} \tag{2.9}$$

$$da' + ad' - cb' - bc' = 0. \tag{2.10}$$

For any (a, b, c, d) satisfying (2.9), there are p^3 solutions to (2.10). Thus,

$$[\mathrm{SL}_2(\mathbb{F}_p(\epsilon)/(\epsilon^2)) : \mathrm{SL}_2\mathbb{F}_p] = p^3.$$

■

Proof of Theorem 2.12. Since the natural map $X_{f^2D} \rightarrow X_D$ is an orbifold covering map, we have

$$\chi(X_{f^2D}) = \chi(X_D)[\mathrm{PSL}_2\mathcal{O}_D : \mathrm{PSL}_2\mathcal{O}_{f^2D}],$$

so the claim follows from Lemma 2.13, Lemma 2.14, and Theorem 2.11 together with

$$\chi(X_1) = \frac{1}{36}$$

because $X_1 \cong \mathbb{H}/\mathrm{SL}_2\mathbb{Z} \times \mathbb{H}/\mathrm{SL}_2\mathbb{Z}$, and

$$\chi(\mathbb{H}/\mathrm{SL}_2\mathbb{Z}) = -\frac{1}{6}.$$

■

Modular forms. For any fundamental discriminant $D > 0$, define

$$H(2, f^2D) = -12\zeta_{K_D}(-1) \sum_{r|f} \mu(r) \left(\frac{D}{r}\right) r\sigma_3\left(\frac{f}{r}\right),$$

where

$$\sigma_m(n) = \sum_{d|n} d^m,$$

and we also adopt the convention that

$$\sigma_m(0) = \frac{1}{2}\zeta_{\mathbb{Q}}(-m),$$

and $\sigma_m(n) = 0$ if $n < 0$. Also define

$$H(2, 0) = \zeta_{\mathbb{Q}}(-3) = \frac{1}{120}.$$

The function $\mathcal{H}: \mathbb{H} \rightarrow \mathbb{C}$,

$$\begin{aligned} \mathcal{H}(\tau) &= \sum_{\substack{D \equiv 0,1 \pmod{4} \\ D \geq 0}} H(2, D)q^D, \quad (\text{with } q(\tau) = e^{2\pi i\tau}) \\ &= -\frac{1}{120} - \frac{1}{12}q - \frac{7}{12}q^4 - \frac{2}{5}q^5 - q^8 - \frac{25}{12}q^9 - 2q^{12} - 2q^{13} - \frac{55}{12}q^{16} - \dots, \end{aligned}$$

was shown by Cohen in [Coh75], to be a modular form of weight $5/2$ for the group $\Gamma_0(4)$. See the discussion in Chapter IX of [vdG88] for more about this form and its relation to X_D .

It is an elementary argument using Möbius inversion to show that for any fundamental discriminant D and $f \in \mathbb{N}$,

$$H(2, f^2 D) = -12 \sum_{s|f} \left(\zeta_{K_D}(-1) s^3 \sum_{r|s} \left(\frac{D}{r} \right) \frac{\mu(r)}{r^2} \right). \quad (2.11)$$

From this and Theorem 2.12, we obtain:

Theorem 2.15. *If $D \neq 1$ is a fundamental discriminant, then*

$$\sum_{r|f} \chi(X_{r^2 D}) = -\frac{1}{6} H(2, f^2 D).$$

Siegel and Cohen gave the following simple formula for $H(2, D)$.

Theorem 2.16 ([Sie69, Coh75]). *If D is not square, then*

$$H(2, D) = -\frac{1}{5} \sum_{e \equiv D(2)} \sigma_1 \left(\frac{D - e^2}{4} \right).$$

When D is square,

$$H(2, D) = -\frac{1}{5} \sum_{e \equiv D(2)} \sigma_1 \left(\frac{D - e^2}{4} \right) - \frac{D}{10}.$$

We will obtain an alternative proof of these formulas in §11.

The product locus. In X_D , define the *product locus* P_D to be the set of all $A \in X_D$ such that A is a polarized product of elliptic curves. The locus P_D was studied in [McM05a] and [McMb]. The locus P_D is also a union of *modular curves*, which were studied in [Hir73] and [vdG88].

Let $\tilde{P}_D \subset \mathbb{H} \times \mathbb{H}$ be the inverse image of P_D in X_D . The locus P_D is a linear subvariety of X_D in the following sense.

Proposition 2.17 ([McM05a]). *$\tilde{P}_D$ is a countable union of graphs of Möbius transformations $\mathbb{H} \rightarrow \mathbb{H}$.*

The goal of the rest of this section is to calculate $\chi(P_D)$. We will work with an auxiliary covering space Q_D of P_D defined as follows.

Let ΩQ be the space of elliptic curves equipped with Abelian differentials; that is,

$$\Omega Q = \Omega \mathcal{A}_1 \times \Omega \mathcal{A}_1 = \mathrm{GL}_2^+ \mathbb{R} / \mathrm{SL}_2 \mathbb{Z} \times \mathrm{GL}_2^+ \mathbb{R} / \mathrm{SL}_2 \mathbb{Z}, \quad (2.12)$$

where $\mathrm{GL}_2^+ \mathbb{R}$ is the subgroup of $\mathrm{GL}_2 \mathbb{R}$ consisting of matrices of positive determinant. In ΩQ , let ΩQ_D be the locus of pairs $((E_1, \omega_1), (E_2, \omega_2))$ such that the product

$$(E_1, \omega_1) \oplus (E_2, \omega_2) := (E_1 \times E_2, \omega_1 + \omega_2) \quad (2.13)$$

is an eigenform for real multiplication by $\mathcal{O}_D$. Let

$$Q_D = \mathbb{P}\Omega Q_D.$$

We now describe the connected components of ΩQ_D following [McM05a]. A *prototype* for real multiplication by $\mathcal{O}_D$ is a triple of integers (e, l, m) such that

$$D = e^2 + 4l^2m, \quad l, m > 0, \quad \text{and } \gcd(e, l) = 1.$$

To each such prototype, we associate a *prototypical eigenform* $\mathcal{Q}(e, l, m)$ as follows. Let λ be the unique positive root of $\lambda^2 - e\lambda - l^2m = 0$, and consider the lattices,

$$\begin{aligned} \Lambda_1 &= \mathbb{Z}(\lambda, 0) \oplus \mathbb{Z}(0, \lambda) \\ \Lambda_2 &= \mathbb{Z}(lm, 0) \oplus \mathbb{Z}(0, l). \end{aligned} \tag{2.14}$$

Let $E_i = \mathbb{C}/\Lambda_i$, and equip E_i with the form ω_i covered by dz on $\mathbb{C}$. Let

$$\mathcal{Q}(e, l, m) = (E_1, \omega_1) \oplus (E_2, \omega_2).$$

From the description of ΩQ in (2.12), we obtain an action of $\mathrm{GL}_2^+\mathbb{R}$ on ΩQ induced by the diagonal action of $\mathrm{GL}_2^+\mathbb{R}$ on $\mathrm{GL}_2^+\mathbb{R} \times \mathrm{GL}_2^+\mathbb{R}$.

Theorem 2.18 ([McM05a, Theorem 2.1]). *The locus ΩQ_D is invariant under the action of $\mathrm{GL}_2^+\mathbb{R}$. Each component is a $\mathrm{GL}_2^+\mathbb{R}$ orbit which contains exactly one prototypical eigenform $\mathcal{Q}(e, l, m)$ and is isomorphic to $\mathrm{GL}_2^+\mathbb{R}/\Gamma_0(m)$.*

Remark. Here $\Gamma_0(m)$ is the congruence subgroup of $\mathrm{SL}_2\mathbb{Z}$ defined by

$$\Gamma_0(m) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2\mathbb{Z} : c \equiv 0 \pmod{m} \right\}.$$

Theorem 2.19. *For any nonsquare discriminant D , we have*

$$\chi(Q_D) = -5\chi(X_D). \tag{2.15}$$

If $D = d^2$ with $d \geq 2$, then we have

$$\chi(Q_{d^2}) = -\frac{1}{72}d^2(5d-6) \sum_{r|d} \frac{\mu(r)}{r^2}. \tag{2.16}$$

Proof. We first claim that

$$\sum_{s|f} \chi(Q_{s^2D}) = -\frac{1}{6} \sum_{\substack{e \equiv f^2D \pmod{2} \\ -f\sqrt{D} < e < f\sqrt{D}}} \sigma_1\left(\frac{f^2D - e^2}{4}\right) \tag{2.17}$$

for any fundamental discriminant D and $f \in \mathbb{N}$. By Theorem 2.18, there is one component of Q_D for each prototype (e, l, m) which is isomorphic to $\mathbb{H}/\Gamma_0(m)$. By [Miy89, Theorem 4.2.5],

$$\chi(\mathbb{H}/\Gamma_0(m)) = \psi(m),$$

where

$$\psi(m) = -\frac{1}{6}m \prod_{\substack{p|m \\ p \text{ prime}}} \left(1 + \frac{1}{p}\right).$$

It is elementary to show that for any $n \in \mathbb{N}$,

$$-\frac{1}{6}\sigma_1(n) = \sum_{\substack{n=l^2m \\ l, m \in \mathbb{N}}} \psi(m).$$

By Theorem 2.18,

$$\begin{aligned} \sum_{r|f} \chi(Q_{f^2D}) &= \sum_{\substack{f^2D=e^2+4l^2m \\ l, m > 0}} \psi(m) \\ &= -\frac{1}{6} \sum_{\substack{e \equiv f^2D \pmod{2} \\ -f\sqrt{D} < e < f\sqrt{D}}} \sigma_1\left(\frac{f^2D - e^2}{4}\right), \end{aligned}$$

which proves (2.17).

If $D \neq 1$ is a fundamental discriminant, then for any $f \in \mathbb{N}$, we have

$$\sum_{r|f} \chi(Q_{r^2D}) = \frac{5}{6}H(2, f^2D) = -5 \sum_{r|f} \chi(X_{r^2D})$$

by Theorem 2.16 and (2.17), which proves (2.15).

Now for any $d \in \mathbb{N}$, we have

$$\frac{1}{72} + \sum_{\substack{r|d \\ r > 1}} \chi(Q_{r^2}) = \frac{5}{6}H(2, d^2) + \frac{d^2}{12} \quad (2.18)$$

by Theorem 2.16 and (2.17), and using the convention that $\sigma_1(0) = -1/24$. From (2.11), we get for any $d \in \mathbb{N}$,

$$H(2, d^2) = \sum_{s|d} \left(-\frac{1}{12}s^3 \sum_{r|s} \frac{\mu(r)}{r^2} \right). \quad (2.19)$$

It follows from Möbius inversion that for any $d \in \mathbb{N}$,

$$d^2 = \sum_{s|d} \left(s^2 \sum_{r|s} \frac{\mu(r)}{r^2} \right). \quad (2.20)$$

Define

$$f(d) = \begin{cases} 1/72 & \text{if } d = 1 \\ \chi(Q_{d^2}) & \text{if } d > 1. \end{cases}$$

Combining (2.18), (2.19), and (2.20), we obtain

$$\sum_{s|d} f(s) = \sum_{s|d} \left(\left(-\frac{5}{72}s^3 + \frac{1}{12}s^2 \right) \sum_{r|s} \frac{\mu(r)}{r^2} \right),$$

which implies (2.16) by Möbius inversion. $\blacksquare$

There is an involution of ΩQ_D which interchanges the order of the two factors in (2.12). This involution preserves ΩQ_D and so restricts to an involution τ of ΩQ_D . The involution τ preserves the fibers of the bundle $\Omega Q_D \rightarrow Q_D$, so it induces an involution $\bar{\tau}$ of Q_D .

Lemma 2.20. *The involution $\bar{\tau}$ fixes pointwise the component of Q_D containing the prototypical eigenform $\mathcal{Q}(e, l, m)$ if and only if*

$$\bar{\tau}\mathcal{Q}(e, l, m) = \mathcal{Q}(e, l, m). \quad (2.21)$$

The only such component on which τ is the identity is the one containing $\mathcal{Q}(0, 1, 1)$.

Proof. First, note that (2.21) holds if and only if $\Lambda_1 = \Lambda_2$ in (2.14), which is true if and only if $m = 1$ and $\lambda = l$. The only prototype for which this holds is $(0, 1, 1)$, so (2.21) is true if and only if $(e, l, m) = (0, 1, 1)$.

Clearly, (2.21) must hold for $\bar{\tau}$ to be the identity on the component containing $\mathcal{Q}(e, l, m)$. Conversely, if (2.21) does hold, then $(e, l, m) = (0, 1, 1)$, and

$$\tau(g \cdot \mathcal{Q}(0, 1, 1)) = g \cdot \mathcal{Q}(0, 1, 1)$$

for any $g \in \mathrm{GL}_2^+ \mathbb{R}$ because τ commutes with the action of $\mathrm{GL}_2^+ \mathbb{R}$ on ΩQ_D . Thus τ fixes every point of the $\mathrm{GL}_2^+ \mathbb{R}$ -orbit of $\mathcal{Q}(0, 1, 1)$, which is exactly the component of ΩQ_4 containing $\mathcal{Q}(0, 1, 1)$. $\blacksquare$

There is natural map $\pi: \Omega Q_D \rightarrow \Omega P_D$ defined by

$$\pi((E_1, \omega_1), (E_2, \omega_2)) = (E_1, \omega_1) \oplus (E_2, \omega_2) \quad (2.22)$$

which descends to a map $\bar{\pi}: Q_D \rightarrow P_D$.

Lemma 2.21. *The map $\bar{\pi}: Q_D \rightarrow P_D$ factors through to an isomorphism of orbifolds, $\bar{\pi}: Q_D/\bar{\tau} \rightarrow P_D$.*

Proof. The map $\bar{\pi}$ factors because the right hand side of (2.22) does not depend on the order of the E_i . It is clearly onto. That $\bar{\pi}$ is one-to-one follows from the following fact about Abelian surfaces: a principally polarized Abelian surface can have at most one representation as a polarized product of elliptic curves. $\blacksquare$

Theorem 2.22. *If D is not square, then*

$$\chi(P_D) = -\frac{5}{2}\chi(X_D).$$

If $D = d^2$, then

$$\chi(P_{d^2}) = -\frac{1}{144}d^2(5d-6) \sum_{r|d} \frac{\mu(r)}{r^2}$$

when $d > 2$, and

$$\chi(P_4) = -\frac{1}{6}.$$

Proof. By Theorem 2.18, Q_4 consists of a single component, which contains $\mathcal{Q}(0, 1, 1)$. Thus by Lemmas 2.20 and 2.21, $P_4 \cong Q_4$, and Q_D is a twofold cover of P_D when $D > 4$. The claim then follows from Theorem 2.19. ■

3 Prototypes

3.1 P_D , W_D , and Y_D -prototypes

Curves in $Y_D \setminus X_D$ will be classified by certain discrete numerical invariants which we call Y_D -prototypes. They are almost (but not quite) the same as the *splitting prototypes* introduced by McMullen in [McM05a] to classify the cusps of W_D .

Definition. A Y_D -prototype of discriminant D is a quadruple, $(a, b, c, \bar{q})$ with $a, b, c \in \mathbb{Z}$ and $\bar{q} \in \mathbb{Z}/\gcd(a, b, c)$ which satisfies the following six properties:

1. $b^2 - 4ac = D$
2. $a > 0$
3. $c \leq 0$
4. $\gcd(a, b, c, \bar{q}) = 1$
5. $a + b + c \leq 0$
6. $a + b + c$ and c are not both zero.

Let $\mathcal{Y}_D$ be the set of Y_D -prototypes. Define a map $\lambda: \mathcal{Y}_D \rightarrow K_D$ by associating to each prototype $P = (a, b, c, \bar{q})$ of discriminant P the unique algebraic number $\lambda(P) \in K_D$ such that $a\lambda(P)^2 + b\lambda(P) + c = 0$ and $\lambda^{(1)} > 0$. This makes sense because if $c < 0$, then the two roots of $ax^2 + bx + c = 0$ have opposite signs, and if $c = 0$, then the condition that $a + b < 0$ implies that the nonzero root of $a\lambda^2 + b\lambda = 0$ is positive. It is easy to check that the last condition $a + b + c \leq 0$ is equivalent to $\lambda^{(1)} \geq 1$.

We say that a prototype is *terminal* if $a + b + c = 0$; it is *initial* if $a - b + c = 0$; and it is degenerate if $c = 0$. Terminal, initial, and degenerate prototypes only arise if D is square. We define the involution,

$$(a, b, c, \bar{q}) \mapsto (-c, -b, -a, \bar{q}), \quad (3.1)$$

on the set of terminal prototypes. We consider two terminal prototypes to be the same if they are related by this involution. Similarly, we consider two degenerate prototypes to be the same if they are related by the involution,

$$(a, b, 0, \bar{q}) \mapsto (-b - a, b, 0, \bar{q}) \quad (3.2)$$

on the set of degenerate prototypes.

Operations on Y_D -prototypes. Given a nonterminal prototype P , define the *next prototype* P^+ by

$$P^+ = \begin{cases} (a, 2a + b, a + b + c, \bar{q}), & \text{if } 4a + 2b + c \leq 0; \\ (-a - b - c, -2a - b, -a, \bar{q}), & \text{if } 4a + 2b + c \geq 0. \end{cases}$$

Given a nondegenerate prototype P , define the *previous prototype* P^- by

$$P^- = \begin{cases} (a, -2a + b, a - b + c, \bar{q}), & \text{if } a - b + c \leq 0; \\ (-c, -b + 2c, -a + b - c, \bar{q}), & \text{if } a - b + c \geq 0. \end{cases}$$

Its easy to check that P^+ and P^- are actually prototypes of the same discriminant and that

$$(P^+)^- = (P^-)^+ = P$$

when these operations are defined.

On the level of $\lambda(P)$, we have,

$$\lambda(P^+) = \begin{cases} \lambda(P) - 1, & \text{if } \lambda(P)^{(1)} - 1 \geq 1; \\ 1/(\lambda(P) - 1), & \text{if } \lambda(P)^{(1)} - 1 < 1; \end{cases}$$

and

$$\lambda(P^-) = \begin{cases} \lambda(P) + 1, & \text{if } N_{\mathbb{Q}}^{K_D}(\lambda(P) + 1) \leq 0; \\ (\lambda(P) + 1)/\lambda(P), & \text{if } N_{\mathbb{Q}}^{K_D}(\lambda(P) + 1) > 0. \end{cases}$$

Define an involution t on the set of prototypes of discriminant D by

$$t(a, b, c, \bar{q}) = \begin{cases} (a, -b, c, \bar{q}), & \text{if } a - b + c \leq 0; \\ (-c, b, -a, \bar{q}), & \text{if } a - b + c \geq 0. \end{cases}$$

Define the *multiplicity* of a nondegenerate prototype P by

$$\text{mult}(P) = \frac{\gcd(a, c)}{\gcd(a, b, c)}.$$

P_D and W_D -prototypes. In addition to the Y_D -prototypes, we will define similar objects which we call P_D and W_D -prototypes. These will classify the cusps of the curves P_D and W_D .

Definition. A P_D -prototype of discriminant D is a quadruple, $(a, b, c, \bar{q})$ with $a, b, c \in \mathbb{Z}$ and $\bar{q} \in \mathbb{Z}/\gcd(a, c)$ which satisfies the following five properties:

1. $b^2 - 4ac = D$
2. $a > 0$
3. $c < 0$
4. $\gcd(a, b, c, \bar{q}) = 1$
5. $a + b + c \leq 0$

Note that the only difference between Y_D and P_D -prototypes are that for P_D -prototypes, $\bar{q}$ ranges in a potentially larger group, and c is not allowed to be zero.

A W_D -prototype of discriminant D is a quadruple, $(a, b, c, \bar{q})$ with $a, b, c \in \mathbb{Z}$ and $\bar{q} \in \mathbb{Z}/\gcd(a, c)$ which satisfies the same properties as above except that the last one is replaced with

$$a + b + c < 0.$$

Let $\mathcal{P}_D$ and $\mathcal{W}_D$ be the sets of P_D and W_D -prototypes. Just as for the Y_D -prototypes, we have natural maps $\lambda: \mathcal{W}_D \rightarrow K_D$ and $\lambda: \mathcal{P}_D \rightarrow K_D$. By reducing $\bar{q}$ modulo $\gcd(a, b, c)$, there are natural maps $\mathcal{P}_D \rightarrow \mathcal{Y}_D$ and $\mathcal{W}_D \rightarrow \mathcal{Y}_D$.

We will see in §8.1 that these maps encode intersections of cusps of W_D and P_D with curves in $Y_D \setminus X_D$.

Relation with splitting prototypes. In [McM05a], McMullen defined a splitting prototype to be a quadruple of integers (a, b, c, e) . Splitting prototypes correspond bijectively to our W_D -prototypes by sending the splitting prototype (a, b, c, e) to the W_D -prototype $(c, e, -b, \bar{a})$.

3.2 Quasi-invertible $\mathcal{O}_D$ -modules

In this section, we will study a class of $\mathcal{O}_D$ -modules which arises naturally in the study of our compactification of the Hilbert modular surface. We will also study a class of bases of these modules whose combinatorics is closely related to the geometry of the compactification. This material will only be used in §7.

Definition. An $\mathcal{O}_D$ -module M is *quasi-invertible* if $M \cong \mathbb{Z} \oplus \mathbb{Z}$ as an Abelian group, and M contains some element x such that $\text{Ann}(x) = 0$.

Examples. If D is a fundamental discriminant, then every quasi-invertible $\mathcal{O}_D$ -module is actually invertible. Since $\mathcal{O}_D$ is a Dedekind domain, the set of isomorphism classes of such modules forms a group, the ideal class group of $\mathcal{O}_D$.

If D is not square, then a quasi-invertible $\mathcal{O}_D$ -module is just an invertible $\mathcal{O}_E$ -module over some order $\mathcal{O}_E$ containing $\mathcal{O}_D$. A quasi-invertible $\mathcal{O}_D$ -module is invertible if and only if it is primitive in the sense that it is not also a module over any $\mathcal{O}_E$ containing $\mathcal{O}_D$.

In this paper, quasi-invertible $\mathcal{O}_D$ -modules will arise in the following way:

Proposition 3.1. *If $M \subset \mathcal{O}_D \oplus \mathcal{O}_D^\vee$ is a Lagrangian $\mathcal{O}_D$ -submodule which has $\mathbb{Z}$ -rank two, then M is quasi-invertible.*

Proof. This is trivial if D is not square because then $\mathcal{O}_D \oplus \mathcal{O}_D^\vee$ is torsion-free, so suppose $D = d^2$. Let $M \subset \mathcal{O}_{d^2} \oplus \mathcal{O}_{d^2}^\vee$ be a rank two Lagrangian submodule which is not quasi-invertible, and let $\{u_1, u_2\}$ be a basis of M . We must then have either $(d, 0) \cdot u_i = 0$ for $i = 1, 2$, or $(0, d) \cdot u_i = 0$ for $i = 1, 2$, or else there

would be some $x \in M$ such that $\text{Ann}(x) = 0$. For concreteness, suppose we are in the second case. Then we must have

$$u_i = ((a_i, 0), (b_i, 0))$$

for some $a_i, b_i \in \mathbb{Q}$.

Since M is Lagrangian, we know that,

$$\begin{aligned} 0 = \langle u_1, u_2 \rangle &= \text{Tr}_{\mathbb{Q} \oplus \mathbb{Q}}^{\mathbb{Q} \oplus \mathbb{Q}} \begin{vmatrix} (a_1, 0) & (a_2, 0) \\ (b_1, 0) & (b_2, 0) \end{vmatrix} \\ &= a_1 b_2 - a_2 b_1, \end{aligned}$$

so $u_1 = r u_2$ for some $r \in \mathbb{Q}$. This contradicts the fact that the u_i are a basis. $\blacksquare$

Embeddings in K_D . A lattice in K_D is a rank two Abelian subgroup. Given a lattice $M \subset K_D$, the set

$$\{x \in K_D : xM \subset M\}$$

is called the *coefficient ring* of M . It is an order in K_D .

Proposition 3.2. *An $\mathcal{O}_D$ -module M is quasi-invertible if and only if it is isomorphic to some lattice $M \subset K_D$ whose coefficient ring contains $\mathcal{O}_D$.*

If M is quasi-invertible, then the embedding $M \rightarrow K_D$ is unique up to multiplication by a non-zero-divisor in K_D .

Proof. If D is not square, then it is clear that a lattice $M \subset K_D$ whose coefficient ring contains $\mathcal{O}_D$ is a quasi-invertible $\mathcal{O}_D$ -module because M is automatically torsion-free. If $D = d^2$, then M could have $\text{Ann}(x) \neq 0$ for every $x \in M$ only if $M \subset \mathbb{Q} \oplus \{0\}$ or $M \subset \{0\} \oplus \mathbb{Q}$. This cannot happen because M has rank two.

Conversely, suppose M is a quasi-invertible $\mathcal{O}_D$ -module with $x \in M$ such that $\text{Ann}(x) = 0$. Since M is torsion-free as an Abelian group, the natural map $M \rightarrow M \otimes \mathbb{Q}$ is injective. The tensor product $M \otimes \mathbb{Q}$ is a K_D -module and a two-dimensional vector space over $\mathbb{Q}$, so the map $K \rightarrow M \otimes \mathbb{Q}$ defined by $r \mapsto r \cdot x$ is an isomorphism. The inverse of this map embeds M as a lattice whose coefficient ring contains $\mathcal{O}_D$. Since the image of x determined the embedding, it is unique up to constant multiple. $\blacksquare$

It is easy to identify the coefficient ring of a lattice in K_D . Let M_λ be the lattice generated by $\{1, \lambda\}$. For $\lambda \in K_D$, let $\phi_\lambda(t)$ be the minimal polynomial of λ , the unique polynomial $\phi_\lambda(t) = at^2 + bt + c$ such that $a, b, c \in \mathbb{Z}$, $a > 0$, $\phi_\lambda(\lambda) = 0$, and $\gcd(a, b, c) = 1$.

Proposition 3.3 ([BS66, p.136]). *If $\lambda \in K_D \setminus \mathbb{Q}$ with $\phi_\lambda(t) = at^2 + bt + c$, then the coefficient ring of M_λ is the order $\mathbb{Z}[a\lambda]$, which is isomorphic to $\mathcal{O}_D$, where $D = b^2 - 4ac$.*

Admissible bases. Since a quasi-invertible $\mathcal{O}_D$ -module M can be embedded in K_D uniquely up to constant multiple, for any $u, v \in M$ with $\text{Ann}(v) \neq 0$, the ratio u/v is a well-defined element of K_D .

Definition. A basis $\{u, v\}$ of a quasi-invertible $\mathcal{O}_D$ -module M is *admissible* if $\text{Ann}(v) = 0$ and $N_{\mathbb{Q}}^{K_D} < 0$.

We will consider two admissible bases of M to be equivalent if they are the same as subsets of $M/\pm 1$. The following proposition gives a complete classification of quasi-invertible $\mathcal{O}_D$ -modules together with an admissible basis.

Proposition 3.4. *Every pair $(M, \{u, v\})$, where M is a quasi-invertible $\mathcal{O}_D$ -module together with an admissible basis $\{u, v\}$, is equivalent to one of the form $(M_\lambda, \{1, \lambda\})$, where $\lambda^{(1)} \geq 1$ and $\psi(\lambda) = 0$, where*

$$\psi(t) = at^2 + bt + c$$

for some integers (a, b, c) such that:

1. $b^2 - 4ac = D$
2. $a > 0$
3. $c < 0$
4. $a + b + c \leq 0$
5. $\mathcal{O}_D = \mathbb{Z}[a\lambda]$

Pairs $(M_\lambda, \{1, \lambda\})$ and $(M_\mu, \{1, \mu\})$ of this form are equivalent if and only if $\lambda = \mu^{-1}$.

Proof. If $\lambda > 1$, then by Proposition 3.2, there is a unique embedding $M \rightarrow K_D$ which – after possibly switching u and v and changing their signs – sends u to 1 and v to some $\lambda \in K$ with $\lambda^{(1)} > 1$. The coefficient ring of the image M_λ is $\mathcal{O}_E$, where $D = s^2 E$ for some quadratic discriminant E . Let $\phi_\lambda(t) = a't^2 + b't + c'$ be the minimal polynomial of λ . By Proposition 3.3, we have $b'^2 - 4a'c' = E$ and $\mathcal{O}_E = \mathbb{Z}[a'\lambda]$. We have $c' < 0$ because $N_{\mathbb{Q}}^{K_D} < 0$. Since $\lambda^{(1)} > 1$, we know that $a' + b' + c' \leq 0$. The integers $(a, b, c) = s(a', b', c')$ then have the required properties.

If $\lambda^{(1)} = 1$, then there are two embeddings $M \rightarrow K_D$ as above, one sending u to 1 and v to λ ; the other sending u to λ^{-1} and v to 1. As in the above paragraph, both embeddings yield a presentation of $(M, \{u, v\})$ in the desired form. ■

We can define a map from the set of nondegenerate Y_D -prototypes to the set of isomorphism classes of admissible bases of quasi-invertible $\mathcal{O}_D$ -modules by sending P to $(M_{\lambda(P)}, \{1, \lambda(P)\})$. This proposition implies that this map is onto and that two prototypes are sent to the same admissible basis if and only if their integers a , b , and c are the same.

Characterization of admissible bases. We now describe a useful characterization of admissible bases. Given two quasi-invertible $\mathcal{O}_D$ -modules M and N , a perfect pairing is a bilinear map

$$h: M \times N \rightarrow \mathbb{Z}$$

such that

$$h(\lambda \cdot x, y) = h(x, \lambda \cdot y)$$

for each $x \in M$, $y \in N$, and $\lambda \in \mathcal{O}_D$, and such that the induced map $M \rightarrow \text{Hom}(N, \mathbb{Z})$ is an isomorphism.

Given such a perfect pairing and a basis $\{u, v\}$ of M , let $\{u^*, v^*\}$ be a dual basis of N , and define

$$\text{sign}(u, v; h) = \text{sign}(\iota_M(u)\iota_M(v)\iota_N(u^*)\iota_N(v^*)),$$

where ι_M and ι_N are ι_1 -linear, nonzero maps $M \rightarrow \mathbb{R}$ and $N \rightarrow \mathbb{R}$. Note that $\text{sign}(u, v; h)$ is invariant under changing the sign of u or v as well as independent of the choice of ι_M and ι_N .

Theorem 3.5. *Given a quasi-invertible $\mathcal{O}_D$ -module M , a basis $\{u, v\}$ is an admissible basis if and only if $\text{sign}(u, v; h) = 1$ for some (or any) perfect pairing $h: M \times N \rightarrow \mathbb{Z}$.*

Proof. Assume that M is embedded in K_D . If $u^{(1)} = 0$ or $v^{(1)} = 0$, then $\text{sign}(u, v; h) = 0$, and the basis is not admissible, so we are done. Since M is quasi-invertible, it can't happen that $u^{(2)} = v^{(2)} = 0$, so assume $u^{(2)} \neq 0$.

Assume without loss of generality that $u = 1$. By possibly changing the sign of v , we can suppose $v^{(1)} > 0$. Then $v^{(2)} < 0$ if and only if $\{u, v\}$ is admissible. Since N and the perfect pairing h are unique up to isomorphism, we can suppose without loss of generality that $N = M^\vee$, the inverse different of M , and that h is the trace pairing $M \times M^\vee \rightarrow \mathbb{Z}$.

Define

$$u^* = \frac{-v'}{v - v'}$$

$$v^* = \frac{1}{v - v'}.$$

It is easy to check that $\{u^*, v^*\}$ is dual to $\{u, v\}$ with respect to the trace pairing. If $\{u, v\}$ is not admissible, then $v^{(2)} < 0$, in which case $(u^*)^{(1)}, (v^*)^{(1)} > 0$, so $\text{sign}(u, v; h) > 0$. If $\{u, v\}$ is not admissible, then $v^{(2)} \geq 0$. If $v^{(2)} > 0$, then $(u^*)^{(1)}$ and $(v^*)^{(1)}$ have opposite signs, so $\text{sign}(u, v; h) = -1$. If $v^{(2)} = 0$, then $(u^*)^{(1)} = 0$, so $\text{sign}(u, v; h) = 0$. ■

Admissible triples. We are also interested in special triples of elements of quasi-invertible $\mathcal{O}_D$ -modules which will arise naturally in the study of the compactification of the Hilbert modular surface.

Definition. An *admissible triple* in a quasi-invertible $\mathcal{O}_D$ -module M is an unordered triple of elements $\{a, b, c\} \subset M/\pm 1$ such that:

- $\pm a \pm b \pm c = 0$ for some choice of signs.
- Some pair of elements of the triple form an admissible basis of M .

Proposition 3.6. *Every pair (M, T) , where M is a quasi-invertible $\mathcal{O}_D$ -module with an admissible triple $T \subset M$, is equivalent to one of the form*

$$(M_\lambda, \{1, \lambda, \lambda - 1\}),$$

where $\lambda^{(1)} \geq 1$ and

$$a\lambda^2 + b\lambda + c = 0$$

for some integers (a, b, c) such that:

1. $b^2 - 4ac = D$
2. $a > 0$
3. $c \leq 0$
4. $a + b + c \leq 0$
5. $a + b + c$ and c are not both zero.
6. $\mathcal{O}_D = \mathbb{Z}[a\lambda]$

Two such pairs $(M_\lambda, \{1, \lambda, \lambda - 1\})$ and $(M_\mu, \{1, \mu, \mu - 1\})$ are equivalent if and only if $\lambda^{(1)} = 1$ and $\mu = \lambda^{-1}$ or if $\lambda^{(2)} = 0$ and $\mu = \lambda/(\lambda - 1)$.

Proof. Let $\{u, v, w\} \subset M$ be an admissible triple and assume that M is embedded in K_D . This triple contains at least one admissible basis, so assume without loss of generality that $\{u, v\}$ is an admissible basis. There are now three cases to consider, depending on whether $\text{Ann}(w) = 0$, $w^{(1)} = 0$, or $w^{(2)} = 0$.

First suppose that $\text{Ann}(w) = 0$. Then $\{u, v, w\}$ contains exactly two admissible bases because

$$N_{\mathbb{Q}}^{K_D} \left(\frac{u}{v} \right) N_{\mathbb{Q}}^{K_D} \left(\frac{v}{w} \right) N_{\mathbb{Q}}^{K_D} \left(\frac{w}{u} \right) = 1.$$

Suppose without loss of generality that u is contained in both bases. Dividing everything by u , we can assume $u = 1$. By possibly switching v and w and changing their signs, we can put the triple uniquely in the form $\{1, \lambda, \lambda - 1\}$ with $\lambda^{(1)} > 0$. Let (a, b, c) be as in Proposition 3.3 applied to M_λ . Since $\lambda^{(1)}\lambda^{(2)} < 0$ and $(\lambda^{(1)} - 1)(\lambda^{(2)} - 1) < 0$, we must actually have $\lambda^{(1)} > 1$, and it follows that $a + b + c < 0$. The other necessary properties of a , b , and c follow from Proposition 3.3.

Now assume $w^{(1)} = 0$. In this case, $K_D = \mathbb{Q} \oplus \mathbb{Q}$, and there are two ways to put the triple in the form, $\{1, \lambda, \lambda - 1\}$, where $\lambda = (1, s)$ with $s < 0$: either divide $\{u, v, w\}$ by u , or divide by v and swap the first two elements. With

(a, b, c) as in Proposition 3.3 applied to M_λ , it follows that $a + b + c = 0$ and $c < 0$. The other necessary properties are clear, and the two λ which arise in this way are u/v and v/u , so they are reciprocal.

Now assume $w^{(2)} = 0$. Dividing the triple by u and switching v and w puts it in the form $\{1, \lambda, \lambda - 1\}$ with $\lambda = (s, 0)$. Since $\lambda - 1 = (s - 1, -1)$ has negative norm, we must have $s > 1$. It follows that if (a, b, c) are as in Proposition 3.3 applied to M_λ , we must have $c = 0$ and $a + b < 0$. Alternatively, we could have divided the triple by v and rearranged the elements to put it in the form $\{1, \mu, \mu - 1\}$ with $\mu = (t, 0)$. Then $t > 1$ by the same argument, and $t = s/(s - 1)$.

The converse statement that triples of the given form are admissible is not hard and will be left to the reader. ■

Just as for admissible pairs, we can define a map from the set of Y_D prototypes to the set of isomorphism classes of quasi-invertible $\mathcal{O}_D$ -modules with admissible triples, sending $(a, b, c, \bar{q})$ to $(M_\lambda, \{1, \lambda, \lambda - 1\})$ as above. Since we identified two terminal or degenerate prototypes if they are related by the involutions (3.1) and (3.2), two prototypes have the same image if and only if they have the same $\bar{q}$.

4 Abelian differentials

In this section, we recall known material about Abelian differentials (holomorphic one-forms) on compact Riemann surfaces. In §4.1, we discuss the flat geometry associated to an Abelian differential. In §4.2, we discuss moduli spaces of Abelian differentials and the action of $\mathrm{SL}_2\mathbb{R}$ on these spaces. In §4.3, we discuss McMullen's results on the dynamics of this action in genus two.

4.1 Flat geometry of Abelian differentials

A Riemann surface with a nonzero Abelian differential (holomorphic one-form) carries a canonical flat geometry, which is closely related to the study of billiards in rational angled polygons, as well as the study of flows on moduli space. In this paper, this geometry will be useful because it will allow us to deform Abelian differentials using some concrete cut-and-paste operations which we will describe in this section.

Translation surfaces. Let X be a Riemann surface with a nonzero Abelian differential ω , and let $Z(\omega)$ be the discrete set of zeros of ω . Every point of X which is not a zero of ω has a neighborhood U and a conformal map $\phi: U \rightarrow \mathbb{C}$ such that $\omega|_U = \phi^*(dz)$. The conformal map ϕ can be defined explicitly by

$$\phi(z) = \int_{z_0}^z \omega$$

for a choice of base point z_0 . These coordinates ϕ are unique up to translation by a constant.

We can also put ω in a standard form in the neighborhood of a zero or a pole at z_0 . There is a neighborhood U of z_0 and conformal map

$$\phi_0: (U, z_0) \rightarrow (\mathbb{C}, 0)$$

such that

$$\omega|_U = \begin{cases} \phi_0^*(z^n dz), & \text{if } z_0 \text{ is not a simple pole;} \\ \phi_0^*(a dz/z), & \text{if } z_0 \text{ is a simple pole with nonzero residue } a. \end{cases} \quad (4.1)$$

Since the local coordinates ϕ away from the zeros of ω are unique up to translation by a constant, any translation invariant geometric structure on $\mathbb{C}$ is inherited by X . In particular, X inherits a flat metric and an oriented foliation $\mathcal{F}_h$ coming from the foliation of the plane by horizontal lines. More generally, for slope $s \in \mathbb{P}^1(\mathbb{R})$, the surface X has an orientable foliation $\mathcal{F}_s$ coming from the foliation of $\mathbb{C}$ by lines of slope s .

In terms of the Abelian differential ω , the flat metric is just $|\omega|$. A vector v is tangent to $\mathcal{F}_h$ in the positive direction if $\omega(v) > 0$.

The flat metric has singularities at the zeros. From the coordinates (4.1), we see that a zero of order n has a neighborhood isometric to a cone with cone




Figure 2: Zeros of order one and two.

angle $2\pi(n+1)$, and the foliation $\mathcal{F}_h$ has $2n+2$ leaves meeting at the zero. Zeros of orders one and two together with the foliation $\mathcal{F}_h$ are pictured in Figure 2.

To summarize, we have associated to an Abelian differential (X, ω) , a flat metric on $X \setminus Z(\omega)$ together with a horizontal foliation $\mathcal{F}_h$ which is parallel with respect to a metric such that the points of $Z(\omega)$ are cone singularities of the metric. A surface with such a structure is sometimes called a *translation surface*. A translation surface is equivalent to a Riemann surface with a nonzero Abelian differential.

A geodesic on (X, ω) is called *straight* if it does not pass through any zeros of ω . A straight geodesic which joins two zeros is called a *saddle connection*. Any straight, closed geodesic is contained in a *cylinder* on X by taking nearby parallel geodesics. If the genus of X is greater than one, any cylinder can be extended until either end contains a zero of ω and each boundary component is a finite union of saddle connections. Such a cylinder is called a *maximal cylinder*.

Plumbing a cylinder. Cylinders arise from simple poles: by (4.1), a simple pole of an Abelian differential has a neighborhood which is isometric to a half-infinite cylinder. Given a meromorphic Abelian differential (X, ω) with simple poles at p and q such that

$$\text{Res}_p \omega = -\text{Res}_q \omega, \quad (4.2)$$

there is a simple surgery operation which allows us to replace the poles at p and q with a cylinder. Cut X along two closed geodesics, one in each of the two half-infinite cylinders around p and q , and then glue X along the resulting boundary components by an isometry. The condition (4.2) means exactly that this gluing map is locally a translation; therefore, we get a new Abelian differential with two fewer poles. Call this operation *plumbing a cylinder*. It depends on two parameters: the height of the resulting cylinder and the amount of twisting of the gluing map.

We can reverse this operation to replace a cylinder with two simple poles. Just cut a cylinder along a closed geodesic, and then glue two half-infinite cylinders to the resulting boundary components. Call this operation *unplumbing a cylinder*. These operations will be used in §6 to define polar coordinates around the boundary of the Deligne-Mumford compactification of moduli space.

Connected Sums. Two Abelian differentials (X_1, ω_1) and (X_2, ω_2) can be combined into one by taking a *connected sum*. This operation is studied in detail in [McMa].

Let I be a line segment in the complex plane. Suppose I is embedded in each of the surfaces (X_i, ω_i) by an isometric embedding $\epsilon_i: I \rightarrow X_i$ preserving the slope of I as well as its length. Now cut each of the X_i along I , and glue the surfaces together by gluing one boundary component along I on one of the surfaces to the opposite boundary component on the other surface. The gluing maps are just translations in the flat structures on the (X_i, ω_i) , so the resulting surface has a flat structure as well away from the ends of I . By the Riemann removable singularity theorem, the conformal structure and Abelian differential can actually be extended to the ends of I . Thus we obtain a new Abelian differential (X, ω) . If the segments $\epsilon_i(I) \subset X_i$ are disjoint from the zeros of ω , then the ω has simple zeros.

We call the resulting Abelian differential,

$$(X, \omega) = (X_1, \omega_1) \#_I (X_2, \omega_2),$$

the *connected sum* of (X_1, ω_1) and (X_2, ω_2) along I .

This construction can be modified in the obvious way to perform a self-connected sum of an Abelian differential with itself, given two parallel embeddings of I in that Abelian differential.

An Abelian differential (X, ω) resulting from a connected sum has two simple zeros p and q and two oriented saddle connections I_1 and I_2 going from p to q such that

$$\int_{I_1} \omega = \int_{I_2} \omega. \quad (4.3)$$

Given any Abelian differential (X, ω) with a pair of oriented, embedded saddle connections I_1 and I_2 both beginning and ending at the same zeros and satisfying (4.3), we can reverse the connected sum operation. To do this, cut X along I_1 and I_2 and then reglue to get a new Abelian differential (X', ω') . Equation (4.3) implies that this gluing can be done by a translation. This operation is called *splitting along I_1 and I_2* and is inverse to the connected sum operation.

In this paper, there are two main cases where we will use these constructions. First, suppose (X_1, ω_1) and (X_2, ω_2) are both genus one, and a segment $I \subset \mathbb{C}$ is embedded in each by embeddings ϵ_i as above. Then we can form the connected sum along I , and the resulting Abelian differential has genus two with two simple zeros. Conversely, splitting a genus two Abelian differential along a pair of saddle connections I_1 and I_2 such that $I_1 \cup I_2$ separates the surface yields a pair of genus one Abelian differentials.

Second, given an genus one Abelian differential with two embedded segments I_i which are parallel and of the same length, we can form a self connected sum along these segments. This again yields a genus two Abelian differential with two simple zeros. Conversely, splitting along a pair of saddle connections on a genus two Abelian differential satisfying (4.3) which don't separate the surface yields a single genus one Abelian differential.

Splitting a double zero. There is also a cut-and-paste operation which replaces a zero of an Abelian differential with two zeros of lower order. This operation is explained in detail in [KZ03] and [EMZ03]. We will describe this operation in the case of a double zero, the only case we need.

Let (X, ω) be an Abelian differential with a double zero at p , and choose a straight geodesic segment I starting at p . The segment I is a leaf of the foliation $\mathcal{F}_\theta$ of some slope θ . On X , draw an “X” composed of the segment I and three other geodesic segments of the same length and slope as in Figure 3. (In this figure, leaves of $\mathcal{F}_\theta$ are represented by dotted or solid lines, with the orientation indicated by an arrow, and the “X” is represented by solid lines. Two consecutive segments meeting at the zero form a 180° angle.) Assume that I was chosen to be short enough so that each arm of the “X” is an embedded straight geodesic on X not meeting any of the other arms.




Figure 3: Splitting a double zero.

Now cut the surface along the “X” to obtain a surface with eight geodesic boundary components, and then glue the boundary components pairwise so that the components marked with the same letter in Figure 3 are glued together. The result is a new Abelian differential with simple zeros at the two points where four leaves of $\mathcal{F}_\theta$ meet in the figure. We will denote this new Abelian differential by

$$(X, \omega) \#_I.$$

This operation can also be reversed in the obvious way. Given a saddle connection of length l joining two distinct zeros, draw two more segments of length $l/2$ emanating from the zeros with the same slopes as in the bottom of Figure 3. If these segments can be drawn without intersecting each other or any other zero, then the process of splitting a double zero can be reversed: cut along these three segments and then reglue, following Figure 3 in reverse. This operation is called *collapsing a saddle connection*.

This operation, together with the connected sum operation, will be used in §12 to parameterize tubular neighborhoods of $\overline{W}_D$ and $\overline{P}_D$ in Y_D .

4.2 Moduli of Abelian differentials

In this subsection, we introduce the moduli space of Abelian differentials and discuss its geometry.

Teichmüller space. Let Σ_g be a connected, closed, oriented, topological surface of genus g , and let $\Sigma_{g,n}$ be a genus g topological surface with n marked points. A marked Riemann surface is a Riemann surface X together with a homeomorphism $\Sigma_{g,n} \rightarrow X$. Two marked Riemann (f, X) and (g, Y) marked by $\Sigma_{g,n}$ are considered to be equivalent if $g \circ f^{-1}$ is homotopic to a conformal isomorphism by a homotopy fixing the marked points.

Let the *Teichmüller space* $\mathcal{T}(\Sigma_{g,n})$ be the space of all Riemann surfaces marked by $\Sigma_{g,n}$ up to equivalence. It has a topology induced by the well-known Teichmüller metric and is homeomorphic to $\mathbb{C}^N$, where

$$N = \begin{cases} 1 & \text{if } g = 1 \text{ and } n = 0 \text{ or } 1; \\ n - 1 & \text{if } g = 1 \text{ and } n > 1; \\ n - 3 & \text{if } g = 0 \text{ and } n > 2; \\ 3g - 3 + n & \text{if } g > 1. \end{cases}$$

We will use the abbreviation $\mathcal{T}_{g,n}$ or $\mathcal{T}_g$ when we don’t need to emphasize the surface Σ_g .

Bers gave $\mathcal{T}_{g,n}$ a complex structure by defining an embedding $B: \mathcal{T}_{g,n} \rightarrow \mathbb{C}^N$ which is a homeomorphism onto its image, a bounded domain in $\mathbb{C}^N$.

Moduli space. The modular group $\text{Mod}(\Sigma_g)$ is the group of all self homeomorphisms of Σ_g up to isotopy. Similarly, $\text{Mod}(\Sigma_{g,n})$ is the group of all self

homeomorphisms which preserve the marked points, up to isotopy preserving the marked points.

An element $\gamma \in \text{Mod}(\Sigma_{g,n})$ acts on $\mathcal{T}(\Sigma_{g,n})$ by replacing a marking

$$f: \Sigma_{g,n} \rightarrow X$$

with $f \circ \gamma^{-1}$. This defines a biholomorphic action of $\text{Mod}(\Sigma_{g,n})$ on $\mathcal{T}(\Sigma_{g,n})$ which is properly discontinuous. Let $Z \subset \Sigma_{g,n}$ be the set of marked points. The stabilizer of a point (f, X) is isomorphic to the group $\text{Aut}(X, f(Z))$ of conformal automorphisms of X preserving the marked points. In genus two, every Riemann surface has an order-two automorphism J , the hyperelliptic involution. This yields an element of order two in $\text{Mod}(\Sigma_2)$ which acts trivially on $\mathcal{T}(\Sigma_2)$.

The *moduli space* of genus g Riemann surfaces with n marked points is the quotient

$$\mathcal{M}_{g,n} = \mathcal{T}(\Sigma_{g,n}) / \text{Mod}(\Sigma_{g,n}).$$

$\mathcal{M}_{g,n}$ is a complex orbifold.

Bundles of Abelian differentials. Given a Riemann surface X , let $\Omega(X)$ be the space of Abelian differentials on X , a rank g complex vector space. Let $\Omega\mathcal{T}_g$ be the space of all pairs (X, ω) with $X \in \mathcal{T}_g$ and $\omega \in \Omega(X)$ a nonzero Abelian differential. We can give $\Omega\mathcal{T}_g$ the structure of a trivial holomorphic punctured vector bundle as follows.

Over $\mathcal{T}_g$, there is the universal curve $\mathcal{CT}_g$, defined by Bers in [Ber73]. It is a complex manifold and comes with a proper map $\pi: \mathcal{CT}_g \rightarrow \mathcal{T}_g$ whose fiber over a Riemann surface X is isomorphic to X itself. The cotangent bundle to the fibers of π is a line bundle $\mathcal{L} \rightarrow \mathcal{CT}_g$. The push-forward $\pi_*\mathcal{O}(\mathcal{L})$ of the sheaf of sections of $\mathcal{L}$ is a sheaf on $\mathcal{T}_g$. The following theorem follows from [Ber61b].

Theorem 4.1. *The sheaf $\pi_*\mathcal{O}(\mathcal{L})$ is the sheaf of sections of a trivial bundle over $\mathcal{T}_g$ whose fiber over a Riemann surface X is $\Omega(X)$.*

The action of the mapping class group Mod_g on $\mathcal{T}_g$ extends to an action on $\Omega\mathcal{T}_g$. The quotient is a rank g orbifold vector bundle $\Omega\mathcal{M}_g$ over $\mathcal{M}_g$ whose fiber over a Riemann surface X is the quotient $\Omega(X)/\text{Aut}(X)$. This bundle is sometimes called the *Hodge bundle*.

In general, when S is any sort of space of Riemann surfaces, ΩS will denote the natural bundle of Abelian differentials over S .

Strata. The bundles $\Omega\mathcal{T}_g$ have a natural stratification in terms of the types of zeros of the Abelian differentials. Each nonzero Abelian differential on a nonsingular Riemann surface has $2g - 2$ zeros, counting multiplicity. Given a sequence of integers $\mathbf{n} = (n_i)_{i=1}^r$ such that $\sum n_i = 2g - 2$, let $\Omega\mathcal{T}_g(\mathbf{n})$ be the locus of all Abelian differentials which have r zeros whose multiplicities are given by the n_i . This locus is a locally closed subset of $\Omega\mathcal{T}_g$, and by Veech [Vee90] it is actually a complex submanifold. The quotient

$$\Omega\mathcal{M}_g(\mathbf{n}) = \Omega\mathcal{T}_g(\mathbf{n}) / \text{Mod}$$

is then a complex suborbifold of $\Omega\mathcal{M}_g$.

Period coordinates. These strata have natural coordinates defined in terms of their periods by Veech [Vee90] and Masur [Mas82] which give the stratum $\Omega\mathcal{T}_g(\mathbf{n})$ the structure of an affine manifold.

There is a fiber bundle of homology groups

$$\mathcal{H}_1 \rightarrow \Omega\mathcal{T}_g(\mathbf{n}),$$

whose fiber over an Abelian differential (X, ω) is $H_1(X, Z(\omega); \mathbb{Z})$, where $Z(\omega)$ is the set of zeros of ω . Given two Abelian differentials (X_1, ω_1) , and $(X_2, \omega_2) \in \Omega\mathcal{T}_g(\mathbf{n})$ with (X_2, ω_2) sufficiently close to (X_1, ω_1) there is a natural isomorphism

$$H_1(X_2, Z(\omega_2)) \rightarrow H_1(X_1, Z(\omega_1)).$$

This defines a flat connection on $\mathcal{H}_1$, the *Gauss-Manin connection*.

Consider $(X, \omega) \in \Omega\mathcal{T}_g(\mathbf{n})$ and a small neighborhood U of (X, ω) . Any $(X', \omega') \in U$ defines an element of $H^1(X', Z(\omega'); \mathbb{C})$ via the periods of ω' . Composing this with the isomorphism

$$H^1(X', Z(\omega'); \mathbb{C}) \rightarrow H^1(X, Z(\omega); \mathbb{C})$$

from the Gauss-Manin connection, we get a map

$$\phi: U \rightarrow H^1(X, Z(\omega); \mathbb{C}),$$

the *period coordinates*. Veech [Vee90] showed that these are in fact biholomorphic coordinate charts.

A choice of basis of $H_1(X, Z(\omega); \mathbb{Z})$ defines an isomorphism

$$H^1(X, Z(\omega); \mathbb{C}) \rightarrow \mathbb{C}^{2g+n-1},$$

where n is the number of zeros of ω . We can suppose that the basis is of the form,

$$\{u_1, \dots, u_{2g}, v_1, \dots, v_{n-1}\}, \quad (4.4)$$

where $\{u_1, \dots, u_{2g}\}$ is a symplectic basis of $H_1(X)$. Two such bases are related by a matrix in the group,

$$G = \left\{ \begin{pmatrix} A & B \\ 0 & C \end{pmatrix} : A \in \mathrm{SP}_{2g}\mathbb{Z}, \quad B \in M_{2g,n}(\mathbb{Z}), \quad C \in \mathrm{GL}_n\mathbb{Z} \right\}.$$

Changing the basis by a matrix in G changes the period coordinates by the transpose of this matrix. This gives $\Omega\mathcal{T}_g(\mathbf{n})$ the structure of an $(G, \mathbb{C}^{2g+n-1})$ -manifold, and $\Omega\mathcal{M}_g(\mathbf{n})$ inherits the structure of a $(G, \mathbb{C}^{2g+n-1})$ -orbifold.

Measures. Since the action of $\mathrm{GL}_n(\mathbb{Z})$ on $\mathbb{C}$ preserves Lebesgue measure, we can pull back this measure by the period coordinates to get the *period measure* $\mu(\mathbf{n})$ on $\Omega\mathcal{M}_g(\mathbf{n})$.

The flat metric defined by an Abelian differential has area given by,

$$\mathrm{Area}(\omega) = \frac{i}{2} \int_X \omega \wedge \bar{\omega}.$$

Let $\Omega_{\leq 1}\mathcal{M}_g(\mathbf{n})$ be the locus of Abelian differentials with $\text{Area}(\omega) \leq 1$, and $\Omega_1\mathcal{M}_g(\mathbf{n})$ the locus of Abelian differentials with $\text{Area}(\omega) = 1$ (we will use the prefixes $\Omega_{\leq 1}$ and Ω_1 to denote the analogous subsets of any space of Abelian differentials). We can define a measure $\mu'(\mathbf{n})$ on $\Omega_1\mathcal{M}_g(\mathbf{n})$ by restricting $\mu(\mathbf{n})$ to $\Omega_{\leq 1}\mathcal{M}_g(\mathbf{n})$ and then projecting by the natural map $\Omega_{\leq 1}\mathcal{M}_g(\mathbf{n}) \rightarrow \Omega_1\mathcal{M}_g(\mathbf{n})$.

The following theorem was proved by Veech and Masur:

Theorem 4.2 ([Mas82],[Vee90]). *The measures $\mu'(\mathbf{n})$ have finite total volume.*

Action of $\text{SL}_2\mathbb{R}$. Given a nonzero Abelian differential $(X, \omega) \in \Omega\mathcal{T}_g$, we can as in §4.1 choose an atlas of coordinate charts $\phi_i: U_i \rightarrow \mathbb{C}$, which cover the complement of the zeros, such that ϕ_i pulls back the form dz on $\mathbb{C}$ to the form ω on X . These coordinate charts differ by translations on their overlap. Now, given an element $A \in \text{SL}_2\mathbb{R}$, define new coordinate charts $\phi'_i = A \circ \phi_i$ by composing with the usual action of A in the complex plane. The new coordinates still differ by translations, and so define a new translation surface. This defines a new Abelian differential $A \cdot (X, \omega) \in \Omega\mathcal{T}_g$ which has a different complex structure than X unless A happens to be a rotation in $\text{SO}_2\mathbb{R}$. If $(X', \omega') = A \cdot (X, \omega)$, then there is a natural real-affine map $h_A: (X, \omega) \rightarrow (X', \omega')$ which takes zeros to zeros. The periods satisfy the relation,

$$\omega'((h_A)_*(\gamma)) = a \cdot \omega(\gamma), \quad (4.5)$$

for any $\gamma \in H_1(X, Z(\omega))$ with $Z(\omega)$ the set of zeros of ω .

This construction defines an action of $\text{SL}_2\mathbb{R}$ on $\Omega\mathcal{T}_g$. This action commutes with the action of Mod and so defines an action of $\text{SL}_2\mathbb{R}$ on $\Omega\mathcal{M}_g$ which preserves to locus $\Omega_1\mathcal{M}_g$ of Abelian differentials with area one.

Equation (4.5) can be used to show that the measure $\mu(\mathbf{n})$ is $\text{SL}_2\mathbb{R}$ invariant, as is $\mu'(\mathbf{n})$.

This action is closely related to the Teichmüller geodesic flow: the projection $\Omega\mathcal{M}_g \rightarrow \mathcal{M}_g$ sends $\text{SL}_2\mathbb{R}$ orbits to copies of the hyperbolic plane in $\mathcal{M}_g$ which are isometrically embedded with respect to the Teichmüller metric, and the restriction of the action to the one parameter subgroup of diagonal matrices is the Teichmüller geodesic flow.

The strata $\Omega\mathcal{M}_g(\mathbf{n})$ and $\Omega_1\mathcal{M}_g(\mathbf{n})$ are invariant under the action of $\text{SL}_2\mathbb{R}$. The measures μ and μ' defined in §5.2 are invariant measures by (4.5).

The natural projection $\Omega_1\mathcal{M}_g \rightarrow \mathbb{P}\Omega\mathcal{M}_g$ sends $\text{SL}_2\mathbb{R}$ orbits to immersed copies of the hyperbolic plane, giving a foliation $\mathcal{FM}_g$ of $\mathbb{P}\Omega\mathcal{M}_g$ by immersed hyperbolic planes. It is possible for a leaf of $\mathcal{FM}_g$ to be a closed subset of $\mathbb{P}\Omega\mathcal{M}_g$. In that case it is called a *Teichmüller curve* because it is an algebraic curve whose projection to $\mathcal{M}_g$ is isometrically immersed with respect to the Teichmüller metric. It is an important unsolved problem to classify Teichmüller curves in $\mathbb{P}\Omega\mathcal{M}_g$ or more generally to classify orbit closures or invariant measures.

4.3 $\mathrm{SL}_2\mathbb{R}$ orbits in genus two.

In genus two, McMullen has completely classified the orbit closures for the action of $\mathrm{SL}_2\mathbb{R}$ on $\Omega_1\mathcal{M}_2$ and the ergodic, invariant measures in the series of papers [McM03a, McMa, McM05a, McM06]. In this subsection, we will discuss the subsets of $\Omega_1\mathcal{M}_2$ which appear in this classification and which will be the focus of this paper.

A $\mathrm{SL}_2\mathbb{R}$ -invariant subset S of $\Omega_1\mathcal{M}_2$ corresponds to a subset $\mathbb{P}S$ of $\mathbb{P}\Omega\mathcal{M}_2$ which is *saturated* in the sense that leaves of the foliation $\mathcal{FM}_2$ are either contained in $\mathbb{P}S$ or disjoint from $\mathbb{P}S$. Invariant measures on $\Omega_1\mathcal{M}_2$ correspond by a disintegration construction to holonomy invariant measures on $\mathcal{FM}_2$. In this paper, we will adopt this point of view and focus on leaves of this foliation rather than on $\mathrm{SL}_2\mathbb{R}$ orbits in $\Omega_1\mathcal{M}_2$.

Eigenform loci. A form $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2$ is an *eigenform for real multiplication by $\mathcal{O}_D$* if the Jacobian of X has real multiplication by $\mathcal{O}_D$ with ω a nonzero eigenform. Let the *eigenform locus* E_D be the locus of all such pairs. The following proposition follows from [McMa, Corollary 5.7].

Proposition 4.3. *E_D is a closed, saturated subset of $\mathbb{P}\Omega\mathcal{M}_2$.*

An embedding $\iota_i: \mathcal{O}_D \rightarrow \mathbb{R}$ determines a map

$$j_i: E_D \rightarrow X_D$$

by sending an eigenform $(X, \omega) \in E_D$ to the pair $(\mathrm{Jac}(X), \rho)$, where

$$\rho: \mathcal{O}_D \rightarrow \mathrm{End} \mathrm{Jac}(X)$$

is chosen so that

$$\omega(\rho(\lambda) \cdot \gamma) = \lambda^{(i)} \omega(\gamma)$$

for each $\gamma \in H_1(X; \mathbb{Z})$. Recall that in §2, we introduced the locus P_D of Abelian varieties in X_D which are polarized products of elliptic curves. The following follows from Proposition 5.4.

Proposition 4.4. *The map j_i is an isomorphism of E_D onto $X_D \setminus P_D$.*

In this paper, we will implicitly identify E_D with $X_D \setminus P_D$ by the isomorphism j_1 .

Let $\mathcal{F}_D$ be the foliation of E_D by Riemann surfaces induced by the foliation $\mathcal{F}$ of $\mathbb{P}\Omega\mathcal{M}_2$. The foliation $\mathcal{F}_D$ of E_D extends to a foliation – which we will continue to call $\mathcal{F}_D$ – of X_D defined by adding the connected components of P_D as leaves of $\mathcal{F}_D$. McMullen [McMb] proved that this is actually a foliation of X_D .

Elliptic differentials. There is a useful alternative characterization of eigenforms (X, ω) for real multiplication by $\mathcal{O}_{d^2}$.

A branched cover $f: X \rightarrow E$ from a Riemann surface to an elliptic curve is said to be primitive if it does not factor through an isogeny of elliptic curves $g: E' \rightarrow E$ of degree greater than one. Equivalently, f is primitive if the map on homology $f_*: H_1(X; \mathbb{Z}) \rightarrow H_1(E; \mathbb{Z})$ is surjective.

An Abelian differential (X, ω) is called an *elliptic differential* if it is the pullback of a nonzero Abelian differential on an elliptic curve by a primitive branched cover. The *degree* of an elliptic differential is the degree of the cover.

Proposition 4.5 ([McMa, Theorem 4.10]). *The locus of degree d elliptic differentials in $\Omega\mathcal{M}_2$ is exactly ΩE_{d^2} .*

There is a special class of elliptic differentials called *square-tiled surfaces*. A square-tiled surface is an Abelian differential which is pulled back from the square elliptic curve $(\mathbb{C}/\mathbb{Z}[i], dz)$ by some (not necessarily primitive) cover branched only over 0. Equivalently, an Abelian differential is square-tiled if and only if its absolute and relative periods all lie in the Gaussian integers $\mathbb{Z}[i]$. All square-tiled surfaces lie on Teichmüller curves.

Weierstrass curves. A genus two Abelian differential is a *Weierstrass form* if it is an eigenform for real multiplication by some quadratic order $\mathcal{O}_D$ and if it has a double zero. Its *discriminant* is the discriminant D of the order $\mathcal{O}_D$. The Weierstrass forms are parameterized by the *Weierstrass form bundle* ΩW_D , which is a line bundle over the *Weierstrass curve*,

$$W_D := \mathbb{P}\Omega W_D \subset X_D.$$

Theorem 4.6 ([McM05a]). *W_D is a union of Teichmüller curves which is nonempty if and only if $D \geq 5$. If $D = 9$ or if $D \not\equiv 1 \pmod{8}$, then W_D is connected. Otherwise W_D has two connected components.*

Remark. When $D = p^2$ for some prime p , this statement was first proved in [HL06].

When W_D has two connected components, they are denoted by W_D^0 and W_D^1 . They are distinguished by a topological invariant called the *spin invariant*. This is easy to describe when $D = d^2$. In that case, an Abelian differential $(X, \omega) \in \Omega W_{d^2}$ is branched over an elliptic curve (E, ν) by a d -fold branched cover $f: X \rightarrow E$. Let p be the unique zero of ω , one of the six Weierstrass points of X . Of the six Weierstrass points, N of them have the same image in E as p , with either $N = 1$ or $N = 3$. If $N = 1$, then (X, ω) lies in ΩW_D^0 , and if $N = 3$, then (X, ω) lies in ΩW_D^1 .

Let $\widetilde{W}_D$ be the inverse image of W_D in the universal cover $\mathbb{H} \times \mathbb{H}$ of X_D .

Proposition 4.7 ([McM03a]). *$\widetilde{W}_D$ is a countable union of graphs of transcendental holomorphic maps $\mathbb{H} \rightarrow \mathbb{H}$.*

Equivalently, W_D is transverse to the absolute period foliation $\mathcal{A}_D$ of X_D introduced in §2.3.

Period coordinates for ΩE_D . Let $E_D(1, 1) = E_D \setminus W_D$, and let $\Omega E_D(1, 1)$ be the bundle of nonzero eigenforms. We can define period coordinates on $\Omega E_D(1, 1)$ in the same way as we defined period coordinates on the strata $\Omega \mathcal{M}_g(\mathbf{n})$. This material will only be used in §15.

Let

$$\tilde{E}_D(1, 1) = \mathbb{H} \times \mathbb{H} \setminus \widetilde{W}_D,$$

the inverse image of $E_D(1, 1)$ in the universal cover of X_D , and let $\Omega \tilde{E}_D(1, 1)$ be the (trivial) bundle of ι_1 -eigenforms. Given an $(X, \omega) \in \Omega \tilde{E}_D(1, 1)$, the homology group $H_1(X, Z(\omega); \mathbb{Z})$ contains $H_1(X; \mathbb{Z})$ as a subgroup which is equipped with an isomorphism

$$\mathcal{O}_D \oplus \mathcal{O}_D^\vee \rightarrow H_1(X; \mathbb{Z}).$$

The condition that ω is an eigenform is equivalent to the period map

$$P_\omega : H_1(X; \mathbb{Z}) \rightarrow \mathbb{C}$$

being $\mathcal{O}_D$ -linear, with $\mathcal{O}_D$ acting on $\mathbb{C}$ via the embedding ι_1 . Let

$$H_{\mathcal{O}_D}^1(X, Z(\omega); \mathbb{C})$$

be the subspace of $H^1(X, Z(\omega); \mathbb{C})$ consisting of all linear maps

$$H_1(X, Z(\omega); \mathbb{Z}) \rightarrow \mathbb{C}$$

that are $\mathcal{O}_D$ -linear on $H_1(X; \mathbb{Z})$.

Now consider $(X, \omega) \in \Omega \tilde{E}_D(1, 1)$. For a sufficiently small neighborhood U of (X, ω) , any $(X', \omega') \in U$ defines an element of $H_{\mathcal{O}_D}^1(X, Z(\omega); \mathbb{C})$ by composing the period map $P_{\omega'}$ with the isomorphism coming from the Gauss-Manin connection,

$$H_1(X, Z(\omega); \mathbb{Z}) \rightarrow H_1(X', Z(\omega'); \mathbb{Z}),$$

which is $\mathcal{O}_D$ -linear on $H_1(X; \mathbb{Z})$. This gives biholomorphic period coordinates

$$\phi : U \rightarrow H_{\mathcal{O}_D}^1(X, Z(\omega); \mathbb{C}).$$

Consider a triple $(\alpha, \beta, \gamma) \subset H_1(X, Z(\omega); \mathbb{Z})$ such that:

- (α, β) is a basis of $H_1(X; \mathbb{Z})$ over $\mathcal{O}_D$.
- $H_1(X; \mathbb{Z}) \oplus \langle \gamma \rangle = H_1(X, Z(\omega); \mathbb{Z})$.

Such a triple determines an isomorphism

$$H_1(X, Z(\omega); \mathbb{Z}) \rightarrow \mathbb{C}^3.$$

Two such triples are related by a matrix in the group,

$$G = \left\{ \begin{pmatrix} A & B \\ 0 & C \end{pmatrix} \in \mathrm{GL}_3 K(\mathcal{O}_D) : A \in \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee), B \in \mathcal{O}_D \oplus \mathcal{O}_D^\vee, C = \pm 1 \right\}.$$

Changing the triple by a matrix in G changes the period coordinates by the transpose of this matrix. This gives $\Omega\tilde{E}_D(1,1)$ the structure of an $(G, \mathbb{C}^3)$ -manifold, and $\Omega E_D(1,1)$ inherits the structure of a $(G, \mathbb{C}^3)$ -orbifold.

These period coordinates are compatible with the action of $\mathrm{SL}_2\mathbb{R}$ in the sense that $\phi: U \rightarrow \mathbb{C}^3$ commutes with the two $\mathrm{SL}_2\mathbb{R}$ actions, where $\mathrm{SL}_2\mathbb{R}$ acts on $\mathbb{C}$ coordinate-wise by identifying each $\mathbb{C}$ factor with $\mathbb{R}^2$.

By pulling back Lebesgue measure on $\mathbb{C}^3$ via these charts, we define a measure μ_D on $E_D(1,1)$. We can use this measure to define a measure μ'_D on $\Omega_1 E_D(1,1)$, using the same trick we used to define the measures $\mu'(\mathbf{n})$ on p. 47.

Theorem 4.8 ([McMa]). *The measures μ'_D are finite, ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measures.*

Cusps of W_D . We describe here McMullen's classification of cusps of W_D from [McM05a], which will play a crucial role in our calculation of $\chi(W_D^c)$.

Given an Abelian differential $(X, \omega) \in \Omega_1 \mathcal{M}_g$ and a slope $s \in \mathbb{P}^1(\mathbb{R})$, the foliation $\mathcal{F}_s$ of X is *completely periodic* if every leaf of the foliation is either a saddle connection joining zeros of ω , or a closed loop on X . A completely periodic foliation divides X into finitely many maximal cylinders C_i foliated by closed leaves of $\mathcal{F}_s$. The complement of $\bigcup C_i$ is a union of saddle connections on X called the spine of (X, ω) .

Two completely periodic foliations $\mathcal{F}_{s_i}$ of $(X_i, \omega_i) \in \Omega_1 \mathcal{M}_g$ are *equivalent* if there is some $A \in \mathrm{SL}_2\mathbb{R}$ such that $A \cdot (X_1, \omega_1) = (X_2, \omega_2)$ and $A \cdot s_1 = s_2$.

Now restrict to the case of Weierstrass forms $(X, \omega) \in W_D$. If D is not square, then a completely periodic foliation decomposes X into two cylinders; if D is square, then a completely periodic foliation decomposes X into either one or two cylinders (see [McM05a, Theorem 4.3]).

A Weierstrass form $(X, \omega) \in \Omega_1 W_D$ together with a completely periodic foliation $\mathcal{F}_s$ determines a cusp of W_D . Let $N \subset \mathrm{SL}_2\mathbb{R}$ be the upper-triangular subgroup consisting of all

$$N_t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}.$$

Let $g \in \mathrm{SL}_2\mathbb{R}$ be some matrix taking s to 0 (so that the horizontal foliation of $g \cdot (X, \omega)$ is completely periodic). The map

$$t \mapsto g^{-1} N_t g$$

defines a path on $\Omega_1 W_D$, which happens to cover a closed horocycle h on W_D . Replacing (X, ω, s) with an equivalent completely periodic foliation gives a horocycle homotopic to h . Homotopy classes of closed horocycles on a Riemann surface S correspond to cusps of S , so this construction associates a cusp of W_D to every equivalence class of completely periodic foliations. This correspondence is in fact a bijection; see, for example, [McM05a, Theorem 4.1].

We say that a cusp of W_D is a *one-cylinder cusp* or a *two-cylinder cusp* if the associated completely periodic foliation has one cylinder or two cylinders respectively. One cylinder cusps only arise on W_D if D is square. An example

of a foliation associated to a one-cylinder cusp is given by gluing the edges of a three-by-one rectangle as in Figure 4 and taking the horizontal foliation.




Figure 4: One cylinder cusp of W_9

McMullen [McM05a] classified the cusps of W_D by identifying them with splitting prototypes. We will describe his classification here using the equivalent W_D -prototypes from §3.

To the prototype $P = (a, b, c, \bar{q})$, we associate the surface (X_P, ω_P) formed by gluing a square in $\mathbb{C}$ with unit length sides to the parallelogram with sides 0 , $\lambda = \lambda(P)$, r , and $\lambda + r$, where λ is the unique positive root of

$$a\lambda^2 + b\lambda + c = 0,$$

and

$$r = -\frac{q}{c}\lambda - i\frac{a}{c}\lambda,$$

and then gluing opposite sides of the resulting polygon (see Figure 5). This (X_P, ω_P) is an eigenform for real multiplication by $\mathcal{O}_D$ where D is the discriminant of P . The horizontal foliation of (X_P, ω_P) is completely periodic and so




Figure 5: Cusp of W_D

determines a cusp w_P of W_D .

Theorem 4.9 ([McM05a, Theorem 4.1]). *The map $P \mapsto w_P$ described above determines a bijection between the set of W_D -prototypes and the set of two-cylinder cusps of W_D .*

We will also need to know which connected component of W_D contains a given cusp w_P . Given an order $\mathcal{O}_D$, define the *conductor* of $\mathcal{O}_D$ to be the integer f such that $D = f^2E$ with E a fundamental discriminant.

Theorem 4.10 ([McM05a, Theorem 5.3]). *The cusp w_P of W_D associated to the W_D -prototype $P = (a, b, c, \bar{q})$ is contained in the component $W_D^{\epsilon(P)}$, where*

$$\epsilon(P) \equiv \frac{b-f}{2} + (a+1)(q+c+qc) \pmod{2}, \quad (4.6)$$

and f is the conductor of $\mathcal{O}_D$.

Other Teichmüller curves. When the discriminant D is square, there is an infinite family of Teichmüller curves on X_{d^2} parameterizing Abelian differentials with two simple zeros. Given $(X, \omega) \in \Omega X_{d^2}$, let $f: (X, \omega) \rightarrow (E, \nu)$ its associated degree d torus cover. Define $\Omega W_{d^2}[n]$ to be the locus of all differentials such that the two branch points of f in E differ by torsion of degree exactly n in the group law on E , and define $W_{d^2}[n] \subset X_{d^2}$ be its projectivization. The locus $W_{d^2}[n]$ is a union of Teichmüller curves. Note that $W_{d^2}[0]$ contains W_{d^2} but is in general larger because X can have distinct zeros which are branched over the same point of E .

There is one more example of a genus two Teichmüller curve which comes from an infinite family discovered by Veech. For even n , consider the Abelian differential obtained by gluing opposite sides of the regular n -gon. Veech [Vee92] showed that the $\mathrm{SL}_2\mathbb{R}$ orbit of this differential lies on a Teichmüller curve D_n . The Abelian differential coming from the decagon is an genus two Abelian differential with two simple zeros. Its orbit is a Teichmüller D_{10} curve lying on X_5 .

Classification of $\mathrm{SL}_2\mathbb{R}$ orbit closures in genus two. It happens that every $\mathrm{SL}_2\mathbb{R}$ orbit-closure on $\Omega_1\mathcal{M}_2$ or ergodic, invariant measure is one of the ones just described.

Theorem 4.11 ([McMa], [McM06]). *Every closure of an $\mathrm{SL}_2\mathbb{R}$ orbit in $\Omega_1\mathcal{M}_2$ is either all of $\Omega_1\mathcal{M}_2$, or one of the following submanifolds: $\Omega_1\mathcal{M}_2(2)$, $\Omega_1 E_D$, $\Omega_1 D_{10}$, a connected component of $\Omega_1 W_D$, or a component of $\Omega_1 W_{d^2}[n]$.*

Furthermore each of these orbit closure carries a unique ergodic, absolutely continuous, $\mathrm{SL}_2\mathbb{R}$ -invariant probability measure. These are all of the ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant probability measures on $\Omega_1\mathcal{M}_2$.

The only gap remaining in this classification is to describe the connected components of $W_{d^2}[n]$.

5 Deligne-Mumford compactification of moduli space

5.1 Stable Riemann surfaces

A *nodal Riemann surface* is a connected, compact, one-dimensional, complex analytic space with only nodes as singularities (a node is a transverse crossing of two nonsingular branches). Equivalently, a nodal Riemann surface can be regarded as a finite type Riemann surface with finitely many cusps which have been identified pairwise to form nodes. A connected component of a nodal Riemann surface X with its nodes removed is called a *part* of X , and the closure of a part of X is an *irreducible component* of X . In this paper the *genus* of a nodal Riemann surface X will mean its arithmetic genus,

$$g = 1 - \chi(\mathcal{O}_X),$$

where $\mathcal{O}_X$ is the structure sheaf of X . In topological terms, the arithmetic genus of a nodal Riemann surface X is the genus of the nonsingular Riemann surface obtained by replacing each node of X with an annulus.

A *stable Riemann surface* is a connected nodal Riemann surface for which each part has nonabelian fundamental group (or equivalently negative Euler characteristic).

A nodal Riemann surface X has a *normalization* $\tilde{X} \rightarrow X$ defined by separating the two branches passing through each node of X .

Stable Abelian differentials. A *stable Abelian differential* on a stable Riemann surface X is a holomorphic 1-form on X minus its nodes such that:

- Its restriction to each part of X has at worst simple poles at the cusps.
- At two cusps which have been identified to form a node, the differential has opposite residues.

These properties can be conveniently rephrased using the normalization. If ω is a meromorphic Abelian differential on X , and $f: \tilde{X} \rightarrow X$ is the normalization, then ω is stable if and only if $f^*\omega$ has at worst simple poles, and for every $q \in X$,

$$\sum_{f(p)=q} \text{Res}_p(f^*\omega) = 0.$$

A stable Riemann surface X has a *dualizing sheaf* ω_X . A stable Abelian differential on a stable Riemann surface is just a global section of ω_X . This is discussed in [HM98] and [Har66].

The stable Abelian differentials on a genus g singular Riemann surface X form a complex vector space which we will write as $\Omega(X)$. The complex dimen-

sion of $\Omega(X)$ is g :

$$\begin{aligned} h^0(X, \omega_X) &= h^1(X, \mathcal{O}_X) \\ &= 1 - \chi(\mathcal{O}_X) \\ &= g. \end{aligned}$$

This fact also follows easily from the Riemann-Roch Theorem. It is also proved in [Ser59, Theorem IV.2].

We will often use the term “Abelian differential” as shorthand for “Riemann surface together with an Abelian differential”. We will call a node of a stable Abelian differential (X, ω) a *polar node* if ω has a pole there and a *holomorphic node* otherwise.

Jacobians. The classical notion of the Jacobian variety of a nonsingular Riemann surface can be extended to singular Riemann surfaces.

If X is a possibly singular Riemann surface, let X_0 be the set of nonsingular points of X . Then there is a natural map $H_1(X_0; \mathbb{Z}) \rightarrow \Omega^*(X)$, given by integrating forms over homology classes. The *Jacobian variety* of X is the variety,

$$\text{Jac}(X) = \Omega^*(X) / H_1(X_0; \mathbb{Z}).$$

In the case of a stable Riemann surface X , it is easy to describe the kernel of the map $H_1(X_0; \mathbb{Z}) \rightarrow \Omega^*(X)$. It is the subgroup of $H_1(X_0; \mathbb{Z})$ generated by the relations $\alpha - \beta$ when α and β are homology classes generated by curves going around the same node of X in the “same direction” on opposite parts of X as in Figure 6.




Figure 6: Curves around a node.

There is an exact sequence which relates the Jacobian of a nodal Riemann surface to the Jacobian of its normalization. If $\tilde{X} \rightarrow X$ is the normalization of X , then holomorphic Abelian differentials on $\tilde{X}$ restrict to stable Abelian differentials on X , so there is a natural map $\text{Jac}(X) \rightarrow \text{Jac}(\tilde{X})$. We get an exact sequence:

$$0 \rightarrow (\mathbb{C}^*)^n \rightarrow \text{Jac}(X) \rightarrow \text{Jac}(\tilde{X}) \rightarrow 0,$$

where n is the difference of the genera of X and $\tilde{X}$. This realizes $\text{Jac}(X)$ as a *semi-Abelian variety*.

For example, if X has genus two and one separating node, then $\tilde{X}$ is the disjoint union of two elliptic curves. The Jacobian of X is then the product of these two elliptic curves.

If X has genus two and one nonseparating node, then the normalization of X is an elliptic curve E . The Jacobian of X is then an extension of E by $\mathbb{C}^*$. The original stable Riemann surface X can be recovered from $\text{Jac}(X)$.

If X has genus two and two (or three) nonseparating nodes, then $\tilde{X}$ is $\mathbb{P}^1$ (or $\mathbb{P}^1 \cup \mathbb{P}^1$ respectively), so $\text{Jac}(X) \cong \mathbb{C}^* \times \mathbb{C}^*$. This example shows that two distinct nodal Riemann surfaces may have the same Jacobian.

The Jacobian of X can also be identified with $\text{Pic}^0(X)$, the group of all line bundles on X which have degree zero on each irreducible component of X [HM98, p.250].

5.2 Deligne-Mumford compactification

Marked stable Riemann surfaces. Given a topological stable surface X of genus g , a *collapse* of Σ_g onto X is a continuous surjection $f: \Sigma_g \rightarrow X$ with the following properties.

- The inverse image of each node of X is a Jordan curve on Σ_g .
- Each component of $\Sigma \setminus f^{-1}(N)$, where N is the set of nodes of X , maps homeomorphically, preserving the orientation, onto a part of X .

If X is nonsingular, a collapse is just a homeomorphism $\Sigma_g \rightarrow X$. These maps were introduced by Bers; in his terminology, a collapse is called a strong deformation.

A *marked stable Riemann surface* is a stable Riemann surface X , together with a collapse $\Sigma_g \rightarrow X$. Two markings $f_i: \Sigma_g \rightarrow X_i$ are *equivalent* if there is a conformal isomorphism $g: X_1 \rightarrow X_2$ such that the following diagram commutes up to homotopy.

$$\begin{array}{ccc} \Sigma_g & \xrightarrow{f_1} & X_1 \\ & \searrow f_2 & \downarrow g \\ & & X_2 \end{array}$$

We will sometimes denote by $[f: \Sigma_g \rightarrow X]$ the class of all marked surfaces equivalent to $f: \Sigma_g \rightarrow X$.

Augmented Teichmüller space. The Teichmüller space $\mathcal{T}(\Sigma_g)$ is contained in the *Augmented Teichmüller space*, $\overline{\mathcal{T}}(\Sigma_g)$, the set of all marked stable Riemann surfaces up to equivalence. Let $\partial\mathcal{T}(\Sigma_g) = \overline{\mathcal{T}}(\Sigma_g) \setminus \mathcal{T}(\Sigma_g)$.

We give $\overline{\mathcal{T}}(\Sigma_g)$ a topology as follows. Given a closed curve γ on Σ_g , define a function $l_\gamma: \overline{\mathcal{T}}(\Sigma_g) \rightarrow \mathbb{R} \cup \{\infty\}$: if γ is not homotopic to a curve on X disjoint

from the nodes, let $l_\gamma(X) = \infty$; if γ is homotopic to a node of X , let $l_\gamma(X) = 0$; otherwise let $l_\gamma(X)$ be the length of the unique geodesic homotopic to γ in the Poincaré metric on X minus its nodes. Give $\overline{\mathcal{T}}(\Sigma_g)$ the smallest topology such that l_γ is continuous for every closed curve γ on Σ_g . The induced subspace topology on $\mathcal{T}(\Sigma_g)$ agrees with that defined by the Teichmüller metric on $\mathcal{T}(\Sigma_g)$.

Abikoff [Abi77] showed that this topology is equivalent to other natural topologies on $\overline{\mathcal{T}}(\Sigma_g)$, such as those obtained by looking at quasiconformal maps or quasi-isometries defined outside a neighborhood of the nodes.

Another useful topology which is equivalent to this one is the *conformal topology* which is defined as follows. Let $[f: \Sigma_g \rightarrow X] \in \partial\mathcal{T}(\Sigma_g)$, and let $U \subset X$ be a neighborhood of the nodes of X . Let V_U be the set of all $[g: \Sigma_g \rightarrow Y] \in \overline{\mathcal{T}}(\Sigma_g)$ for which f and g can be adjusted by homotopies so that $g \circ f^{-1}|_{X \setminus \overline{U}}$ is conformal. The set of all V_U as U runs over all neighborhoods of the nodes of X define a neighborhood basis of X . Together with the open sets of $\mathcal{T}(\Sigma_g)$, this defines a topology on $\overline{\mathcal{T}}(\Sigma_g)$. It is well-known that this topology is equivalent to the one defined above; however, a proof does not exist in the literature.

A *curve system* on Σ_g is a collection of simple closed curves on Σ_g , none of which are isotopic to any other or to a point. For each curve system S , there is a subspace $\mathcal{T}(\Sigma_g, S) \subset \partial\mathcal{T}(\Sigma_g)$ consisting of marked stable Riemann surfaces $[f: \Sigma_g \rightarrow X]$ which collapse each curve of a curve system homotopic to S to a point. For each connected component Σ_g^i of $\Sigma_g \setminus S$, let S_i be the closed surface with marked points obtained by collapsing the boundary components of Σ_g^i to points and regarding the images of the boundary components as marked points. There is a natural isomorphism,

$$\mathcal{T}(\Sigma_g, S) \cong \prod_i \mathcal{T}(S_i),$$

Deligne-Mumford compactification. The action of $\text{Mod}(\Sigma_g)$ of $\mathcal{T}(\Sigma_g)$ extends to an action on $\overline{\mathcal{T}}(\Sigma_g)$. The quotient,

$$\overline{\mathcal{M}}_g = \overline{\mathcal{T}}(\Sigma_g) / \text{Mod}(\Sigma_g),$$

is the Deligne-Mumford compactification of moduli space. It is a compact orbifold whose points naturally parameterize stable Riemann surfaces of genus g .

Given a curve system $S \subset \Sigma_g$, let $\mathcal{M}_g(S) \subset \overline{\mathcal{M}}_g$ be the stratum of stable Riemann surfaces homeomorphic to the topological stable surface Σ_g/S . This is a locally closed subset of $\overline{\mathcal{M}}_g$, and $\overline{\mathcal{M}}_g$ is the disjoint union of all of the $\mathcal{M}_g(S)$ as S ranges over all isotopy classes of curve systems on Σ_g up to the action of the modular group.

Dehn space. Given a curve system S on Σ_g , let $\text{Tw}(S)$ be the group generated by Dehn twists around the curves of Σ . It is an Abelian group isomorphic to $\mathbb{Z}^h$, where h is the number of curves in S . Define the *Dehn space* $\mathcal{D}(\Sigma_g, S)$ to be

$$\mathcal{D}(\Sigma_g, S) = (\mathcal{T}(\Sigma_g) \cup \mathcal{T}(\Sigma_g, S)) / \text{Tw}(S).$$

We will sometimes use the notation $\mathcal{D}_g(S) = \mathcal{D}(\Sigma_g, S)$ when we don't need to emphasize the surface Σ_g . Bers called $\mathcal{D}(\Sigma_g, S)$ the strong deformation space.

Let $\text{Stab}(S)$ be the subgroup of $\text{Mod}(\Sigma_g)$ which maps S to an isotopic curve system. $\text{Tw}(S)$ is a normal subgroup of $\text{Stab}(S)$, and the quotient,

$$\text{Mod}(\Sigma_g, S) = \text{Stab}(S) / \text{Tw}(S),$$

is the mapping class group of the topological stable surface obtained by collapsing each curve in S to a point.

The group $\text{Mod}(\Sigma_g, S)$ acts on $\mathcal{D}(\Sigma_g, S)$, and the natural map

$$\pi: \mathcal{D}(\Sigma_g, S) \rightarrow \overline{\mathcal{M}}_g$$

is invariant under this action. Bers showed:

Proposition 5.1 ([Ber74, p. 1221]). *If $X \in \mathcal{D}(\Sigma_g, S)$, and $G \subset \text{Mod}(\Sigma_g, S)$ is the stabilizer of X , then there is a neighborhood $U \subset \mathcal{D}(\Sigma_g, S)$ of X , stable under the action of G such that π factors through to a map, $\bar{\pi}: U/G \rightarrow \overline{\mathcal{M}}_g$, which is homeomorphic onto its image.*

Complex structure. We now describe the complex structure on $\overline{\mathcal{M}}_g$, following Bers's approach [Ber81].

In [Ber61a], Bers defined an embedding $B: \mathcal{T}(\Sigma_g) \rightarrow \mathbb{C}^n$, where

$$n = \begin{cases} 1 & \text{if } g = 1; \\ 3g - 3 & \text{if } g > 1. \end{cases}$$

It is a homeomorphism onto a bounded domain, and $\mathcal{T}(\Sigma_g)$ inherits the complex structure on $\mathbb{C}^n$. Ahlfors [Ahl60] showed that this is the unique complex structure on $\mathcal{T}(\Sigma_g)$ for which the periods of Abelian differentials vary holomorphically.

The modular group $\text{Mod}(\Sigma_g)$ acts biholomorphically on $\mathcal{T}(\Sigma_g)$ with this complex structure, so $\mathcal{M}_g$ inherits the structure of a complex orbifold.

Bers gives $\overline{\mathcal{M}}_g$ a complex structure by first giving $\mathcal{D}(\Sigma_g, S)$ a complex structure for each curve system S . First define a sheaf of rings $\mathcal{O}$ on $\mathcal{D}(\Sigma_g, S)$ to be the sheaf consisting of all continuous functions $\mathcal{D}(\Sigma_g, S) \rightarrow \mathbb{C}$ which are holomorphic on

$$\mathcal{D}_0(\Sigma_g, S) = \mathcal{T}(\Sigma_g) / \text{Tw}(S) \subset \mathcal{D}(\Sigma_g, S)$$

with respect to the complex structure on $\mathcal{D}_0(\Sigma_g, S)$ induced by the Bers embedding of Teichmüller space. Given a domain, $U \subset \mathcal{D}(\Sigma_g, S)$, we consider a function $f: U \rightarrow \mathbb{C}$ to be holomorphic if it is in $\mathcal{O}$. Bers showed that this defines an integrable complex structure on $\mathcal{D}(\Sigma_g, S)$ by giving a biholomorphic isomorphism of $\mathcal{D}(\Sigma_g, S)$ with a bounded domain in $\mathbb{C}^n$ parameterizing a certain family of Kleinian groups:

Proposition 5.2 ([Ber81]). *Let $S = \bigcup S_i$ be a curve system on Σ_g . There is a biholomorphic map*

$$B: \mathcal{D}(\Sigma_g, S) \rightarrow \mathbb{C}^n$$

onto a bounded domain in $\mathbb{C}^n$. A curve S_i is homotopic to a node on a marked stable Riemann surface $X \in \mathcal{D}(\Sigma_g, S)$ if and only if

$$z_i(B(X)) = 0,$$

where the z_i are the coordinates on $\mathbb{C}^n$.

Similar embeddings of $\mathcal{D}(\Sigma_g, S)$ are constructed in [Mar87] and [Kra90].

Using Proposition 5.1, we give $\overline{\mathcal{M}}_g$ the unique complex structure which makes all of the maps $\mathcal{D}(\Sigma_g, S) \rightarrow \overline{\mathcal{M}}_g$ holomorphic.

Baily showed in [Bai60, Bai62] that $\mathcal{M}_g$ also has the structure of a quasi-projective variety. Wolpert [Wol85] showed using the Weil-Petersson metric on $\mathcal{M}_g$ that $\overline{\mathcal{M}}_g$ has the structure of a projective variety. Alternatively, Deligne-Mumford [DM69, Mum71] constructed $\overline{\mathcal{M}}_g$ algebraically and showed that it is a coarse moduli space for stable genus g curves, and Knudsen-Mumford [KM76] showed that $\overline{\mathcal{M}}_g$ is a projective variety.

Jacobians in genus two. By associating a Riemann surface to its Jacobian, there is a natural morphism $\text{Jac}: \mathcal{M}_g \rightarrow \mathcal{A}_g$.

Theorem 5.3 ([Nam73]). *The morphism Jac extends to a morphism*

$$\overline{\text{Jac}}: \overline{\mathcal{M}}_g \rightarrow \widehat{\mathcal{A}}_g,$$

which sends a Riemann surface X to the Jacobian of the normalization of X .

Determining the image of Jac map is in general very difficult; however, this is simple in genus two. Let $\widetilde{\mathcal{M}}_2$ be the subvariety of $\overline{\mathcal{M}}_2$ consisting of nonsingular Riemann surfaces together with pairs of elliptic curves joined at a single node. The following proposition is well known.

Proposition 5.4. *The image of $\text{Jac}: \mathcal{M}_2 \rightarrow \mathcal{A}_2$ is exactly those Abelian varieties which are not polarized products of elliptic curves. Furthermore, Jac extends to an isomorphism $\widetilde{\text{Jac}}: \widetilde{\mathcal{M}}_2 \rightarrow \mathcal{A}_2$.*

Sketch of proof. The image of $\overline{\text{Jac}}$ contains an open set because these varieties have the same dimension, and Jac restricted to $\mathcal{M}_2$ is injective by the Torelli theorem. Any map between complete irreducible varieties of the same dimension whose image contains an open set is onto, so $\overline{\text{Jac}}$ is onto.

It's easy to check that the Jacobian of a genus two Riemann surface X can't be the polarized product of two elliptic curves (for example this would give a degree one map of X to an elliptic curve), so the image of $\mathcal{M}_2$ is exactly the complement of the locus of polarized products.

The map $\widetilde{\text{Jac}}$ is also injective (since the Jacobian of a Riemann surface X formed from two elliptic curves joined at a node is X itself), so it is a bijection. It follows that $\widetilde{\text{Jac}}$ is an isomorphism because it is a bijection between two normal varieties. ■

Abelian differentials over $\overline{\mathcal{M}}_g$. The bundle $\Omega\mathcal{T}(\Sigma_g)$ extends to a trivial bundle $\Omega\overline{\mathcal{T}}(\Sigma_g)$ over $\overline{\mathcal{T}}(\Sigma_g)$. The quotient,

$$\Omega\overline{\mathcal{M}}_g = \Omega\overline{\mathcal{T}}(\Sigma_g) / \text{Mod}(\Sigma_g),$$

is the moduli space of stable Abelian differentials of genus g . It is an orbifold vector bundle over $\overline{\mathcal{M}}_g$ whose fiber over X is $(\Omega(X) \setminus \{0\}) / \text{Aut}(X)$.

Similarly, define the bundle $\Omega\mathcal{D}(\Sigma_g, S)$ to be the restriction of

$$\Omega\overline{\mathcal{T}}(\Sigma_g, S) / \text{Tw}(S)$$

to $\mathcal{D}(\Sigma_g, S)$. This is a trivial bundle whose fiber over a marked, stable Abelian differential (f, X) is $\Omega(X)$. To see that it is trivial, choose a Lagrangian subspace $\Lambda \subset H_1(\Sigma_g; \mathbb{Z})$ which is orthogonal to the homology class of each of the curves in S . Then Λ is fixed pointwise by the action of $\text{Tw}(S)$, and

$$\Omega\mathcal{D}(\Sigma_g, S) \cong \mathcal{D}(\Sigma_g, S) \times \text{Hom}(\Lambda, \mathbb{C}).$$

5.3 Degenerating Abelian differentials

We now study in more detail the geometry of Abelian differentials as they approach the boundary of moduli space. For example, we will see that Abelian differentials close to a stable Abelian differential (X, ω) with a polar node develop very long cylinders which are pinched off to form a node in the limit. We will then use our understanding of this geometry to study the behavior of Jacobians of Riemann surfaces near the boundary of moduli space. Finally, we will give two applications concerning real multiplication which will be used in §7.

Long cylinders. The conformal topology on $\overline{\mathcal{T}}(\Sigma_g)$ defined on p. 57 is compatible with the topology on the trivial bundle $\Omega\overline{\mathcal{T}}(\Sigma_g)$ in the following sense. Consider $[f: \Sigma_g \rightarrow X] \in \partial\mathcal{T}(\Sigma_g)$ with $\omega \in \Omega(X)$. Let $U \subset X$ be a neighborhood of the nodes of X , and let $K > 1$. Let $V_{U,K}$ be the set of all $([g: \Sigma_g \rightarrow Y], \eta) \in \Omega\overline{\mathcal{T}}(\Sigma_g)$ for which f and g can be adjusted by homotopies so that $g \circ f^{-1}|_{X \setminus \overline{U}}$ is conformal, and

$$K^{-1} < \left| \frac{(g \circ f^{-1})^* \eta}{\omega} \right| < K$$

on $X \setminus \overline{U}$. It is an unpublished result that the $V_{U,K}$ are a neighborhood basis of $([f: \Sigma_g \rightarrow X], \omega)$ in $\Omega\overline{\mathcal{T}}(\Sigma_g)$.

We can use this description of the topology on $\Omega\overline{\mathcal{T}}(\Sigma_g)$ to prove the following theorem which gives a more precise description of the shape of an Abelian differential close to a stable Abelian differential (X, ω) with $X \in \partial\mathcal{M}_2$.

Theorem 5.5. *Let $(X, \omega) \in \Omega\overline{\mathcal{T}}(\Sigma_g)$ with $X \in \overline{\mathcal{T}}(\Sigma_g, S)$, and let $C \subset X$ be a compact subset disjoint from the nodes. For any $\epsilon, h > 0$ and $K > 1$, there is a neighborhood U of (X, ω) in $\Omega\overline{\mathcal{T}}(\Sigma_g)$ such that each $(Y, \eta) \in U$ has the following properties.*

- The collapse $Y \rightarrow X$ induced by the markings is homotopic to a collapse $f: Y \rightarrow X$ which is conformal on C and satisfies

$$K^{-1} < \left| \frac{f^* \omega}{\eta} \right| < K \quad (5.1)$$

on C .

- Each curve in S which represents a polar node of (X, ω) is homotopic to a cylinder on Y of height at least h .
- When $g = 2$, for each curve γ in S which represents a polar node of X , if

$$\int_{\gamma} \eta = 0. \quad (5.2)$$

then there are saddle connections I_1 and I_2 on (Y, η) joining two distinct zeros of η of length less than ϵ such that γ is homotopic to $I_1 \cup I_2$ and (Y, η) can be split along $I_1 \cup I_2$.

Proof. The first statement follows directly from the definition of the conformal topology on $\Omega\overline{T}(\Sigma_g)$ above.

Now let p be a polar node of (X, ω) . By possibly enlarging the compact set C , we can suppose that C contains a cylinder D of height $2h$ around p . If $f: Y \rightarrow X$ is a collapse satisfying (5.1) with $K < 2$, then the two boundary components of $f^{-1}(D)$ are at least distance h apart. Furthermore, if K is sufficiently small, then the boundary components of $f^{-1}(D)$ both have winding number zero, so $f^{-1}(D)$ contains no zeros. Therefore $f^{-1}(D)$ contains a cylinder of height at least h , which proves the second statement.

Now suppose that p is a holomorphic node of (X, ω) and suppose that (5.2) holds. If ϵ is sufficiently small, then there is a neighborhood U of p isomorphic to two disks of radius ϵ/π joined at a point. By possibly enlarging the compact subset $C \subset X$, we have $\partial U \subset C$. If $([f: \Sigma_2 \rightarrow Y], \eta) \in V_{X \setminus C, K}$ with $K < 2$, then $f^{-1}(U)$ is bounded by two curves β_i of circumference less than 2ϵ . Furthermore, if K is sufficiently small, then the derivatives of f will be close to the identity, and we can assume that the curves β_i have winding number one and positive curvature. Since the β_i have winding number one, it follows from the Gauss-Bonnet Theorem that η has two zeros in $f^{-1}(U)$, counting multiplicity. Since the β_i have positive curvature, there is a shortest geodesic $\delta \in f^{-1}(U)$ generating $\pi_1(f^{-1}(U))$. By (5.2), we have

$$\int_{\delta} \eta = 0, \quad (5.3)$$

which implies that δ is not a straight geodesic, so it must be a union of saddle connections joining distinct zeros. As there are at most two zeros in $f^{-1}(U)$, the curve δ must be a union of two saddle connections I_1 and I_2 going from a zero q to a zero r of η . By (5.3), we have

$$\int_{I_1} \eta = \int_{I_2} \eta. \quad (5.4)$$

This means that we can split along $I_1 \cup I_2$ by the discussion of splitting along a union of saddle connections in §4.1. Finally, since δ is the shortest curve in its homotopy class in $f^{-1}(U)$ and the length of β_i is less than 2ϵ , the length of I_i must be less than ϵ as claimed. $\blacksquare$

Semi-Abelian varieties. We now introduce a topology on the space of semi-Abelian varieties and show that the map which associates to a marked stable Riemann surface its Jacobian is continuous.

A *semi complex torus* is a quotient $A = V/\Lambda$, where V is a complex vector space containing a discrete subgroup Λ , such that there is an exact sequence,

$$0 \rightarrow (\mathbb{C}^*)^n \rightarrow A \xrightarrow{\phi} B \rightarrow 0,$$

where $B = V'/\Lambda'$ is a compact complex torus. A principal polarization h' on B induces a degenerate Hermitian form h on V by pulling back h' by the lift $\tilde{\phi}: V \rightarrow V'$. Such a Hermitian form on V is a *principal polarization* on A .

Equip $\mathbb{Z}^{2g}$ with the usual symplectic form defined by (2.1). A *marked semi-Abelian variety* is a triple (A, Γ, ϕ) , where

- A is a semi-Abelian variety,
- $\Gamma \subset \mathbb{Z}^{2g}$ is a subgroup of $\mathbb{Z}^{2g}$ on which the symplectic form vanishes and which is saturated in the sense that if $x \in \mathbb{Z}^{2g}$ and $mx \in \Gamma$ for some $m \in \mathbb{Z}$, then $x \in \Gamma$, and
- ϕ is a symplectic isomorphism $\delta: \Gamma^\perp \rightarrow H_1(A; \mathbb{Z})$.

Let $\mathcal{H}_g^\dagger$ be the space of all marked semi-Abelian varieties. The subspace of $\mathcal{H}_g^\dagger$ consisting of marked Abelian varieties is naturally isomorphic to the Siegel upper half plane $\mathcal{H}_g$ by the discussion in §2.1.

We give $\mathcal{H}_g^\dagger$ a topology as follows. If

$$(A_n = V_n/\Lambda_n, \Gamma_n, \phi_n)$$

is a sequence in $\mathcal{H}_g^\dagger$, then we say that this sequence converges to $(A = V/\Lambda, \Gamma, \phi)$ if eventually $\Gamma_n \subset \Gamma$ and there are linear isomorphisms $\psi_n: V_n \rightarrow V$ with the following properties:

- For all $\alpha \in \Gamma^\perp$,
- $$\lim_{n \rightarrow \infty} \psi_n \circ \phi_n(\alpha) = \phi(\alpha),$$
- For all $\alpha \in \mathbb{Z}^{2g} \setminus \Gamma^\perp$, the sequence $\psi_n \circ \phi_n(\alpha)$ eventually leaves every compact subset of V .
 - If h_n and h are the polarizations on A_n and A , then

$$\lim_n (\phi_n)_*(h_n) = h$$

as Hermitian forms on V ,

This defines a Hausdorff topology on $\mathcal{H}_g^\dagger$.

Over $\mathcal{H}_g^\dagger$ is the trivial rank-two bundle $\Omega\mathcal{H}_h^\dagger$ consisting of pairs (A, ω) , where $A \in \mathcal{H}_g^\dagger$ and $\omega \in \Omega(A)$, the space of holomorphic one-forms on A .

A choice of a symplectic isomorphism $H_1(\Sigma_2; \mathbb{Z}) \rightarrow \mathbb{Z}^4$ defines natural maps $\text{Jac}: \overline{\mathcal{T}}(\Sigma_2) \rightarrow \mathcal{H}_2^\dagger$ and $\Omega \text{Jac}: \Omega\overline{\mathcal{T}}(\Sigma_2) \rightarrow \Omega\mathcal{H}_2^\dagger$, sending a marked stable Riemann surface to its Jacobian with the induced marking.

Theorem 5.6. *The maps Jac and ΩJac defined above are continuous.*

We will sketch the proof of this theorem below.

Given any $X \in \mathcal{T}(\Sigma_2, S)$, we can define a norm $\|\cdot\|_X$ on $H_1(\Sigma_2; \mathbb{R})$, the Hodge norm as follows. Let $V_S \subset H_1(\Sigma_2; \mathbb{R})$ be the subspace generated by the curves in S . If $\gamma \notin V_S^\perp \subset H_1(\Sigma_2; \mathbb{R})$, set $\|\gamma\|_X = \infty$. On $V_S^\perp$, let $\|\cdot\|_X$ be the norm induced by the Hermitian metric on $\Omega(X)^*$ coming from the polarization via the embedding $V_S^\perp \rightarrow \Omega(X)^*$. Alternatively, for $\gamma \in V_S^\perp$,

$$\|\gamma\|_X = \sup_{\substack{\omega \in \Omega(X) \\ \|\omega\|=1}} |\omega(\gamma)|,$$

where

$$\|\omega\| = \left(\int_X |\omega|^2 \right)^{1/2}.$$

Theorem 5.7. *Let $V_{S, \mathbb{Z}} \subset H_1(\Sigma_2; \mathbb{Z})$ be the subgroup generated by the curves in S . If $\{X_n\}$ is a sequence in $\overline{\mathcal{T}}(\Sigma_2)$ converging to X in $\mathcal{T}(\Sigma_2, S)$, then*

- *There exists a $C > 0$ such that $\|\gamma\|_{X_n} > C$ for all $\gamma \in V_{S, \mathbb{Z}}^\perp \setminus V_{S, \mathbb{Z}}$.*
- *For all $D > 0$, there exists $N > 0$ such that $\|\gamma\|_{X_n} > D$ for all $\gamma \in H_1(\Sigma_2; \mathbb{Z}) \setminus V_{S, \mathbb{Z}}^\perp$ if $n > N$.*

Define for any curve system S on Σ_2 ,

$$\overline{\mathcal{T}}(\Sigma_2, S) = \mathcal{T}(\Sigma_2) \cup \bigcup_{T \subset S} \mathcal{T}(\Sigma_2, T),$$

where the union is over all curve systems $T \subset S$. The proof of these theorems will rely on the following lemma.

Lemma 5.8. *Given any linear map $R: V_S \rightarrow \mathbb{R}$, there is a unique section $Z \mapsto \omega_X$ of $\Omega\overline{\mathcal{T}}(\Sigma_2, S)$ over $\overline{\mathcal{T}}(\Sigma_2, S)$ such that:*

- $\omega_X(\gamma) = R(\gamma)$ for all $\gamma \in V_S$, and
- $\text{Im } \omega_X(\gamma) = 0$ for all $\gamma \in V_S^\perp$.

Proof. Let $X \in \overline{\mathcal{T}}(\Sigma_2, S)$. We have an exact sequence,

$$0 \rightarrow \Omega(\tilde{X}) \rightarrow \Omega(X) \rightarrow \text{Hom}(V_S, \mathbb{C}) \rightarrow 0,$$

where $\tilde{X}$ is the normalization of X . Choose some $\eta \in \Omega(X)$ such that $\omega(\gamma) = R(\gamma)$ for each $\gamma \in V_S$. The form η defines a map,

$$S: V_S^\perp / V_S \rightarrow \mathbb{R},$$

by $S(\gamma) = \text{Im } \eta(\gamma)$. We have isomorphisms,

$$H_1(\tilde{X}; \mathbb{R}) \rightarrow V_S^\perp / V_S,$$

and

$$\Omega(\tilde{X}) \rightarrow \text{Hom}(H_1(\tilde{X}; \mathbb{R}), \mathbb{R}),$$

defined by $\omega \mapsto (\gamma \mapsto \text{Im } \omega(\gamma))$. Let $\nu \in \Omega(\tilde{X})$ induce the map S above. Then $\omega_X = \eta - \nu$ is the desired form, giving the desired section over $\overline{\mathcal{T}}(\Sigma_2, S)$. ■

Sketch of Proof of Theorem 5.6. Suppose we have a sequence $\{X_n\}$ in $\overline{\mathcal{T}}(\Sigma_2)$ converging to some $X \in \mathcal{T}(\Sigma_2, S)$. To prove that Jac is continuous, we need to show:

- For every section $Y \mapsto \omega_Y$ of $\overline{\Omega\mathcal{T}}(\Sigma_2, S)$ and $\gamma \in V_S^\perp$, we have

$$\omega_{X_n}(\gamma) \rightarrow \omega_X(\gamma),$$

and

- For every $\gamma \in H_1(\Sigma_2; \mathbb{R}) \setminus V_S^\perp$, there is a section $Y \mapsto \eta_Y$ of $\overline{\Omega\mathcal{T}}(\Sigma_2, S)$ such that $|\eta_{X_n}(\gamma)| \rightarrow \infty$.

We should also show that the polarizations converge, but we will omit the proof of this.

If $X \mapsto \omega_X$ is a section of $\overline{\Omega\mathcal{T}}(\Sigma_2, S)$, then ω_{X_n} converges uniformly to ω_X away from the nodes of X as in Theorem 5.5. If $\gamma \in V_S^\perp$, then γ is represented by a curve on X which is disjoint from the nodes, so $\omega_{X_n}(\gamma) \rightarrow \omega(\gamma)$ by uniform convergence.

Suppose $\gamma \in H_1(\Sigma_2; \mathbb{R}) \setminus V_S^\perp$. By Lemma 5.8, there is a section $X \mapsto \eta_X$ of $\overline{\Omega\mathcal{T}}(\Sigma_2, S)$ such that

$$\text{sign } \eta(\alpha) = \text{sign}(\alpha \cdot \gamma)$$

for every $\alpha \in H_1(\Sigma_2)$ representing a curve of S . If n is large, then each $\alpha \in H_1(\Sigma_2)$ representing a curve of S is homologous to a very tall cylinder C_α on (X_n, η_{X_n}) , and this cylinder contributes positively to $\text{Im } \eta_{X_n}(\gamma)$. If n is sufficiently large, then the contributions from these cylinders will be much greater than that from the rest of (X_n, η_{X_n}) , and we will have $\text{Im } \eta_{X_n}(\gamma) \rightarrow \infty$. ■

Proof of Theorem 5.7. Let $\{\eta^i\}$ be a basis of $\Omega(\tilde{X})$ such that $\|\eta^i\| = 1$. There is some $C' > 0$ such that

$$\sup_i |\eta^i(\gamma)| > C'$$

for all $\gamma \in V_{S, \mathbb{Z}}^\perp / V_{S, \mathbb{Z}}$. Let

$$C = \frac{1}{2} \sup_i C_i.$$

For each i , there is a sequence $\eta_n^i \in \Omega(X_n)$ such that

$$\eta_n^i \rightarrow \eta^i. \quad (5.5)$$

By Theorem 5.5, we have

$$\|\eta_n^i\| \rightarrow \|\eta^i\|,$$

so we can normalize η_n^i so that $\|\eta_n^i\| = 1$ for all n , and (5.5) still holds. By the uniform convergence statement of Theorem 5.5, we have for each i and $\gamma \in V_{S,\mathbb{Z}}^\perp$,

$$|\eta_n^i(\gamma)| > \frac{1}{2}|\eta^i(\gamma)|$$

for n sufficiently large. The first claim of the Theorem follows.

Let $\alpha_i \in H_1(\Sigma_2; \mathbb{Z})$ be homology classes representing each nonseparating curve S_i of S . Choose a section $Y \mapsto \omega_Y$ of $\Omega\overline{T}(\Sigma_2; S)$ as in Lemma 5.8 such that

$$\omega(\alpha_i) = \mathbb{R} \setminus \{0\}$$

for each i . We claim that for every $C > 0$ and $\gamma \in H_1(\Sigma_2; \mathbb{Z}) \setminus V_{S,\mathbb{Z}}^\perp$ such that

$$\text{sign } \alpha \cdot \gamma = \text{sign } \omega_X(\alpha), \quad (5.6)$$

there is some N for which we have

$$\frac{|\omega_{X_n}(\gamma)|}{\|\omega_{X_n}\|} > C$$

for all $n > N$. The second claim of the Theorem follows because we can choose finitely many such sections so that for all $\gamma \in H_1(\Sigma_2; \mathbb{Z}) \setminus V_{S,\mathbb{Z}}^\perp$, (5.6) holds for some section.

Let $M \subset X$ be a compact subset which is a deformation retract of the complement of the nodes. By Theorem 5.5, there are collapses $g_n: X_n \rightarrow X$ such that

$$K_n^{-1} < \left| \frac{(g_n)^* \omega_X}{\omega_{X_n}} \right| < K_n$$

on $M_n = g_n^{-1}(M)$ with $K_n \geq 1$ and $K_n \rightarrow 1$. There are also cylinders $C_{i,n}$ in $X_n \setminus M_n$ which represent the homology classes α_i such that if $H_{i,n} = \text{height } C_{i,n}$, then $\lim H_{i,n} = \infty$.

The class γ is represented on each X_n by a simple closed curve γ_n . On X_n , each cylinder $C_{i,n}$ contributes

$$|\alpha_i \cdot \gamma| H_{i,n}$$

to $\text{Im} \int_\gamma \omega_{X_n}$. There are also segments of γ_n which pass through M_n to get from one cylinder to another. Let β be such a segment. Since we chose ω_{X_n} so that its periods over $V_{S,\mathbb{Z}}^\perp$ are all real, $\text{Im} \int_\beta \omega_{X_n}$ doesn't depend on the path β takes. If n is large, then this integral will be uniformly bounded over each such path β , say by

$$\left| \text{Im} \int_\beta \omega_{X_n} \right| < c = \text{diameter}(M) + 1.$$

We have

$$\|\omega_{X_n}\| = \left(\sum_i H_{i,n} + \text{Area}(M) \right)^{1/2} \leq \left(r \max_i H_{i,n} + \text{Area}(M) \right)^{1/2},$$

where r is the number of curves in S . Also, we have

$$\text{Im} \int_{\gamma_n} \omega \geq \max_i H_{i,n} - c.$$

Thus

$$\|\gamma\|_{X_n} \geq \frac{\left| \int_{\gamma_n} \omega_{X_n} \right|}{\|\omega_{X_n}\|} \geq \frac{\text{Im} \int_{\gamma_n} \omega_{X_n}}{\|\omega_{X_n}\|} \geq \frac{\max_i H_{i,n} - c}{\sqrt{r \max_i H_{i,n} + \text{Area}(M)}} \rightarrow \infty$$

as $n \rightarrow \infty$. ■

Real multiplication near $\partial\mathcal{M}_2$. We now use the two previous theorems to study eigenforms for real multiplication near $\partial\mathcal{M}_2$. As in the previous paragraph, we will write $V_{S,\mathbb{Z}}$ for the subgroup of $H_1(\Sigma_2; \mathbb{Z})$ generated by the curves in a curve system S on Σ_2 .

Theorem 5.9. *Let $X \in \mathcal{T}(\Sigma_2, S)$ be a stable Riemann surface whose Jacobian has real multiplication by $\mathcal{O}_D$. There is a neighborhood U of X in $\overline{\mathcal{T}}(\Sigma_2)$ such that for each stable Riemann surface $Y \in U$ which has real multiplication by $\mathcal{O}_D$, the real multiplication preserves the two subspaces $V_{S,\mathbb{Z}}$ and $V_{S,\mathbb{Z}}^\perp$ of $H_1(\Sigma_2; \mathbb{Z})$.*

Proof. If $\gamma \in V_{S,\mathbb{Z}}$, then we have $\|\gamma\|_X = 0$, so $\|\gamma\|_{X_n} \rightarrow 0$ as $n \rightarrow \infty$. Let $\rho_n: \mathcal{O}_D \rightarrow \text{End Jac}(X_n)$ be a choice of real multiplication, and let $\lambda \in \mathcal{O}_D$ be a generator of $\mathcal{O}_D$ over $\mathbb{Z}$. If the claim of the Theorem is false, then taking a subsequence, we can assume that $\rho_n(\lambda) \cdot \gamma \notin V_{S,\mathbb{Z}}$ for all n . We have

$$\|\rho_n(\lambda) \cdot \gamma\|_{X_n} \leq \sup_i |\lambda^{(i)}| \cdot \|\gamma\|_{X_n}.$$

This is a contradiction because the right hand side of this equation goes to zero while the left hand side is bounded below by Theorem 5.7.

The assertion that $V_{S,\mathbb{Z}}^\perp$ is preserved is proved in the same way. ■

Recall that we have the subspace $\Omega E_D \subset \Omega\mathcal{M}_2$ of eigenforms for real multiplication by $\mathcal{O}_D$. We can define real multiplication on semi-Abelian varieties just as we did for Abelian varieties in §2.2; however, we will not require real multiplication on a semi-Abelian variety to be proper. Let $\overline{\Omega E_D} \subset \overline{\Omega\mathcal{M}_2}$ be the closure of ΩE_D .

Theorem 5.10. *Every stable Abelian differential $(X, \omega) \in \overline{\Omega E_D}$ is a (not necessarily proper) eigenform for real multiplication by $\mathcal{O}_D$.*

Proof. Suppose (X_n, ω_n^1) is a sequence in $\Omega\mathcal{T}(\Sigma_2)$ is a sequence of ι_1 -eigenforms for real multiplication by $\mathcal{O}_D$ converging to some (X, ω^1) with $X \in \mathcal{T}(\Sigma_2, S)$. Let $\omega_n^2 \in \Omega(X_n)$ be a sequence of ι_2 -eigenforms. Taking a subsequence and suitably normalizing the ω_n^2 , we can assume that $\omega_n^2 \rightarrow \omega^2$ for some nonzero $\omega^2 \in \Omega(X)$.

We claim that the ω^i span $\Omega(X)$. Suppose not. Then $\omega^1 = c\omega^2$ for some $c \in \mathbb{C}$. Choose some $\alpha \in V_{S, \mathbb{Z}}^\perp$ such that $\omega^1(\alpha) \neq 0$, and let $\lambda \in \mathcal{O}_D \setminus \mathbb{Z}$ be a generator of $\mathcal{O}_D$ over $\mathbb{Z}$. Then for n large,

$$\lambda^{(1)} \omega^1(\alpha) \sim \omega_n^1(\rho_n(\lambda) \cdot \alpha) \sim c \omega_n^2(\rho_n(\lambda) \cdot \alpha) = c \lambda^{(2)} \omega_n^2(\alpha) \sim \lambda^{(2)} \omega^1(\alpha),$$

a contradiction because $\lambda^{(1)} \neq \lambda^{(2)}$.

Now via the dual bases to $\{\omega_n^i\}$ and $\{\omega^i\}$, we can identify $\Omega(X_n)^*$ and $\Omega(X)^*$ with $\mathbb{C}^2$. We then have natural embeddings

$$\begin{aligned} \phi_n &: H_1(\Sigma_2; \mathbb{Z}) \rightarrow \mathbb{C}^2, \quad \text{and} \\ \phi &: V_{S, \mathbb{Z}}^\perp \rightarrow \mathbb{C}^2 \end{aligned}$$

such that

$$\begin{aligned} \text{Jac}(X_n) &\cong \mathbb{C}^2 / \text{Im } \phi_n, \quad \text{and} \\ \text{Jac}(X) &\cong \mathbb{C}^2 / \text{Im } \phi, \end{aligned}$$

and $\phi_n \rightarrow \phi$ as $n \rightarrow \infty$. The real multiplication on $\text{Jac}(X_n)$ lifts to

$$\tilde{\rho}(z_1, z_2) = (\lambda^{(1)} z_1, \lambda^{(2)} z_2)$$

on $\mathbb{C}^2$. Since $\phi_n \rightarrow \phi$, the map $\tilde{\rho}$ preserves $\text{Im } \phi$, so defines real multiplication on $\text{Jac}(X)$ with ω^1 an eigenform as desired. $\blacksquare$

6 Local coordinates for strata in $\Omega\overline{\mathcal{M}}_2$

While the stratification of $\Omega\mathcal{M}_2$ is very simple, consisting of only two strata, the natural stratification of $\Omega\overline{\mathcal{M}}_2$ is much more complicated. With the appropriate definition of a stratum, there are seventeen different strata in $\Omega\overline{\mathcal{M}}_2$. It will be necessary for our study of the compactification of X_D to understand them. In particular, we will need to give local coordinates on $\Omega\overline{\mathcal{M}}_2$ around most of these strata. In §6.1, we will define the stratification of $\Omega\overline{\mathcal{M}}_2$, give notation for the various strata, and list the seventeen strata in $\Omega\overline{\mathcal{M}}_2$. In the rest of this section, we will then discuss each stratum in turn.

6.1 Strata in $\Omega\overline{\mathcal{M}}_2$

Given two stable Abelian differentials (X_1, ω_1) and $(X_2, \omega_2) \in \Omega\overline{\mathcal{M}}_2$, say that they are in the same *stratum* of $\Omega\overline{\mathcal{M}}_2$ if there is a homeomorphism $f: X_1 \rightarrow X_2$ with the following properties:

- f takes zeros of ω_1 to zeros of ω_2 of the same order.
- f takes polar nodes of (X_1, ω_1) to polar nodes of (X_2, ω_2) .
- f takes irreducible components of X_1 on which ω_1 vanishes to irreducible components of X_2 on which ω_2 vanishes.

Notation for strata. We now introduce notation for the various strata in $\Omega\overline{\mathcal{M}}_2$. Let $S \subset \Sigma_2$ be a curve system, and define $\Omega\mathcal{T}_2^0(S)$ to be the locus of all $(X, \omega) \in \Omega\overline{\mathcal{T}}_2$ such that:

- $X \in \mathcal{T}_2(S)$;
- The form ω has poles at the nodes corresponding to nonseparating curves in S .

If $T \in S$ is a nonseparating curve, then define $\Omega\mathcal{T}_2^0(S, T)$ to be the locus of all $(X, \omega) \in \Omega\overline{\mathcal{T}}_2$ such that:

- $X \in \mathcal{T}_2(S)$;
- The form ω has poles at the nodes corresponding to nonseparating curves in $S \setminus T$;
- The form ω is holomorphic at the node corresponding to the curve T .

If $\mathbf{n}$ is either (2) or (1, 1), then let

$$\Omega\mathcal{T}_2^0(S; \mathbf{n}) \subset \Omega\mathcal{T}_2^0(S)$$

be the locus of $(X, \omega) \in \Omega\mathcal{T}_2^0(S)$ where in addition, ω has zeros whose orders are given by $\mathbf{n}$.

Let $\Omega\mathcal{M}_2^0(S, T)$ and $\Omega\mathcal{M}_2^0(S)$ and $\Omega\mathcal{M}_2^0(S; \mathbf{n})$ be the images in $\Omega\overline{\mathcal{M}}_2(S)$ of the corresponding spaces defined above.

Let $\Omega\mathcal{D}'_2(S, T)$ be the locus of Abelian differentials $(X, \omega) \in \Omega\mathcal{D}_2(S)$ such that either $\int_T \omega = 0$ or T represents a holomorphic node on (X, ω) . This is a nonsingular hypersurface in $\Omega\mathcal{D}_2(S)$.

Let $\Omega\mathcal{D}_2^0(S)$ be the locus of Abelian differentials $(X, \omega) \in \Omega\mathcal{D}_2(S)$ such that each nonseparating curve of $S \setminus T$ is represented by either a polar node of (X, ω) or a cylinder on (X, ω) . Given a curve $T \in S$, let

$$\Omega\mathcal{D}_2^0(S, T) \subset \Omega\mathcal{D}'_2(S, T)$$

be the locus of (X, ω) with the following properties:

- Each nonseparating curve of $S \setminus T$ is represented by either a polar node of (X, ω) or a cylinder on (X, ω) .
- A curve γ which is either T or a separating curve in S is either represented by a holomorphic node of (X, ω) or is homotopic to a union $I \cup J(I)$ of a saddle connection I with its image under the hyperelliptic involution J .

By Theorem 5.5, $\Omega\mathcal{D}_2^0(S, T)$ is a neighborhood of $\Omega\mathcal{T}_2^0(S, T)$ in $\Omega\mathcal{D}'_2(S, T)$.

Also, let

$$\Omega\mathcal{D}_2^0(S, \mathbf{n}) \subset \Omega\mathcal{D}_2^0(S)$$

be those $(X, \omega) \in \Omega\mathcal{D}_2^0(S, T)$ such that ω has zeros whose orders are given by $\mathbf{n}$.

Up to the action of the mapping class group, there are six curve systems in Σ_2 , shown in Figure 7. We will denote by $T_{i,j}$ a curve system with i separating curves and j nonseparating curves. Such a curve system is unique up to the action of the mapping class group. We will write $T_{i,j}^k$ with $k = 1, \dots, n$ for the n individual curves of $T_{i,j}$, and we will always order the curves so that if $T_{i,j}$ contains a separating curve, then it is the last curve $T_{i,j}^n$.

The seventeen strata. Here is a list of the seventeen strata in $\Omega\overline{\mathcal{M}}_2$.

1. $\Omega\mathcal{M}_2(1, 1)$ is the stratum of nonsingular Abelian differentials with two simple zeros.
2. $\Omega\mathcal{M}_2(2)$ is the stratum of nonsingular Abelian differentials with a double zero.
3. $\Omega\mathcal{M}_2^0(T_{0,1})$ is the stratum of stable Abelian differentials consisting of two nonzero genus one Abelian differentials joined at a node.
4. $\Omega\mathcal{M}_2^0(T_{1,0}; 1, 1)$ is the stratum of stable Abelian differentials with one nonseparating polar node and two simple zeros.
5. $\Omega\mathcal{M}_2^0(T_{1,0}; 2)$ is the stratum of stable Abelian differentials with one nonseparating polar node and one double zero.




Figure 7: Six curve systems on Σ_2 .

6. $\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$ is the stratum of stable Abelian differentials with one nonseparating holomorphic node. These can be regarded as genus one Abelian differentials with two points joined to a node, and they have no zeros.
7. $\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$ is the stratum of stable Abelian differentials with one nonseparating holomorphic node and one nonseparating polar node. These can be regarded as the differential $a dz/z$ on $\mathbb{C}^*$ with the points 0 and ∞ identified to form a polar node, and with two points in $\mathbb{C}^*$ identified to form a holomorphic node.
8. $\Omega\mathcal{M}_2^0(T_{2,0}; 1, 1)$ is the stratum of stable Abelian differentials with two nonseparating polar nodes and two simple zeros.
9. $\Omega\mathcal{M}_2^0(T_{2,0}; 2)$ is the stratum of stable Abelian differentials with two nonseparating polar nodes and a double zero.
10. $\Omega\mathcal{M}_2^0(T_{2,1})$ is the stratum of stable Abelian differentials with two nonseparating polar nodes and one separating holomorphic node. These can be regarded as two infinite cylinders with the ends of the cylinders identified to form polar nodes and one point from each cylinder identified to form a holomorphic node.
11. $\Omega\mathcal{M}_2^0(T_{3,0})$ is the stratum of stable Abelian differentials (X, ω) with three nonseparating polar nodes. These have two simple zeros, one in each irreducible component of X .
12. $\Omega\mathcal{M}_2^0(T_{3,0}, T_{3,0}^3)$ is the stratum of stable Abelian differentials with two nonseparating polar nodes and one nonseparating holomorphic node. These differentials have no zeros.
13. $\Omega\mathcal{M}_2^0(T_{1,1})$ is the stratum of stable Abelian differentials with one nonseparating polar node and one separating holomorphic node.
- 14-17. There are four other strata consisting of stable Abelian differentials (X, ω) which vanish on some irreducible component of X . There is one such stratum corresponding to the curve system $T_{0,1}$, one corresponding to the curve system $T_{2,1}$, and two corresponding to the curve system $T_{1,1}$.

We will now go down this list and discuss each of these strata. We will not discuss the stratum $\Omega\mathcal{M}_2(1, 1)$ because we don't need to know anything more about it, and we will not discuss the strata 13-17 on this list because they do not arise in this paper.

6.2 The strata $\Omega\mathcal{M}_2(2)$ and $\Omega\mathcal{M}_2(T_{0,1})$

The strata $\Omega\mathcal{M}_2(2)$ and $\Omega\mathcal{M}_2(T_{0,1})$ are suborbifolds of $\Omega\overline{\mathcal{M}}_2$. We will see this explicitly by extending the period coordinates on these strata to period coordinates on a neighborhood of these strata.

Period coordinates for $\Omega\mathcal{M}_2(2)$. The map $\text{Per}: \Omega\mathcal{T}_2 \rightarrow H^1(\Sigma_2; \mathbb{C})$ is a submersion by the period coordinates from §4.2, so the fibers of this map are the leaves of a foliation $\mathcal{A}$ of $\Omega\mathcal{T}_2$ by Riemann surfaces along whose leaves the absolute periods of Abelian differentials are constant. This foliation meets the stratum $\Omega\mathcal{T}_2(2)$ transversely because the restriction of Per to this stratum is locally biholomorphic.

We can locally parameterize the leaf L of $\mathcal{A}$ through some $(X, \omega) \in \Omega\mathcal{T}_2(2)$ using the operation of splitting a double zero from §4.1 as follows. Choose $\epsilon > 0$ such that every geodesic segment starting at the zero p of (X, ω) of length less than ϵ^3 is embedded in X . Since p is a cone point of cone angle 6π , we can parameterize short geodesic segments starting at p by the ϵ -disk $\Delta_\epsilon \subset \mathbb{C}$ by associating continuously to a point $z \in \Delta_\epsilon$ a geodesic segment $I(z) \subset X$ such that

$$\int_{I(z)} \omega = z^3. \quad (6.1)$$

The map $\phi: \Delta_\epsilon \rightarrow L$ defined by

$$\phi(z) = (X, \omega) \#_{I(z)}$$

is holomorphic and satisfies $\phi(z) = \phi(-z)$ because the segments $I(z)$ and $I(-z)$ differ by an angle of 3π , and so these segments determine the same “X” in (X, ω) (see Figure 3), along which we perform surgery to form $\phi(z)$. Furthermore, $\phi(z) = \phi(w)$ if and only if $z = \pm w$, so ϕ lifts to a map $\tilde{\phi}$ defined by

$$\tilde{\phi}(z) = (X, \omega) \#_{I(z^{1/2})},$$

which maps Δ_ϵ conformally onto a neighborhood of (X, ω) in L .

Now let $U \subset \Omega\mathcal{T}_2(2)$ be a small neighborhood of (X, ω) . We can parameterize the above construction to define continuously for each $(Y, \eta) \in U$ and $z \in \Delta_\epsilon$ (possibly making ϵ smaller) a geodesic segment $I_{(Y, \eta)}(z)$ on Y starting at the zero of η such that

$$\int_{I_{(Y, \eta)}(z)} \eta = z^3.$$

Define $\Phi: U \times \Delta_\epsilon \rightarrow \Omega\mathcal{T}_2$ by

$$\Phi((Y, \eta), z) = (Y, \eta) \#_{I_{(Y, \eta)}(z^{1/2})}.$$

Φ gives a conformal isomorphism of $\{(Y, \eta)\} \times \Delta_\epsilon$ onto a neighborhood of (Y, η) in the leaf of $\mathcal{A}$ through (Y, η) . Since $\mathcal{A}$ is transverse to $\Omega\mathcal{T}_2(2)$, it follows that Φ is biholomorphic onto its image in $\Omega\mathcal{T}_2(2)$ if U and ϵ are sufficiently small. By considering the inverse of Φ , we obtain the following local coordinates around points in $\Omega\mathcal{T}_2(2)$.

Proposition 6.1. *Every $(X, \omega) \in \Omega\mathcal{T}(2)$ has a neighborhood V such that:*

- *Each $(Y, \eta) \in V$ has either a shortest saddle connecting distinct zeros or a double zero.*

- The multi-valued function,

$$(Y, \eta) \mapsto \left(\int_I \eta \right)^{2/3},$$

has a single-valued branch y .

- The product $\text{Per} \times y: V \rightarrow H^1(\Sigma_2; \mathbb{C}) \times \mathbb{C}$ is biholomorphic onto its image.

Period coordinates for $\Omega\mathcal{M}_2(T_{0,1})$. The stratum $\Omega\mathcal{M}_2(T_{0,1})$ is a suborbifold of $\overline{\Omega\mathcal{M}}_2$ isomorphic to the symmetric square of $\Omega\mathcal{M}_{1,1}$. We give local coordinates for this stratum just as for $\Omega\mathcal{M}_2(2)$ using the connected sum construction. As above, let $\text{Per}: \Omega\mathcal{D}_2(T_{0,1}) \rightarrow H^1(\Sigma_2; \mathbb{C})$ be the natural period map.

Each $(X, \omega) \in \Omega\mathcal{T}_2(T_{0,1})$ is a one point connected sum of two genus one differentials:

$$(X, \omega) = (X_1, \omega_1) \# (X_2, \omega_2).$$

Given a small open set $U \subset \Omega\mathcal{T}_2(T_{0,1})$ and a sufficiently small $\epsilon > 0$, we define a holomorphic map $\Phi: U \times \Delta_\epsilon \rightarrow \Omega\mathcal{D}_2(T_{0,1})$ by

$$\Phi((X, \omega), z) = (X_1, \omega_1) \#_{I(z^{1/2})} (X_2, \omega_2),$$

where $I(z) \subset \mathbb{C}$ is the segment joining 0 to ϵ .

If $(Y, \eta) \in \Omega\mathcal{D}_2(T_{0,1})$ has a unique shortest saddle connection I , then define

$$y(Y, \eta) = \left(\int_I \eta \right)^2.$$

By taking an inverse of Φ , we obtain local coordinates around $\Omega\mathcal{T}_2(T_{0,1})$ as above.

Proposition 6.2. *Every $(X, \omega) \in \Omega\mathcal{T}_2(T_{0,1})$ has a neighborhood $V \subset \Omega\mathcal{D}_2(T_{0,1})$ such that:*

- Each $(Y, \eta) \in V$ has either a shortest saddle connecting distinct zeros or a separating node.
- The function $\text{Per} \times y: V \rightarrow H^1(\Sigma_2; \mathbb{C}) \times \mathbb{C}$ is biholomorphic onto its image.

6.3 The strata $\Omega\mathcal{M}_2^0(T_{1,0}; 1, 1)$ and $\Omega\mathcal{M}_2^0(T_{1,0}; 2)$

The locus $\Omega\mathcal{M}_2^0(T_{1,0})$ – the union of the strata $\Omega\mathcal{M}_2^0(T_{1,0}; 1, 1)$ and $\Omega\mathcal{M}_2^0(T_{1,0}; 2)$ – is a four dimensional orbifold. It can be naturally embedded in the bundle $\Omega'\mathcal{M}_{1,2}$ consisting of meromorphic Abelian differentials (X, ω) on an elliptic curve X with simple poles at two marked points p and q such that

$$\text{Res}_p \omega = -\text{Res}_q \omega.$$

Its image is the complement of the sub-line-bundle where these residues are zero. The stratum $\Omega\mathcal{M}_2^0(T_{1,0}; 1, 1)$ is an open, dense suborbifold of $\Omega\mathcal{M}_2^0(T_{1,0})$, and $\Omega\mathcal{M}_2^0(T_{1,0}; 2)$ is a closed, three-dimensional suborbifold.

Local coordinates. We now give local coordinates on neighborhoods of these strata. Choose a symplectic basis $\{\alpha_i, \beta_i\}_{i=1}^2$ of $H_1(\Sigma_2; \mathbb{Z})$ such that α_1 represents the single curve in $T_{1,0}$.

Define functions on $\Omega\mathcal{D}_2^0(T_{1,0})$ as follows: given $(Y, \eta) \in \Omega\mathcal{D}_2^0(T_{1,0})$, define

$$\begin{aligned} v &= \eta(\alpha_1) & w &= \eta(\alpha_2) \\ x &= \eta(\beta_2) & z &= e^{2\pi i \eta(\beta_1)/\eta(\alpha_1)}, \end{aligned}$$

where we consider $z(Y, \eta)$ to be 0 if Y has a node. These are well-defined functions because α_1 , α_2 , and β_2 give well-defined homology classes on Y via the marking, and β_1 gives a homology class on Y which is well-defined up to adding a multiple of α_1 . These are all holomorphic functions on U . This follows from the fact that these functions are all holomorphic on $\Omega\mathcal{D}_2^0(T_{1,0}) \setminus \Omega\mathcal{T}_2^0(T_{1,0})$ because periods vary holomorphically, together with the fact that these functions are all continuous on U , which follows from Theorem 5.5. Functions which are continuous and holomorphic on an open dense subset of their domain are holomorphic, so these functions v , w , x , and z are all holomorphic.

Now, let $(X, \omega) \in \Omega\mathcal{T}_2^0(T_{1,0})$, and let $U \subset \Omega\mathcal{D}_2^0(T_{1,0})$ be an open neighborhood of (X, ω) with a continuous choice of a saddle connection $I_{(Y, \eta)}$ on each $(Y, \eta) \in U$ joining distinct zeros. If (X, ω) has two simple zeros, then we can also continuously orient these saddle connections, but this is not possible if (X, ω) has a double zero. For $(Y, \eta) \in U$, define

$$y = \begin{cases} \int_{I_{(Y, \eta)}} \eta & \text{if } (X, \omega) \text{ has two simple zeros;} \\ \left(\int_{I_{(Y, \eta)}} \eta \right)^{2/3} & \text{if } (X, \omega) \text{ has a double zero,} \end{cases}$$

If U is sufficiently small, then the functions (v, w, x, y) are a system of holomorphic local coordinates on $\Omega\mathcal{T}_2^0(T_{1,0}) \cap U$ by (a simple generalization of) the period coordinates from [Vee90] together with the argument of Proposition 6.1.

Recall that in §4.1, we defined the unplumbing operation which replaced an Abelian differential with a choice of a cylinder with a new meromorphic Abelian differential with two simple poles. Since these poles have opposite residues, we can join them to a node and obtain a stable Abelian differential.

Define the *unplumbing map* to be the function,

$$\psi: \Omega\mathcal{D}_2^0(T_{1,0}) \rightarrow \Omega\mathcal{T}_2^0(T_{1,0}),$$

defined by sending (Y, η) to the Abelian differential obtained by unplumbing the cylinder on Y which is homotopic to the single curve in the curve system $T_{1,0}$. The function ψ is continuous and is holomorphic on

$$\Omega\mathcal{D}_2^0(T_{1,0}) \setminus \Omega\mathcal{T}_2^0(T_{1,0}),$$

as can easily be seen using period coordinates, so ψ is holomorphic on all of $\Omega\mathcal{D}_2^0(T_{1,0})$. Let $\Psi = \psi \times z$.

Proposition 6.3. *The map Ψ is a biholomorphic map of $\Omega\mathcal{D}_2^0(T_{1,0})$ onto its image in $\Omega\mathcal{T}_2^0(T_{1,0}) \times \mathbb{C}$.*

It follows that if U is sufficiently small, the functions (v, w, x, y, z) are a holomorphic system of local coordinates on the neighborhood U of (X, ω) .

Proof. We can show that Ψ is biholomorphic onto its image by constructing a holomorphic inverse. Given $(Y, \eta) \in \Omega\mathcal{T}_2^0(T_{1,0})$ and $u \in \mathbb{C}^*$, if $|u|$ is sufficiently small, then we can uniquely plumb a cylinder into the node of (Y, η) to form a new nonsingular Abelian differential (Y', η') such that $z(Y', \eta') = u$. The point is that the plumbing construction is determined by two quantities: the height of the resulting cylinder, and the twist of the gluing map; the height is determined by $|u|$, and the twist is determined by $\arg u$.

Let $V \subset \Omega\mathcal{T}_2^0(T_{1,0}) \times \mathbb{C}$ be the open neighborhood of $\Omega\mathcal{T}_2^0(T_{1,0}) \times \{0\}$ where this operation is defined, and let $\Phi: V \rightarrow \Omega\mathcal{D}_2^0(T_{1,0})$ be the plumbing map. The map Φ is continuous and is holomorphic on $V \setminus (\Omega\mathcal{T}_2^0(T_{1,0}) \times \{0\})$ by period coordinates, so it is everywhere holomorphic. The maps Φ and Ψ are inverse to each other, so both are biholomorphic.

The last statement then follows from the fact that (v, w, x, y) are a system of local coordinates on $\Omega\mathcal{T}_2^0(T_{1,0})$ on a neighborhood of (X, ω) . $\blacksquare$

Cylinder covers. The stratum $\mathcal{M}_2(T_{1,0}) \subset \overline{\mathcal{M}}_2$ of stable Riemann surfaces with one nonseparating node is naturally isomorphic to the moduli space $\mathcal{M}_{1,2}$ of elliptic curves with two marked points, and we will implicitly identify them. In $\mathcal{M}_2(T_{1,0})$, let $\mathcal{M}_2(T_{1,0})(d)$ be the locus of all X such that the two marked points on X differ by exactly d -torsion in the group law on the underlying elliptic curve. This locus $\mathcal{M}_2(T_{1,0})(d)$ is isomorphic to modular curve $\mathbb{H}/\Gamma_1(d)$.

Given a marked elliptic curve X in $\mathcal{M}_{1,2}$, call a meromorphic function $f: X \rightarrow \mathbb{P}^1$ a *cylinder cover* if the poles and zeros of f are located at the two marked points of X . By Riemann-Hurwitz, such an f has two other branch points counting multiplicity. We say that f is primitive if f is not of the form g^r for some meromorphic function g and $r > 1$. A meromorphic Abelian differential (X, ω) in $\Omega'\mathcal{M}_{1,2}$ is a *cylinder covering differential* if $\omega = df/f$ for some primitive cylinder cover. The *degree* of ω is the degree of the primitive cover. In terms of stable Riemann surfaces, a cylinder cover is a morphism $f: X \rightarrow C$, where $X \in \mathcal{M}_2(T_{1,0})$, the curve C is $\mathbb{P}^1$ with 0 and ∞ identified to form a node, and the inverse image of the node of C is the node of X .

Let $\Omega\mathcal{M}_2^0(T_{1,0})(d)$ be the locus of degree d cylinder covering differentials in $\Omega\mathcal{M}_2^0(T_{1,0})$.

Proposition 6.4. *A stable Riemann surface $X \in \mathcal{M}_2(T_{1,0})$ has a degree d cylinder covering differential if and only if $X \in \mathcal{M}_2(T_{1,0})(d)$, in which case it is unique up to constant multiple.*

Proof. To prove uniqueness, assume that f and g are two degree d cylinder covers. This means that f and g have the same zeros and poles, so $f = cg$ for some $c \in \mathbb{C}$. Thus $df/f = dg/g$.

Recall that if E is an elliptic curve and $\text{Div}^0(E)$ is the group of divisors of degree 0 on E , then there is a natural map $\mu: \text{Div}^0(E) \rightarrow E$, defined via the group law on E . Abel's Theorem says that a divisor $D \in \text{Div}^0(E)$ is the divisor of a meromorphic function on E if and only if $\mu(D) = 0$ (for example, see [GH78, p. 235]).

Now let $X \in \mathcal{M}_2(T_{1,0})$, which we'll regard as an elliptic curve with two marked points p and q . Choose an identity point 0 on X , fixing the group law. If X has a degree d cylinder cover $f: X \rightarrow \mathbb{P}^1$, with its zeros at p and its poles at q , then $(f) = d(p - q)$ (where (f) denotes the divisor associated to f), and by Abel's Theorem, $d(p - q) = 0$ in the group law on E ; that is, p and q differ by d -torsion.

We need to show that there is no $e < d$ with $e|d$ and $e(p - q) = 0$. Suppose there is such an e . Then by Abel's Theorem again, there would be a meromorphic function h on X with $(h) = e(p - q)$, which implies that h is a degree e cylinder cover. By the uniqueness statement, $f = ch^{d/e}$ for some $c \in \mathbb{C}$, contradicting primitivity of f .

The proof of the converse of this proposition is also a straightforward application of Abel's Theorem and will be left to the reader. $\blacksquare$

Corollary 6.5. *The suborbifold $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})(d)$ of $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})$ is isomorphic to $\mathbb{H}/\Gamma_1(d)$.*

6.4 The stratum $\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$

We now give local coordinates around the stratum $\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$. We will actually only give local coordinates on a hypersurface containing this stratum, but this will be sufficient for our purposes.

Let $\{\alpha_i, \beta_i\}_{i=1}^2$ be a symplectic basis of $H_1(\Sigma_2; \mathbb{Z})$ as in the previous section. Given $(Y, \eta) \in \Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$, define

$$\begin{aligned} w &= \eta(\alpha_2) & x &= \eta(\beta_2) \\ y &= \eta(\beta_1) & z &= \left(\int_I \eta \right)^2, \end{aligned}$$

where I is one of the two saddle connections on (Y, η) such that $I \cup J(I)$ is homotopic to the curve $T_{1,0}^1$ in $T_{1,0}$. These are holomorphic functions on $\Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$. If $(Y, \eta) \in \Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$ is regarded as a genus one differential with two marked points, then $y(Y, \eta)$ is the integral of η along a path joining the marked points. The functions (w, x, y) are a system of holomorphic local coordinates on $\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$.

Proposition 6.6. *For any $(X, \omega) \in \Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$, the functions (w, x, y, z) define a system of holomorphic local coordinates on a sufficiently small neighborhood of (X, ω) in $\Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$.*

Sketch of proof. The proof of this proposition is completely analogous to the proof of Proposition 6.3. Let

$$\psi: \Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1) \rightarrow \Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$$

be the map which sends an Abelian differential to the one obtained by splitting along the union of saddle connections $I \cup J(I)$ homotopic to the curve $T_{1,0}^1$. Let $\Psi = \psi \times z$. This is a holomorphic map,

$$\Psi: \Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1) \rightarrow \Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1) \times \mathbb{C}$$

Given any $u \in \mathbb{C}$, let $I(u) \subset \mathbb{C}$ be the segment connecting 0 to u . For any $(Y, \eta) \in \Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$, regarded as a genus one differential with two marked points, if $|u|$ is sufficiently small, then there are two parallel embeddings

$$\epsilon_i: I(u^{1/2}) \rightarrow (Y, \eta),$$

each sending 0 to one of the marked points of (Y, η) . Let

$$V \subset \Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1) \times \mathbb{C}$$

be the neighborhood of $\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1) \times \{0\}$ where these two embeddings ϵ_i exist, and define

$$\Phi: V \rightarrow \Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$$

to be the map which replaces (Y, η) with the self connected sum

$$(Y, \eta) \#_{I(z^{1/2})}.$$

Φ is a holomorphic map, and since Φ and Ψ are inverse to each other, both are biholomorphic. It then follows from the fact that (w, x, y) give local coordinates on $\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$ around (X, ω) that (w, x, y, z) give local coordinates on $\Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$ around (X, ω) . $\blacksquare$

The d -torsion locus. Let

$$\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d) \subset \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$$

be the locus of genus one differentials such that the two marked points differ by torsion of degree exactly d . One can show using period coordinates that $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$ is a suborbifold of $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$. We have the isomorphism,

$$\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d) \cong \mathcal{M}_2(T_{1,0})(d) \cong \mathbb{H}/\Gamma_1(d).$$

6.5 The stratum $\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$

We now define local coordinates on a hypersurface containing the stratum $\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$. Choose a symplectic basis $\{\alpha_i, \beta_i\}_{i=1}^2$ of $H_1(\Sigma_2; \mathbb{Z})$ such that α_i represents the curve $T_{2,0}^i$ in $T_{2,0}$. Given $(Y, \eta) \in \Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$, define

$$\begin{aligned} w &= \eta(\alpha_1) & x &= \eta(\beta_2) \\ y &= e^{2\pi i \eta(\beta_1)/\eta(\alpha_1)} & z &= \left(\int_I \eta \right)^2, \end{aligned}$$

where I is a saddle connection joining distinct zeros on (Y, η) such that $I \cup J(I)$ is homotopic to the curve $T_{2,0}^2$. These are holomorphic functions on $\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$, and the functions (w, x) are holomorphic local coordinates on $\Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)$.

Proposition 6.7. *Given $(X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)$, the functions (w, x, y, z) give a system of holomorphic local coordinates on a sufficiently small neighborhood U of (X, ω) in $\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$.*

The proof is a straightforward combination of the proofs of Propositions 6.3 and 6.6, so we will omit it as well of the proofs of further similar descriptions of local coordinates.

Points in $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$. Every Abelian differential in $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$ can be obtained by the following construction: given $r \in \mathbb{C}/\mathbb{Z}$, let f_r be the Abelian differential obtained from the cylinder $(\mathbb{C}/\mathbb{Z}, dz)$ by identifying the ends of the cylinder to form a polar node and identifying the points 0 and r to form a holomorphic node. Two Abelian differentials f_r and $f_{r'}$ are the same if and only if r and r' are related by the group G generated by the transformations $z \mapsto z + 1$ and $z \mapsto -z$. This gives an isomorphism of $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$ with

$$(\mathbb{C} \setminus \mathbb{Z})/G \cong (\mathbb{C} \setminus \{0, 1\})/(z \mapsto z^{-1}).$$

In $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$, let

$$\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d) = \bigcup_{q \in \Lambda} f_{q/d},$$

where

$$\Lambda = \{q \in \mathbb{Z} : \gcd(q, d) = 1\}.$$

We can regard $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d)$ as the locus of stable Abelian differentials $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$ such that – if we regard (X, ω) as $(\mathbb{C}^*, dz/z)$ with two marked points – the two marked points differ by exactly d -torsion in $\mathbb{C}^*$.

$\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$ contains N points, where

$$N = \begin{cases} 1 & \text{if } d = 2; \\ \frac{1}{2}\phi(d) & \text{if } d > 2. \end{cases}$$

6.6 The strata $\Omega\mathcal{M}_2^0(T_{2,0}; 1, 1)$, $\Omega\mathcal{M}_2^0(T_{2,0}; 2)$, and $\Omega\mathcal{M}_2^0(T_{2,1})$

We now give local coordinates around the strata $\Omega\mathcal{M}_2^0(T_{2,0}; 1, 1)$, $\Omega\mathcal{M}_2^0(T_{2,0}; 2)$, and $\Omega\mathcal{M}_2^0(T_{2,1})$. In each of these cases, let $\{\alpha_i, \beta_i\}_{i=1}^2$ be a symplectic basis of $H_1(\Sigma_2; \mathbb{Z})$ such that the α_i represent the nonseparating curves of the curve system. Fix an Abelian differential (X, ω) in $\Omega\mathcal{T}_2^0(T_{2,0}; 1, 1)$, $\Omega\mathcal{T}_2^0(T_{2,0}; 2)$, or $\Omega\mathcal{T}_2^0(T_{2,1})$, and choose a small neighborhood U of (X, ω) in $\Omega\mathcal{D}_2^0(T_{2,0}; 1, 1)$, $\Omega\mathcal{D}_2^0(T_{2,0})$, or $\Omega\mathcal{D}_2^0(T_{2,1})$ respectively. If U is sufficiently small, then we can continuously choose for each $(Y, \eta) \in U$ having distinct zeros a saddle connection $I_{(Y, \eta)}$ connecting those zeros. If (X, ω) has two distinct zeros, then we can also continuously choose an orientation for these saddle connections, but this is not possible if it has a double zero or a separating node. Given $(Y, \eta) \in U$, define

$$\begin{aligned} v &= \eta(\alpha_1) & w &= \eta(\alpha_2) \\ y &= e^{2\pi i \eta(\beta_1)/\eta(\alpha_1)} & z &= e^{2\pi i \eta(\beta_2)/\eta(\alpha_2)} \\ x &= \begin{cases} \int_I \eta & \text{if } (X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,0}; 1, 1); \\ (\int_I \eta)^{2/3} & \text{if } (X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,0}; 2); \\ (\int_I \eta)^2 & \text{if } (X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,1}). \end{cases} \end{aligned}$$

These are holomorphic functions on U , and (v, w, x) give a holomorphic system of local coordinates on $\Omega\mathcal{T}_2^0(T_{2,0})$ in the first two cases. In the last case where $(X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,1})$, the (v, w, x) give a system of local coordinates near (X, ω) on the subspace of $\Omega\mathcal{D}_2^0(T_{2,1})$ where the nonseparating curves of $T_{2,1}$ remain nodes.

Proposition 6.8. *The functions (v, w, x, y, z) give a system of holomorphic local coordinates on a neighborhood U of (X, ω) in $\Omega\mathcal{D}_2^0(T_{2,0}; 1, 1)$, $\Omega\mathcal{D}_2^0(T_{2,0}; 2)$, or $\Omega\mathcal{D}_2^0(T_{2,1})$.*

Curves in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$. We now discuss some curves in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$ which will play a prominent role in the rest of this paper. Given $\lambda \in \mathbb{C}^*$, let C_λ be the locus of points $(X, [\omega]) \in \mathbb{P}\Omega\overline{\mathcal{M}}_2$ having two nonseparating nodes and possibly a separating node such that the ratio of the residues of ω at the two nonseparating nodes of X is $\pm\lambda^{\pm 1}$. Two such curves C_λ and $C_{\lambda'}$ coincide if and only if $\lambda' = \pm\lambda^{\pm 1}$. Also, let $c_\lambda \in \mathbb{P}\Omega\overline{\mathcal{M}}_2$ be the Abelian differential with three nonseparating nodes and residues 1, λ , and $\lambda - 1$ as in Figure 8.

Proposition 6.9. *The curve $C_\lambda \subset \mathbb{P}\Omega\overline{\mathcal{M}}_2$ is a twice-punctured sphere. If $\lambda = \pm 1$, then it is an orbifold locus of order two; otherwise it contains no orbifold points of $\mathbb{P}\Omega\overline{\mathcal{M}}_2$. The punctures of C_λ are the two points c_λ and $c_{\lambda+1}$. The curve C_λ contains exactly one point of $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,1})$.*

Proof. Let $C_\lambda^0 \subset C_\lambda$ be the subset of differentials with no separating node. We claim that C_λ^0 is a thrice-punctured sphere. The natural map $C_\lambda^0 \rightarrow \mathcal{M}_2(T_{2,0})$ is an isomorphism. Let $\mathcal{M}_{0,4}^{\text{num}}$ be the moduli space of four *numbered* points on $\mathbb{P}^1$. There is a natural map $\pi: \mathcal{M}_{0,4}^{\text{num}} \rightarrow \mathcal{M}_2(T_{2,0})$ which sends the point




Figure 8: The point $c_\lambda \in \mathbb{P}\Omega\overline{\mathcal{M}}_2$.

$(p_1, p_2, p_3, p_4) \in \mathcal{M}_{0,4}^{\text{num}}$ to the stable Riemann surface obtained by identifying (p_1, p_2) and (p_3, p_4) to nodes. There are four ways to renumber the points p_i to get the same stable Riemann surface, so a fiber of π potentially contains four points; however, the automorphism group of the p_i contains a Klein four-group whose action on the p_i realizes any renumbering of the p_i which preserves the decomposition into the two sets $\{p_1, p_2\}$ and $\{p_3, p_4\}$. Therefore π is an isomorphism. There is thus an isomorphism $\tilde{\pi}: \mathcal{M}_{0,4}^{\text{num}} \rightarrow C_\lambda^0$, defined by sending the point $Q = (p_1, p_2, p_3, p_4) \in \mathcal{M}_{0,4}^{\text{num}}$ to (X, ω) , where $X = \pi(Q)$, and ω is induced by the unique meromorphic Abelian differential η on $\mathbb{P}^1$ having a simple pole at each p_i with residue given by

$$\begin{aligned} \text{Res}_{p_1} \eta &= 1 & \text{Res}_{p_2} \eta &= -1 \\ \text{Res}_{p_3} \eta &= \lambda & \text{Res}_{p_4} \eta &= -\lambda. \end{aligned}$$

It is well known that $\mathcal{M}_{0,4}^{\text{num}}$ is a thrice-punctured sphere, so C_λ^0 is as well.

Cusps of $\mathcal{M}_{0,4}^{\text{num}}$ correspond to isotopy classes of simple closed curves in $\mathbb{P}^1 \setminus \{p_i\}_{i=1}^4$ up to the action of the modular group $\text{Mod}_{0,4}^{\text{num}}$ of self-homeomorphisms of $\mathbb{P}^1$ fixing each of the p_i up to isotopy fixing the p_i . There are three such isotopy classes of curve. These are shown in Figure 9 with the corresponding limiting Abelian differential in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$. Thus we see that C_λ contains one point in $\mathbb{P}\Omega\overline{\mathcal{M}}_2^0(T_{2,1})$, and the two cusps of C_λ are as claimed.

To obtain the statement about orbifold points, note that for each $(X, [\omega]) \in C_\lambda$, the automorphism group of X is the Klein four group. If $\lambda = \pm 1$, then all of these automorphisms stabilize the projective class of ω , while if $\lambda \neq \pm 1$, only the hyperelliptic involution stabilizes $[\omega]$. ■

Proposition 6.10. *If $\lambda \neq \pm 1$, then $\mathbb{P}\Omega\overline{\mathcal{M}}_2^0(T_{2,0}; 2)$ meets C_λ in exactly one point. Otherwise they are disjoint.*

Proof. Define an isomorphism $\pi: \mathbb{C} \setminus \{0, 1\} \rightarrow C_\lambda^0$ by $p(w) = (X_w, \omega_w)$, where X_w is the stable Riemann surface obtained by identifying the points $\{0, 1\}$ and




Figure 9: Three curves determining cusps of C_λ^0 .

$\{w, \infty\}$ to nodes, and ω_w is induced by the meromorphic Abelian differential,

$$\begin{aligned}\omega'_w &= \left(\frac{1}{z} - \frac{1}{z-1} + \frac{\lambda}{z-w} \right) dz \\ &= \frac{\lambda z^2 + (-1-\lambda)z + w}{z(z-1)(z-w)} dz,\end{aligned}$$

on $\mathbb{P}^1$.

The differential ω'_w has a double zero if and only if

$$\lambda z^2 + (-1-\lambda)z + w$$

has a double zero in $\mathbb{C} \setminus \{0, 1, w\}$, which happens if and only if the discriminant,

$$\Delta_w = (1+\lambda)^2 - 4\lambda w,$$

vanishes for some $w \in \mathbb{C} \setminus \{0, 1\}$. This happens if and only if $\lambda \neq \pm 1$, in which case, there is a unique w such that $\Delta_w = 0$. Thus there is a unique Abelian differential in C_λ with a double zero if $\lambda \neq \pm 1$, and there is no such Abelian differential if $\lambda = \pm 1$. $\blacksquare$

For $\lambda \neq 0$ or ± 1 , let $w_\lambda \in C_\lambda$ be the unique point representing a stable Abelian differential with a double zero. Also, for $\lambda \neq 0$, let $p_\lambda \in C_\lambda$ be the unique point representing a stable Abelian differential with a nonseparating node.

6.7 The stratum $\Omega\mathcal{M}_2^0(T_{3,0})$

We now give local coordinates around points in $\Omega\mathcal{M}_2^0(T_{3,0})$, the stratum of stable Abelian differentials with three nonseparating polar nodes. There is a unique stable Riemann surface $X \in \mathcal{M}_2(T_{3,0})$ consisting of two thrice-punctured spheres with the cusps joined together to form three separating nodes. The automorphism group of X is $S_3 \times \mathbb{Z}/2$, with S_3 acting on each irreducible component of X by permuting the nodes and $\mathbb{Z}/2$ acting by permuting the irreducible components of X . This means that the stratum $\Omega\mathcal{M}_2^0(T_{3,0})$ is isomorphic to $\mathbb{C}^2/(S_3 \times \mathbb{Z}/2)$.

Choose two points $p, q \in \Sigma_2 \setminus T_{3,0}$. Let $\alpha_i \in H_1(\Sigma_2 \setminus \{p, q\}; \mathbb{Z})$ be a homology class representing the curve $T_{3,0}^i \in T_{3,0}$. Let $\gamma_i \in H_1(\Sigma_2, \{p, q\}; \mathbb{Z})$ be homology classes such that $\alpha_i \cdot \gamma_j = \delta_{ij}$ (see Figure 10).

Any marked stable Abelian differential $(f, (Y, \eta)) \in \Omega\mathcal{D}_2^0(T_{3,0})$ has three cylinders C_i homotopic to the curves $T_{3,0}^i$ of $T_{3,0}$ and two simple zeros, one in each component of the complement of the cylinders. We can change f to a marking f_1 by an isotopy so that f_1 takes each curve $T_{3,0}^i$ into the cylinder C_i and takes the points p and q to zeros of η . Two such markings f_1 and f_2 which are isotopic to f are isotopic to each other by an isotopy sending for all time the points p and q to zeros of η and sending the curve $T_{3,0}^i$ into the cylinder C_i . This means that we can regard the classes α_i as homology classes in $H_1(Y \setminus Z(\eta); \mathbb{Z})$




Figure 10: Homology classes in $H_1(\Sigma_2, \{p, q\}; \mathbb{Z})$.

and the γ_i as homology classes in $H_1(Y, Z(\eta); \mathbb{Z})$, well-defined up to adding a multiple of α_i (because the marking f was only defined up to Dehn twist around the curves of $T_{3,0}$).

Given $(Y, \eta) \in \Omega\mathcal{D}_2^0(T_{3,0})$, define

$$\begin{aligned} v &= \eta(\alpha_1) & w &= \eta(\alpha_2) \\ x &= e^{2\pi i \eta(\gamma_1) / \eta(\alpha_1)} & y &= e^{2\pi i \eta(\gamma_2) / \eta(\alpha_2)} \\ z &= e^{2\pi i \eta(\gamma_3) / \eta(\alpha_3)} \end{aligned}$$

These are well-defined holomorphic functions on $\Omega\mathcal{D}_2^0(T_{3,0})$, and the coordinates (v, w) define a map $\Omega\mathcal{T}_2^0(T_{3,0}) \rightarrow \mathbb{C}^2$ which is biholomorphic onto its image.

Proposition 6.11. *The functions (v, w, x, y, z) define a map $\Omega\mathcal{D}_2^0(T_{3,0}) \rightarrow \mathbb{C}^5$ which is biholomorphic onto its image.*

6.8 The stratum $\Omega\mathcal{M}_2^0(T_{3,0}, T_{3,0}^3)$

Finally, consider the stratum $\Omega\mathcal{M}_2^0(T_{3,0}, T_{3,0}^3)$. An Abelian differential in this stratum can be regarded as a pair of infinite cylinders with an end of each cylinder identified with an end of the other cylinder to form two polar nodes, and with a point from each cylinder identified to form a holomorphic node. The stratum $\Omega\mathcal{M}_2^0(T_{3,0}, T_{3,0}^3)$ is isomorphic to $\mathbb{C}^* / \pm 1$, with the isomorphism sending (X, ω) to the residue of ω at one of the polar nodes. The whole stratum is an orbifold locus of order two in $\Omega\overline{\mathcal{M}}_2$, because each (X, ω) in this stratum has an involution which switches the irreducible components of X and preserves ω .

Choose a symplectic basis $\{\alpha_i, \beta_i\}_{i=1}^2$ of $H_1(\Sigma_2; \mathbb{Z})$ such that α_i represents the curve $T_{3,0}^i$, and let $\alpha_3 \in H_1(\Sigma_2; \mathbb{Z})$ be a class representing $T_{3,0}^3$. Given a

marked stable Abelian differential $(Y, \eta) \in \Omega\mathcal{D}_2^0(T_{3,0}, T_{3,0}^3)$, the class β_i determines a class in $H_1(Y; \mathbb{Z})$ which is well defined up to adding multiples of α_i and α_3 . This means that

$$e^{2\pi i \eta(\beta_i)/\eta(\alpha_i)}$$

is a well defined complex number because $\eta(\alpha_3) = 0$.

Given $(Y, \eta) \in \Omega\mathcal{D}_2^0(T_{3,0}, T_{3,0}^3)$, define

$$\begin{aligned} w &= \eta(\alpha_1) & x &= e^{2\pi i \eta(\beta_1)/\eta(\alpha_1)} \\ y &= e^{2\pi i \eta(\beta_2)/\eta(\alpha_2)} & z &= \left(\int_I \eta \right)^2, \end{aligned}$$

where I is a saddle connection on (Y, η) such that $I \cup J(I)$ is homotopic to the curve $T_{3,0}^3$. The coordinate w defines an isomorphism $\Omega\mathcal{T}_2^0(T_{3,0}, T_{3,0}^3) \rightarrow \mathbb{C}$.

Proposition 6.12. *The coordinates (w, x, y, z) give a biholomorphic isomorphism of $\Omega\mathcal{D}_2^0(T_{3,0}, T_{3,0}^3)$ onto its image in $\mathbb{C}^4$.*

7 Limits of eigenforms

7.1 Introduction

We now begin the study of the compactification of X_D . Recall that we have the locus $E_D \subset \mathbb{P}\Omega\overline{\mathcal{M}}_2$ of eigenforms for real multiplication by $\mathcal{O}_D$, and there is the isomorphism $j_1: E_D \rightarrow X_D \setminus P_D$ from Proposition 4.4. The inverse of j_1 extends to an embedding $k_1: X_D \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$. In this section we will study the closure of X_D in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$, which we denote by $\overline{X}_D$. Our goal is to classify exactly which stable Abelian differentials lie in $\overline{X}_D \setminus X_D$ and to understand the local structure of $\overline{X}_D$ and its strata around these points. More precisely, for each $(X, [\omega]) \in \overline{X}_D$ we will:

- Choose a neighborhood of $(X, [\omega])$ in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$ of the form $U/\text{Aut}(X, [\omega])$, where U is a neighborhood of $(X, [\omega])$ in an appropriate Dehn space;
- Give local coordinates on U as in §6;
- Give explicit equations for the inverse image $\pi^{-1}(\overline{X}_D)$ in U .

This procedure will be slightly modified when $(X, [\omega])$ has a nonseparating holomorphic node. In that case, we will only give local coordinates on a hypersurface V in U which we will show contains the inverse image $\pi^{-1}(\overline{X}_D)$ in U .

We will see that $\overline{X}_D$ in general has non-normal singularities along curves in $\overline{X}_D \setminus X_D$. To get a less singular compactification, we will pass to the normalization Y_D of $\overline{X}_D$, which we will study in §8 using the results of this section.

Stratification of $\overline{X}_D$. The stratification of $\Omega\overline{\mathcal{M}}_2$ which we discussed in §6 induces a stratification of $\overline{X}_D$. Given a stratum $\Omega\mathcal{M}_2^0(S, T; \mathbf{n})$, let $X_D(S, T, \mathbf{n})$ denote the stratum,

$$X_D(S, T, \mathbf{n}) = \overline{X}_D \cap \mathbb{P}\Omega\mathcal{M}_2^0(S, T, \mathbf{n}),$$

with the analogous notation if T or $\mathbf{n}$ is omitted. Here is a list of all of these strata with a summary of what we will prove about each one:

1. $X_D(1, 1)$ is the stratum of nonsingular eigenforms with two simple zeros. We called this locus $E_D(1, 1)$ in §4.3. It is an open, dense subset of X_D .
2. $X_D(2)$ is the stratum of nonsingular eigenforms with a double zero, otherwise known as W_D .
3. $X_D(T_{0,1})$ is the stratum of eigenforms consisting of two genus one differentials joined at a node. This is the curve P_D , which we studied in §2.3.
- 4-5. $X_D(T_{1,0})$ is the locus of limiting eigenforms with one nonseparating polar node, containing the stratum $X_D(T_{1,0}; 1, 1)$ as an open dense set and the stratum $X_D(T_{1,0}; 2)$ as a finite subset. We will show in §7.2 that $X_D(T_{1,0})$

is empty unless D is square. We will show in §7.3 that $\overline{X}_{d^2}$ is an orbifold around $X_{d^2}(T_{1,0})$, and $X_{d^2}(T_{1,0})$ is a suborbifold isomorphic to $\mathbb{H}/\Gamma_1(d)$. We will also show that

$$X_{d^2}(T_{1,0}) = \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})(d)$$

as subsets of $\mathbb{P}\Omega\overline{\mathcal{M}}_2$, where $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$ is the locus of degree d cylinder covering differentials discussed in §6.3. We will show that $X_{d^2}(T_{1,0}; 2)$ consists of the intersection points of $\overline{W}_{d^2}$ with $X_{d^2}(T_{1,0})$ and that these intersections are transverse if $d > 3$.

6. $X_D(T_{1,0}, T_{1,0}^1)$ is the stratum of limiting eigenforms with one nonseparating holomorphic node. We will show in §7.2 that this stratum is empty unless D is square. We will show in §7.4 that $\overline{X}_{d^2}$ is an orbifold around $X_{d^2}(T_{1,0}, T_{1,0}^1)$, and $X_{d^2}(T_{1,0}, T_{1,0}^1)$ is a suborbifold isomorphic to $\mathbb{H}/\Gamma_1(d)$. We will also show that

$$X_{d^2}(T_{1,0}, T_{1,0}^1) = \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$$

as subsets of $\mathbb{P}\Omega\overline{\mathcal{M}}_2$, where $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$ is the d -torsion locus introduced in §6.4.

7. $X_D(T_{2,0}, T_{2,0}^2)$ is the stratum of limiting eigenforms with one nonseparating holomorphic node and one nonseparating polar node. We will show in §7.2 that this stratum is empty unless D is square. In §7.5 we will show that when $D = d^2$, this stratum is the finite set,

$$X_{d^2}(T_{2,0}, T_{2,0}^2) = \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d),$$

introduced in §6.5. We will also show that $\overline{X}_{d^2}$ is nonsingular at these points and that each of these points is a transverse intersection point of the closures of the strata $X_{d^2}(T_{1,0})$ and $X_{d^2}(T_{1,0}, T_{1,0}^1)$.

- 8-10. The strata $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, and $X_D(T_{2,1})$ together consist of the limiting eigenforms with two nonseparating polar nodes, possibly with a separating node. We will show in §7.6 that the union of these strata is the union of the curves $C_{\lambda(P)}$, where P is a nondegenerate Y_D -prototype. We will see that $\overline{X}_D$ is in general singular along these curves and can even have several branches passing through them. We will assign a Y_D -prototype to each of these branches, and we will give explicit equations for these branches in local coordinates.
- 11-12. The strata $X_D(T_{3,0})$ and $X_D(T_{3,0}, T_{3,0}^3)$ together consist of the limiting eigenforms with three nonseparating nodes. We will show in §7.7 that $X_D(T_{3,0})$ is the union of the points $c_{\lambda(P)}$ over all nonterminal Y_D -prototypes P , and $X_D(T_{3,0}, T_{3,0}^3)$ is the union of the points $c_{\lambda(P)}$ over all terminal Y_D -prototypes P . We will see that $\overline{X}_D$ is in general singular at these points. We will assign Y_D -prototypes to the branches of $\overline{X}_D$ through each $c_{\lambda(P)}$, and we will give equations for these branches in local coordinates.

13-17. We will show in §7.2 that the stratum $X_D(T_{1,1})$ as well as the four strata consisting of stable Abelian differentials (X, ω) where ω vanishes on some irreducible component of X are all empty.

Bundles. Over any of the bundles of projective spaces $\mathbb{P}\Omega\overline{\mathcal{M}}_2$, $\mathbb{P}\Omega\mathcal{T}_2$, or $\mathbb{P}\Omega\mathcal{D}_2(S)$, there is a canonical line bundle $\mathcal{O}(-1)$, whose fiber over a projective class of Abelian differentials, $(X, [\omega])$ is the space of constant multiples of $[\omega]$. Define a Hermitian metric on each of these bundles by defining on the fiber over (X, ω) ,

$$h(\eta, \eta) = \int_X |\eta|^2.$$

This metric is singular over the $(X, [\omega])$ which have infinite area. For use in §9, we will give sections for $\mathcal{O}(-1)$ around points in $\overline{X}_D$ and calculate the norms of these sections.

7.2 Empty strata

Proposition 7.1. *If D is not square, and $(X, [\omega]) \in \overline{X}_D \setminus X_D$, then $\text{Jac}(X) \cong (\mathbb{C}^*)^2$, and the period map $\text{Per}_\omega: H_1(\text{Jac}(X); \mathbb{Z}) \rightarrow \mathbb{C}$ is injective.*

Proof. If $(X, [\omega]) \in \overline{X}_D$, then $(X, \omega) \in \overline{\Omega X}_D$ is a nonzero eigenform for real multiplication by $\mathcal{O}_D$ by Theorem 5.10, so $M = H_1(\text{Jac}(X); \mathbb{Z})$ is a torsion-free $\mathcal{O}_D$ -module. If D is not square, this implies that the $\mathbb{Z}$ -rank of M is even because $M \otimes \mathbb{Q}$ is a K_D -vector space. This means that either $\text{Jac}(X)$ is compact, or $\text{Jac}(X) \cong (\mathbb{C}^*)^2$. Since for any $X \in \overline{X}_D \setminus X_D$, the Jacobian $\text{Jac}(X)$ is noncompact, we must have $\text{Jac}(X) \cong (\mathbb{C}^*)^2$.

Since ω is an eigenform, $\text{Per}_\omega: M \rightarrow \mathbb{C}$ is $\mathcal{O}_D$ -linear, so $K = \text{Ker}(\text{Per}_\omega)$ is an $\mathcal{O}_D$ -submodule of M . If $K \neq 0$, then K must be a submodule of M of $\mathbb{Z}$ -rank two, so M/K would be a finite Abelian group. If $K \neq M$, then this contradicts the fact that Per_ω embeds M/K into $\mathbb{C}$; if $K = M$, then this contradicts the fact that ω is nonzero. ■

Corollary 7.2. *If D is not square, then all of the strata in $\overline{X}_D \setminus X_D$ besides $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, $X_D(T_{2,1})$, and $X_D(T_{3,0})$ are empty.*

Proposition 7.3. *The stratum $X_{d^2}(T_{1,1})$ as well as the four strata of $\overline{X}_{d^2}$ consisting of stable Abelian differentials (X, ω) where ω vanishes on some irreducible component of X are all empty.*

Proof. We know that each $(X, \omega) \in X_{d^2}$ is a degree d branched cover of a genus one Abelian differential by Proposition 4.5. For any stable Abelian differential $(X, \omega) \in \overline{X}_{d^2}$, we can take a limit of these branched covers to get a branched covering of degree at most d from (X, ω) to a genus one stable Abelian differential (Y, η) . If $(X, \omega) \in X_{d^2}(T_{1,1})$, then it consists of a one point connected sum of a genus one nonsingular Abelian differential (having finite area) with an infinite cylinder (having infinite area). This is impossible because the same

(Y, η) cannot be finitely covered by both an Abelian differential with infinite area and one with finite area.

The proofs that the other four strata are empty are all similar, so for concreteness we will show that there is no stable Abelian differential $(X, \omega) \in \overline{X}_{d^2}$ which is the one point connected sum,

$$(X_1, \omega_1) \# (X_2, 0),$$

where X_i are elliptic curves with ω_1 nonzero. Suppose $(X, \omega) = \lim(X_n, \omega_n)$ with $(X_n, \omega_n) \in X_{d^2}$. There is a sequence of separating curves $\gamma_n \subset X_n$ which are pinched to the node of X in the limit. If n is large, then γ_n is homotopic to a union of short saddle connections $I_1 \cup I_2$ by Theorem 5.5, and we can split along these saddle connections to obtain a pair of genus one forms (E_n^i, ω_n^i) . We have a degree d branched covering $f_n: (X_n, \omega_n) \rightarrow (F_n, \nu_n)$ over some genus one form (F_n, ν_n) . Taking a subsequence, (F_n, ν_n) converges to some genus one form (F, ν) which is nonsingular because otherwise E_n^1 would converge to a Riemann surface with a node which would contradict the assumption that X has only one node. Each of the genus one forms (E_n^i, ω_n^i) is then an unbranched cover of (F_n, ν_n) of degree less than d . Thus we have

$$\frac{1}{d} \leq \frac{\text{Area}(\omega_n^1)}{\text{Area}(\omega_n^2)} \leq d,$$

which is a contradiction because exactly one of the sequences ω_n^i must converge to zero. $\blacksquare$

7.3 The strata $X_D(T_{1,0}; 1, 1)$ and $X_D(T_{1,0}, 2)$

In this section, we will study the strata $X_D(T_{1,0}; 1, 1)$ and $X_D(T_{1,0}, 2)$, whose union is the locus $X_D(T_{1,0})$, consisting of Abelian differentials with one polar node. Since these are empty when D is not square, we will restrict to the case $D = d^2$. The goal of the rest of this section is to prove the following description of these strata.

Theorem 7.4. *The locus $X_{d^2}(T_{1,0})$ is exactly the locus $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})(d)$ of degree d cylinder covering differentials discussed in §6.3. The variety $\overline{X}_{d^2}$ is an orbifold around $X_{d^2}(T_{1,0})$ with $X_{d^2}(T_{1,0})$ a suborbifold isomorphic to $\mathbb{H}/\Gamma_1(d)$. The stratum $X_{d^2}(T_{1,0}; 2) \subset X_{d^2}(T_{1,0})$ is a finite subset equal to*

$$\overline{W}_{d^2} \cap X_{d^2}(T_{1,0}).$$

These intersections are transverse if $d > 3$.

Local coordinates. Let $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})$, and choose some marking $f: \Sigma_2 \rightarrow X$, so that we can regard $(X, [\omega])$ as a point in $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})$. Take a symplectic basis $\{\alpha_i, \beta_i\}$ of $H_1(\Sigma_2; \mathbb{Z})$ as in §6.3. On a small neighborhood U of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0})$, we can normalize each projective class $(Y, [\eta]) \in U$

so that $\eta(\alpha_1) = 1$. This allows us to identify U with the hypersurface $U_1 \subset \Omega\mathcal{D}_2^0(T_{1,0})$ consisting of all (Y, η) with $\eta(\alpha_1) = 1$. The restriction of the local coordinates (v, w, x, y, z) from §6.3 to U_1 then gives a system of local coordinates (w, x, y, z) on U .

Assume that U is small enough that the natural map $U/\text{Aut}(X, [\omega]) \rightarrow \mathbb{P}\Omega\mathcal{M}_2$ is an isomorphism onto its image, so that we can regard U as an orbifold coordinate chart around $(X, [\omega])$. By the definition of $\mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0})$, for each $(Y, \eta) \in U$, the curve $T_{1,0}^1$ is represented by a maximal cylinder C on (Y, η) . Using Theorem 5.5, by possibly shrinking U , we can assume that:

- $|\eta(\alpha_2)| < \frac{1}{d+1} \text{height}(C)$
- $|\eta(\beta_2)| < \frac{1}{d+1} \text{height}(C)$
- $\text{Im } \eta(\beta_1) > 0$

$X_{d^2}(T_{1,0})$ in local coordinates. We now turn to the question of when our $(X, [\omega])$ lies in the union of strata $X_{d^2}(T_{1,0})$ and what these strata look like in coordinates around $(X, [\omega])$. We will continue to work in the fixed neighborhood U of some $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})$ which we chose above.

For any (Y, η) in U , we have a marking $f: \Sigma_2 \rightarrow Y$, which is defined up to Dehn twist around $T_{1,0}$. Choosing a particular marking, we regard the symplectic basis $\{\alpha_i, \beta_i\}_{i=1}^2$ of $H_1(\Sigma_2; \mathbb{Z})$ as also a symplectic basis of $H_1(Y; \mathbb{Z})$.

Lemma 7.5. *Let $(Y, \eta) \in U \cap X_{d^2}$, and let $g: (Y, \eta) \rightarrow (E, \nu)$ be a primitive, degree d , branched cover of a genus one differential E . Then there is a basis $\{a, b\}$ of $H_1(E; \mathbb{Z})$ and integers p, q , and r such that $\gcd(d, p, q) = 1$, and*

$$\begin{aligned} g_*(\alpha_1) &= da & g_*(\beta_1) &= ra + b \\ g_*(\alpha_2) &= pa & g_*(\beta_2) &= qa \end{aligned}$$

Proof. There is some primitive $a \in H_1(E; \mathbb{Z})$ such that $g_*(\alpha_1) = na$ for some $n \in \mathbb{N}$. Since $\eta(\alpha_1) = 1$, we must have $\nu(a) > 0$. It follows that we can choose some $b \in H_1(E; \mathbb{Z})$ such that $\text{Im } \nu(b) > 0$, and such that $\{a, b\}$ form a basis of $H_1(E; \mathbb{Z})$. The cohomology class b is represented by some closed geodesic B on E .

We claim that

$$\text{Im } \nu(b) > \frac{1}{d+1} \text{height}(C),$$

where C is the maximal cylinder on Y homotopic to the single curve of $T_{1,0}$. To see this, consider $S = g^{-1}(A) \cap C$. The set S consists of r parallel closed curves $\{S_i\}_{i=1}^r$ on C . Since each point of B has d preimages under g , we must have $r \leq d$. Two consecutive curves S_i and S_{i+1} bound a cylinder C_i which has the same height as the cylinder obtained by cutting E along A , so $\text{height}(C_i) = \text{Im } \nu(b)$. There are also two more cylinders, C_0 and C_r bounded by S_1 and S_r respectively and the boundary curves of C . These cylinders have height less

than $\text{Im } \nu(b)$ since they are mapped injectively into E by g . Together, these cylinders fill out C , so

$$\text{height}(C) = \sum_{i=0}^r \text{height}(C_i) < (r+1) \text{Im } \nu(b) \leq (d+1) \text{Im } \nu(b),$$

which proves the claim.

It follows from this claim that $g_*(\alpha_2) = pa$ for some $p \in \mathbb{Z}$. To see this, assume that $g_*(\alpha_2) = pa + sb$ with $s \neq 0$, then

$$\begin{aligned} \frac{1}{d+1} \text{height}(C) &> |\eta(\alpha_2)| && \text{(by the definition of } U) \\ &= |\nu(g_*(\alpha_2))| \\ &\geq |s| \text{Im } \nu(b) && \text{(because } \nu(a) \text{ is real)} \\ &> \frac{1}{d+1} \text{height}(C), \end{aligned}$$

a contradiction. Similarly $g_*(\beta_2) = qa$ for some $q \in \mathbb{Z}$.

The map $g_*: H_1(Y; \mathbb{Z}) \rightarrow H_1(E; \mathbb{Z})$ is onto because $g: Y \rightarrow E$ is primitive. Since we've shown that each basis element of $H_1(Y; \mathbb{Z})$ except β_1 maps to a multiple of a , we must have $g_*(\beta_1) = ra \pm b$ for some $r \in \mathbb{Z}$, and because $\text{Im } \eta(\beta_1) > 0$ and $\text{Im } \nu(b) > 0$, we actually have $g_*(\beta_1) = xa + b$. It follows that

$$g_*(\alpha_1) \cdot g_*(\beta_1) = (na) \cdot (ra + b) = n,$$

and similarly $g_*(\alpha_2) \cdot g_*(\beta_2) = 0$. It follows easily from the fact that g is degree d that

$$g_*(\alpha_1) \cdot g_*(\beta_1) + g_*(\alpha_2) \cdot g_*(\beta_2) = d.$$

so $n = d$. Because g is primitive, we must have $\gcd(d, p, q) = 1$. ■

Let $\psi: U \rightarrow \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})$ be the unplumbing map defined in §6.3, which sends $(Y, \eta) \in U$ to the Abelian differential obtained by unplumbing the cylinder on (Y, η) homotopic to the curve $T_{1,0}^1$. In our local coordinates, ψ is the projection $\psi(w, x, y, z) = (w, x, y, 0)$. In $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})$, let $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$ be the inverse image of $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})(d)$, the locus of degree d cylinder covering differentials, under the natural projection.

Theorem 7.6. *An Abelian differential $(Y, \eta) \in U \setminus \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})$ is in X_{d^2} if and only if $\psi(Y, \eta) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$.*

Proof. Let $(Y, \eta) \in U$, and let C be the maximal closed cylinder on Y containing a closed geodesic A homotopic to the curve of $T_{1,0}$. Let (Z, ν) be the stable Abelian differential obtained by cutting Y along A to obtain a surface (Y', η') with geodesic boundaries A_1 and A_2 , and then gluing in an infinite cylinder to each resulting boundary component of Y' . The surface Y' naturally lies in both Y and Z , and (Z, ν) is the unplumbing of (Y, η) along C .

We need to show that (Y, η) is a degree d elliptic differential if and only if (Z, ν) is a degree d cylinder covering differential.

First, assume that (Z, ν) is a degree d cylinder covering differential. Let $g: (Z, \nu) \rightarrow (D, \xi)$ be a map to a cylinder $(D, \xi) = (\mathbb{C}/\mathbb{Z}, dz)$ realizing Z as a degree d cylinder covering differential. The map g sends the horizontal geodesics A_1 and A_2 to horizontal geodesics B_1 and B_2 on C , and by the Open Mapping Theorem, B_1 and B_2 must be distinct (or else the point $p \in Y'$ for which $g(p)$ has largest imaginary part would lie in the interior of Y' , which contradicts the Open Mapping Theorem). The geodesics B_1 and B_2 bound a subcylinder D' of D , and g maps Y' onto D' with degree d . The induced map in homology, $g_*: H_1(Y'; \mathbb{Z}) \rightarrow H_1(D'; \mathbb{Z})$ is onto because $g_*: H_1(Z; \mathbb{Z}) \rightarrow H_1(D; \mathbb{Z})$ is onto by primitivity of g . Now, we can recover Y by gluing together A_1 and A_2 by an appropriate gluing map. By gluing D' along B_1 and B_2 in a way compatible with the map g , we get an elliptic curve E with Abelian differential ζ and a primitive degree d map $h: (Y, \eta) \rightarrow (E, \zeta)$, realizing (Y, η) as a elliptic differential.

Now assume that (Y, η) is a degree d elliptic differential. Let $g: (Y, \eta) \rightarrow (E, \zeta)$ be a map realizing (Y, η) as an elliptic differential. Take a basis $\{a, b\}$ of $H_1(E; \mathbb{Z})$ as in Lemma 7.5. Let $B = g(A)$, a horizontal closed geodesic on E . The geodesics A and B are homologous to α_1 , and a respectively. Since $g_*(\alpha_1) = da$ by Lemma 7.5, the restriction $g|_A: A \rightarrow B$ is degree d , so $g^{-1}(B) = A$.

Cut E along B , and call the resulting cylinder with boundary (D', ξ') . Let (D, ξ) be the infinite cylinder obtained by gluing half-infinite cylinders to each boundary component of (D', ξ') . Since $g^{-1}(B) = A$, there is a natural degree d map $h': (Y', \eta') \rightarrow (D', \xi')$ and thus a degree d map $h: (Z, \nu) \rightarrow (D, \xi)$.

To show that (Z, ν) is a degree d cylinder covering differential, it just remains to show that h is primitive, and it is enough to show that $h'_*: H_1(Y'; \mathbb{Z}) \rightarrow H_1(D'; \mathbb{Z})$ is onto. This follows easily from Lemma 7.5. Since the homology classes α_i and β_2 can be represented by closed curves disjoint from A , they naturally define classes on Y' . Then the fact that $\gcd(d, p, q) = 1$ from Lemma 7.5, implies that h'_* is onto. $\blacksquare$

Corollary 7.7. *We have*

$$X_{d^2}(T_{1,0}) = \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})(d). \quad (7.1)$$

Given $(X, [\omega]) \in X_{d^2}(T_{1,0})$ contained in a sufficiently small neighborhood $U \subset \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0})$ with coordinates on U as defined above, $\overline{X}_{d^2} \cap U$ is cut out by the equations,

$$\begin{aligned} w &= \omega(\alpha_2)/\omega(\alpha_1) \\ x &= \omega(\beta_2)/\omega(\alpha_1), \end{aligned} \quad (7.2)$$

and $X_{d^2}(T_{1,0})$ is cut out by the additional equation $z = 0$. If ω has a double zero, then $\overline{W}_{d^2}$ is cut out by the equations (7.2) together with the additional equation $y = 0$.

In these coordinates, the foliation $\mathcal{A}_{d^2}$ of X_{d^2} has leaves given by $z = \text{const.}$

Proof. It follows immediately from Theorem 7.6 that $(w, x, y, z) \in U$ is in X_{d^2} if and only if $(w, x, y, 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$ and $z \neq 0$, and (7.1) follows by taking the closure.

If $(w, x, y, 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$, then $(w, x, y', 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$ for any y' because being a cylinder covering differential only depends on absolute periods. Furthermore, we must have $w, x \in \frac{1}{d}\mathbb{Z}$ by Lemma 7.5, so the coordinates w and x are locally constant along $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$. It follows that if $(X, [\omega]) = (w, x, y, 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$, then around this point, $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0})(d)$ is cut out by the equations (7.2) together with $z = 0$, and X_{d^2} is cut out by the equations (7.2).

The equations for $X_{d^2}(T_{1,0})$ and $\overline{W}_{d^2}$ are clear from the definitions of the coordinates. The leaves of the foliation $\mathcal{A}_{d^2}$ are determined by the condition that the absolute periods are locally constant along the leaves, which means in these coordinates that v , w , x , and z are all constant. Since we're in U_1 , we have $v = 1$, and w and x are locally constant by (7.2). Thus the foliation is given by $z = \text{const}$. ■

Theorem 7.4 follows directly from this corollary. Note that the statement in Theorem 7.4 that the intersections of $\overline{W}_{d^2}$ and $X_{d^2}(T_{1,0})$ are transverse if $d > 3$ follows from the coordinates above together with the fact that $\Gamma_1(d)$ is torsion-free if $d > 3$.

A section of $\mathcal{O}(-1)$. Recall that we normalized each projective class $(Y, [\eta]) \in \mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0})$ so that each $\eta(\alpha_1) = 1$. We can regard this as defining a section s of the canonical line bundle $\mathcal{O}(-1)$ over $\mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0})$.

Proposition 7.8. *The norm of this section s of $\mathcal{O}(-1)$ is given by,*

$$h(s, s) = -\frac{1}{2\pi} \log |z| + wx. \quad (7.3)$$

Proof. For any genus two Riemann surface X with a symplectic basis $\{\alpha_i, \beta_i\}$ of $H_1(X; \mathbb{Z})$, it is well known that for any $\omega \in \Omega(X)$,

$$\int_X |\omega|^2 = \text{Im}(\overline{\omega(\alpha_1)}\eta(\beta_1) + \overline{\omega(\alpha_2)}\eta(\beta_2)). \quad (7.4)$$

Equation (7.3) follows from this formula using the fact that w and x are real on $\overline{X}_{d^2}$ in these coordinates. ■

7.4 The stratum $X_D(T_{1,0}, T_{1,0}^1)$

We now study the stratum $X_D(T_{1,0}, T_{1,0}^1)$ of Abelian differentials in $\overline{X}_D$ with one nonseparating holomorphic node. Since this stratum is empty if D is not square, we will restrict to the case $D = d^2$.

Recall that the d -torsion locus $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$ is the locus of $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$ such that, if we regard $(X, [\omega])$ as a genus one differential with two marked points, then the marked points differ by exactly d -torsion on X . The goal of this section is to prove the following theorem.

Theorem 7.9. $\overline{X}_{d^2}$ is an orbifold around the stratum $X_{d^2}(T_{1,0}, T_{1,0}^1)$, which is a suborbifold equal to

$$\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d) \cong \mathbb{H}/\Gamma_1(d).$$

Vanishing of periods. We now prove that for Abelian differentials $(X, [\omega]) \in X_{d^2}$ close to some $(X_0, [\omega_0]) \in \overline{X}_{d^2}$ which has a holomorphic node, periods along curves of $(X, [\omega])$ which are close to a holomorphic node of $(X_0, [\omega_0])$ must vanish.

Proposition 7.10. Let S be a curve system on Σ_2 , let $\pi: \mathbb{P}\Omega\mathcal{D}_2(S) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$ be the natural projection, and let $(X_0, [\omega_0]) \in \mathbb{P}\Omega\mathcal{D}_2(S) \cap \pi^{-1}(\overline{X}_{d^2})$ have a nonseparating holomorphic node represented by a homology class $\alpha \in H_1(\Sigma_2; \mathbb{Z})$. Then for $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{D}_2(S)$ sufficiently close to $(X_0, [\omega_0])$, we must have $\omega(\alpha) = 0$.

Proof. Choose a representative ω_0 of the projective class $[\omega_0]$, and suppose there is a sequence $(X_n, \omega_n) \rightarrow (X_0, \omega_0)$ with $(X_n, \omega_n) \in \pi^{-1}(X_{d^2})$ and $\omega_n(\alpha) \neq 0$ for all n . There are degree d branched covers $f_n: (X_n, \omega_n) \rightarrow (E_n, \eta_n)$ over elliptic curves. Taking a subsequence, we can assume that $(E_n, \eta_n) \rightarrow (E, \eta)$, with E either a cylinder or an elliptic curve and that f_n converges to a branched cover $f: (X_0, \omega_0) \rightarrow (E, \eta)$ of degree at most d . We must then have $\eta \neq 0$.

The injectivity radius $I(E_n, \eta_n)$ of (E_n, η_n) is bounded by the absolute value of any nonzero period of η . Since $\eta((f_n)_*\alpha) = \omega_n(\alpha)$, and $\omega_n(\alpha) \rightarrow 0$, this means that $I(E_n, \eta_n) \rightarrow 0$. This is a contradiction because injectivity radius is continuous, and (E, η) has nonzero injectivity radius. $\blacksquare$

Local coordinates. Choose some $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)$. By choosing a marking $f: \Sigma_2 \rightarrow X$, we can regard $(X, [\omega])$ as a point in $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$. Let $\{\alpha_i, \beta_i\}_{i=1}^2$ be a symplectic basis of $H_1(\Sigma_2; \mathbb{Z})$ such that α_1 represents the curve $T_{1,0}^1$, as in §6.4.

Let $U \subset \mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$ be a neighborhood of $(X, [\omega])$ small enough that the natural map $U/\text{Aut}(X, [\omega]) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$ is an isomorphism onto its image, and $\eta(\alpha_2) \neq 0$ for every $(Y, [\eta]) \in U$. By normalizing each $(Y, [\eta]) \in U$ so that $\eta(\alpha_2) = 1$, we can identify U with an open set U_1 in the hypersurface in $\Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$ of those Abelian differentials (Z, ζ) such that $\zeta(\alpha_2) = 1$. The coordinates (w, x, y, z) from Proposition 6.6 then restrict to local coordinates (x, y, z) on U .

$X_{d^2}(T_{1,0}, T_{1,0}^1)$ in local coordinates. The set U is not a neighborhood of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0})$, but rather it is a neighborhood of $(X, [\omega])$ in the hypersurface $\mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$. Proposition 7.10 implies that in a sufficiently small neighborhood V of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0})$,

$$V \cap X_{d^2} \subset U.$$

This means that we can restrict to this hypersurface and still understand the local structure of $\overline{X}_{d^2}$ around $(X, [\omega])$.

Let $\psi: U \rightarrow \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$ be the projection (defined in the proof of Proposition 6.6) which splits each $(X, [\omega]) \in U$ along a union of saddle connections $I \cup J(I)$ which is homotopic to α_1 . In our local coordinates on U , this is the map $\phi(x, y, z) = (x, y, 0)$.

Let $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)(d)$ be the inverse image under the natural projection of the d -torsion locus $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$.

Theorem 7.11. *A $(Y, \eta) \in U \setminus \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$ is in X_{d^2} if and only if $\psi(Y, \eta) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)(d)$.*

Proof. Let $(Y, \eta) \in U$, with saddle connections I_1 and I_2 , which we split along to form an elliptic curve (E, ν) . Let p and q be the two zeros of η . Let J_1 and J_2 be the two segments which are the images of the saddle connections I_i in E . The J_i start at points p_i and end at points q_i such that upon taking the connected sum along J_1 and J_2 , the p_i are identified to form the zero p and the q_i are identified to form the zero q (see Figure 11).




Figure 11: Split along $I_1 \cup I_2$ and then reglue.

We need to show that (Y, η) is a degree d elliptic differential if and only if the points p_i differ by exactly d -torsion in the group law on E .

First, suppose that (Y, η) is a degree d elliptic differential. Then (Y, η) is branched over an elliptic curve by some $g: (Y, \eta) \rightarrow (F, \xi)$. The saddle connections I_i must have as their image the same segment K in F because they have the same direction and length, and they start at the same zero p . It follows that when we cut Y along I_1 and I_2 , points which are then glued together to form E map to the same point of F under g . Thus g defines an isogeny $g': E \rightarrow F$ of the same degree, sending p_1 and p_2 to the same point. This implies that p_1 and p_2 differ by d -torsion on E .

Conversely, if p and q differ by d -torsion, then there is a degree d -isogeny $g: (E, \nu) \rightarrow (F, \xi)$; it must send points of J_1 and J_2 which are glued together to form Y to the same point of Y , so we get a branched cover $(Y, \eta) \rightarrow (F, \xi)$ which realizes η as a degree d elliptic differential. ■

Corollary 7.12. *The stratum $X_{d^2}(T_{1,0}, T_{1,0}^1)$ is exactly $\mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$. If we have $(X, [\omega]) \in X_{d^2}(T_{1,0}, T_{1,0}^1)$, then in the (x, y, z) coordinates defined above on the neighborhood U of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{1,0}, T_{1,0}^1)$, the variety $\overline{X}_{d^2}$ is cut out by the equation,*

$$dy = ax + b, \quad (7.5)$$

for some relatively prime integers a and b .

In these coordinates, the stratum $X_{d^2}(T_{1,0}, T_{1,0}^1)$ is cut out by the additional equation $z = 0$, and leaves of the foliation $\mathcal{A}_{d^2}$ are determined by either of the equivalent equations, $x = \text{const}$ or $y = \text{const}$.

Proof. It follows immediately from Theorem 7.11 that $(x, y, z) \in U$ is in X_{d^2} if and only if $(x, y, 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)(d)$ and $z \neq 0$. Thus

$$U \cap \overline{X}_{d^2} = \{(x, y, z) \in U : (x, y, 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)(d)\},$$

and $X_{d^2}(T_{1,0}, T_{1,0}^1) = \mathbb{P}\Omega\mathcal{M}_2^0(T_{1,0}, T_{1,0}^1)(d)$ as claimed.

It remains to show that if $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)(d)$, then near $(X, [\omega])$, the locus $\mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)(d)$ is cut out by the equation (7.5). This is because for some $(Y, \eta) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{1,0}, T_{1,0}^1)$, a relative period joining the marked points is given by $y = \eta(\beta_1)$, so the marked points differ by d -torsion if and only if $d\eta(\beta_1) = a\eta(\beta_2) + b\eta(\alpha_2)$ for some relatively prime integers a and b , which is equivalent to (7.5).

The equation for $X_{d^2}(T_{1,0}, T_{1,0}^1)$ is clear from the definition of the coordinates, and the equations for the leaves of $\mathcal{A}_{d^2}$ hold because either of these equations together with (7.5) implies that the absolute periods are all constant. ■

Theorem 7.9 follows immediately from this corollary.

7.5 The stratum $X_D(T_{2,0}, T_{2,0}^2)$

We now study the stratum $X_D(T_{2,0}, T_{2,0}^2)$ consisting of limits of eigenforms with one nonseparating polar node and one nonseparating holomorphic node. The arguments in this section are straightforward combinations of those in the previous two sections, so we will omit some of the proofs. This stratum is empty when D is not square, so we will restrict to the case $D = d^2$.

Recall that in §6.5 we defined the finite set $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d)$, consisting of the stable Abelian differentials $f_{q/d} \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$, where q is an integer relatively prime to d . The following is the main theorem of this section.

Theorem 7.13. *The stratum $X_{d^2}(T_{2,0}, T_{2,0}^2)$ is the locus $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d)$. The variety $\overline{X}_{d^2}$ is nonsingular at these points, and each $f_{q/d} \in X_{d^2}(T_{2,0}, T_{2,0}^2)$ is a transverse intersection point of the closures of the strata $X_{d^2}(T_{1,0})$ and $X_{d^2}(T_{1,0}, T_{1,0}^1)$.*

Local coordinates. Choose some $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)$. By choosing a marking $f: \Sigma_2 \rightarrow X$, we can regard $(X, [\omega])$ as a point in $\mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)$. Let $\{\alpha_i, \beta_i\}_{i=1}^2$ be a symplectic basis of $H_1(\Sigma_2; \mathbb{Z})$ such that α_i represents the curve $T_{2,0}^i$ in $T_{2,0}$ as in §6.5.

Let $U \subset \mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$ be a neighborhood of $(X, [\omega])$ small enough that the natural map $U/\text{Aut}(X, [\omega]) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$ is an isomorphism onto its image. By normalizing each $(Y, [\eta]) \in U$ so that $\eta(\alpha_1) = 1$, we can identify U with an open set U_1 in the hypersurface in $\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$ consisting of those Abelian differentials (Z, ζ) such that $\zeta(\alpha_1) = 1$. The coordinates (w, x, y, z) from Proposition 6.7 then restrict to coordinates (x, y, z) on U .

$X_{d^2}(T_{2,0}, T_{2,0}^2)$ in local coordinates. Let $\psi: U \rightarrow \mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)$ be the projection which sends an Abelian differential (Y, η) to the one obtained by unplumbing the cylinder on (Y, η) homotopic to the curve $T_{2,0}^1$ and splitting along the pair of saddle connections $I \cup J(I)$ homotopic to $T_{1,0}^2$. In local coordinates, $\psi(x, y, z) = (x, 0, 0)$. Let $\mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)(d)$ be the inverse image under the natural projection to $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d)$.

Theorem 7.14. *If the neighborhood U of $(X, [\omega])$ is sufficiently small, then a nonsingular $(Y, \eta) \in U$ is in X_{d^2} if and only if $\phi(Y, \eta) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)(d)$.*

The proof of this theorem is a straightforward combination of the proofs of Theorems 7.6 and 7.11, so we will omit the proof. The idea is that $f_{q/d}$ is a primitive degree d branched cover of a cylinder, and given a (Y, η) such that $\psi(Y, \eta) = f_{q/d}$, we can use this cover to exhibit (Y, η) as a degree d torus cover.

Corollary 7.15. *The stratum $X_{d^2}(T_{2,0}, T_{2,0}^2)$ is equal to $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d)$. If $(X, [\omega]) \in X_{d^2}(T_{2,0}, T_{2,0}^2)$, then in a neighborhood V of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$,*

$$V \cap \overline{X}_{d^2} \subset \mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2).$$

In the coordinates (x, y, z) on a neighborhood U of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$, the variety $\overline{X}_{d^2}$ is cut out by the equation,

$$x = \frac{q}{d}, \tag{7.6}$$

for some integer q relatively prime to d .

The intersection of the closure of $X_{d^2}(T_{1,0})$ with U is cut out by the additional equation $y = 0$, and the intersection of the closure of $X_{d^2}(T_{1,0}, T_{1,0}^1)$ with U is cut out by the additional equation $z = 0$. The leaves of the foliation $\mathcal{A}_{d^2}$ of X_{d^2} are given by $y = \text{const}$.

Proof. Proposition 7.10 implies directly that in a neighborhood V of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$,

$$V \cap \overline{X}_{d^2} \subset \mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2).$$

It follows immediately from Theorem 7.14 that $(x, y, z) \in U$ is in X_{d^2} if and only if $(x, 0, 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)(d)$ and $y, z \neq 0$. Thus

$$U \cap \overline{X}_{d^2} = \{(x, y, z) \in U : (x, 0, 0) \in \mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}, T_{2,0}^2)(d)\},$$

and $X_{d^2}(T_{2,0}, T_{2,0}^2) = \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d)$ as claimed.

Suppose $(X, \omega) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}, T_{2,0}^2)(d)$. The coordinate $x = \omega(\beta_2)$ measures a relative period on (X, ω) joining the two marked points which are identified to form a holomorphic node. Since $\omega(\alpha_1) = 1$, we have $x = \omega(\beta_2) = q/d$ for some integer q relatively prime to d if and only if the marked points differ by d -torsion. Thus $\overline{X}_{d^2}$ is cut out by (7.6) as claimed.

The intersection of the closure of $X_{d^2}(T_{1,0})$ with U is cut out by the additional equation $y = 0$ because $y(Y, \eta) = 0$ exactly when (Y, η) has a polar node, and the intersection of the closure of $X_{d^2}(T_{1,0}, T_{1,0}^1)$ with U is cut out by the additional equation $z = 0$ because $z(Y, \eta) = 0$ exactly when (Y, η) has a holomorphic node.

The equation $y = \text{const}$ defines the foliation $\mathcal{A}_{d^2}$ because this makes the period along β_1 constant, and we have already seen that the other periods are locally constant along X_{d^2} with our normalization for η . $\blacksquare$

Theorem 7.13 follows directly from this corollary.

A section of $\mathcal{O}(-1)$. Recall that we normalized each projective class $(Y, [\eta]) \in \mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$ so that each $\eta(\alpha_1) = 1$. We can regard this as defining a section s of the canonical line bundle $\mathcal{O}(-1)$ over $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0}, T_{2,0}^2)$.

The following follows directly from (7.4) together with the definition of our coordinates.

Proposition 7.16. *The norm of this section s of $\mathcal{O}(-1)$ is given by,*

$$h(s, s) = -\frac{1}{2\pi} \log |z|. \quad (7.7)$$

7.6 The strata $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, and $X_D(T_{2,1})$

We now turn to the strata $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, and $X_D(T_{2,1})$ which together consist of those limiting eigenforms which have exactly two nonseparating polar nodes, possibly with a separating node.

Recall the definitions from §6.6 of the curves C_λ and the points p_λ and w_λ in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$. The following is the classification of points in the strata $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, and $X_D(T_{2,1})$, which we will prove over the course of this subsection.

Theorem 7.17. *The union of the three strata $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, and $X_D(T_{2,1})$ is equal to the union,*

$$\bigcup_{\lambda} C_\lambda, \quad (7.8)$$

where the union is over all λ such that $\lambda = \lambda(P)$ for some nondegenerate Y_D -prototype P . The stratum $X_D(T_{2,0}; 2)$ is the finite set,

$$\bigcup_{\lambda} w_\lambda,$$

where the union is over all λ such that $\lambda = \lambda(P)$ for some W_D -prototype P . The stratum $X_D(T_{2,1})$ is the finite set,

$$\bigcup_{\lambda} p_{\lambda},$$

where the union is over all λ such that $\lambda = \lambda(P)$ for some P_D -prototype P . The stratum $X_D(T_{2,0}; 1, 1)$ is the complement of these finite sets in (7.8).

We will also see that in general $\overline{X}_D$ is singular along the curves C_{λ} , and we will give equations for $\overline{X}_D$ in local coordinates around C_{λ} .

Local coordinates. Choose an $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}; 1, 1)$, $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}; 2)$ or $\mathbb{P}\Omega\mathcal{M}_2^0(T_{2,1})$. By choosing a marking $f: \Sigma_2 \rightarrow X$, we can regard $(X, [\omega])$ as a point in $\mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}; 1, 1)$, $\mathbb{P}\Omega\mathcal{T}_2^0(T_{2,0}; 2)$, or $\mathbb{P}\Omega\mathcal{T}_2^0(T_{2,1})$.

Our first goal is to show that if $(X, [\omega]) \in \overline{X}_D$, then $(X, [\omega])$ lies in the union (7.8). If $(X, [\omega]) \in \overline{X}_D$, then ω is an eigenform for real multiplication of $\mathcal{O}_D$ on $\text{Jac}(X)$ by Theorem 5.10, so assume that it is such an eigenform. It follows that the ratio of the residues of the two poles of ω is real. Thus we can choose a representative ω of the projective class $[\omega]$ and choose homology classes α_1 and α_2 in $H_1(\Sigma_2, \mathbb{Z})$ such that the following properties hold:

- The α_i represent the two nonseparating nodes of X .
- $\omega(\alpha_1) = 1$.
- $\omega(\alpha_2) = \lambda$ with $\lambda \geq 1$.

If $\lambda > 1$, then this choice of the α_i and ω is uniquely determined up to the hyperelliptic involution by these properties. If $\lambda = 1$, then there are two possible choices for the α_i and ω up to the hyperelliptic involution, and we choose one arbitrarily (the other one is then obtained by swapping the α_i).

Extend $\{\alpha_1, \alpha_2\}$ to a basis $\{\alpha_i, \beta_i\}_{i=1}^2$ of $H_1(\Sigma_2; \mathbb{Z})$ as follows:

- If $(X, \omega) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}; 1, 1)$, let β_i be an arbitrary pair which is dual to the α_i with respect to the intersection pairing.
- If $(X, \omega) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,1})$, let β_i be a pair dual to the α_i such that each β_i is represented by a simple closed curve disjoint from the separating curve $T_{2,1}^3$. This determines each β_i up to adding a multiple of α_i .
- If $(X, \omega) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{2,0}; 2)$, then the horizontal foliation of (X, ω) consists of two horizontal cylinders separated by a “figure-eight”; let $F \subset \Sigma_2$ be the inverse image of this figure-eight with p the singular point of F . In $H_1(\Sigma_2; \mathbb{Z})$, let the β_i be a pair which is dual to the α_i such that β_1 is represented by a simple closed curve which passes through p and is disjoint from $F \setminus p$, and β_2 is represented by a simple closed curve which is disjoint from F (see Figure 12). These conditions determine each β_i uniquely up to adding a multiple of α_i .




Figure 12: Symplectic basis for $H_1(\Sigma_2; \mathbb{Z})$.

Let U be a small neighborhood of (X, ω) in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0})$ or $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,1})$ such that the natural map $U/\text{Aut}(X, \omega) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$ is an isomorphism onto its image. Also choose U small enough so that the conclusion of Theorem 5.9 holds. That is, if $(Y, \eta) \in U$ is an eigenform for real multiplication by $\mathcal{O}_D$, then the induced action of $\mathcal{O}_D$ on $H_1(\Sigma_2; \mathbb{Z})$ preserves the subgroup S spanned by the α_i , giving S the structure of an $\mathcal{O}_D$ -module.

By normalizing each projective class $(Y, [\eta]) \in U$ so that $\eta(\alpha_1) = 1$, we can identify U with an open set U' in the hypersurface in $\Omega\mathcal{D}_2^0(T_{2,0})$ or $\Omega\mathcal{D}_2^0(T_{2,1})$ of those Abelian differentials (Z, ζ) such that $\zeta(\alpha_1) = 1$. For any $\mu \in \mathbb{R}$, let

$$U_\mu = \{(Y, \eta) \in U : \eta(\alpha_1) = 1, \text{ and } \eta(\alpha_2) = \mu\},$$

a hypersurface in U . We then have $(X, \omega) \in U_\lambda$.

Proposition 7.18. *If the neighborhood U of (X, ω) is taken sufficiently small, then for each $(Y, \eta) \in U \cap X_D$, the $\mathcal{O}_D$ -submodule S of $H_1(\Sigma_2; \mathbb{Z})$ is a quasi-invertible $\mathcal{O}_D$ -module with $\{\alpha_1, \alpha_2\}$ an admissible basis; furthermore, $U \cap X_D \subset U_\lambda$.*

It follows that $\lambda = \lambda(P)$ for some nondegenerate Y_D -prototype P .

Proof. For an Abelian differential $(Y, \eta) \in U$ which is sufficiently close to (X, ω) , the classes α_i will be homologous to core curves of very tall horizontal cylinders on Y by Theorem 5.5. Since $\alpha_i \cdot \beta_i = 1$, a curve representing β_i passes vertically through the cylinder, so the cylinder makes a large positive contribution to $\text{Im } \eta(\beta_i)$. Thus, if $(Y, \eta) \in U$ with U sufficiently small, we must have $\text{Im } \eta(\beta_i) > 0$.

The intersection pairing is an unimodular pairing between the $\mathcal{O}_D$ -modules S and $H_1(\Sigma_2; \mathbb{Z})/S$, and the α_i and β_i represent bases of these modules dual with respect to this pairing. The period maps associated to η and $\text{Im } \eta$ are nonzero ι_1 -linear maps of these modules to $\mathbb{R}$ which send the α_i and the β_i

respectively to positive reals. It then follows from Theorem 3.5 that the α_i form an admissible basis of S .

By Proposition 3.4, for any $(Y, \eta) \in U \cap X_D$ there are integers a, b and c as in the conclusion of that proposition with $a\eta(\alpha_2)^2 + b\eta(\alpha_2) + c = 0$. Since there are only finitely many such integers, and $\eta(\alpha_2) \rightarrow \lambda$ as $(Y, \eta) \rightarrow (X, \omega)$, we must have $\eta(\alpha_2) = \lambda$ for (Y, η) sufficiently close to (X, ω) .

Finally, $\lambda = \lambda(P)$ for any Y_D -prototype $(a, b, c, \bar{q})$ with a, b , and c as above. $\blacksquare$

Corollary 7.19. *The strata $X_D(T_{2,0}; 1, 1)$, $X_D(T_{2,0}; 2)$, and $X_D(T_{2,1})$ are contained in the union (7.8).*

In the coordinates (v, w, x, y, z) on $\Omega\mathcal{D}_2^0(T_{2,0})$ or $\Omega\mathcal{D}_2^0(T_{2,1})$ from §6.6, the subspace U_λ is cut out by the equations,

$$v = 1 \text{ and } w = \lambda,$$

so the (x, y, z) restrict to give local coordinates on U_λ , which send $(Y, \eta) \in U$ to

$$\begin{aligned} x &= \begin{cases} \int_I \eta & \text{if } (X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,0}; 1, 1); \\ \left(\int_I \eta\right)^{2/3} & \text{if } (X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,0}; 2); \\ \left(\int_I \eta\right)^2 & \text{if } (X, \omega) \in \Omega\mathcal{T}_2^0(T_{2,1}), \end{cases} \\ y &= e^{2\pi i \eta(\beta_1)} \\ z &= e^{2\pi i \eta(\beta_2)/\lambda}, \end{aligned}$$

where I is a saddle connection joining distinct zeros as in §6.6.

Prototypes. Now assume that our $(X, [\omega]) \in C_\lambda$, where $\lambda = \lambda(P)$ for some nondegenerate Y_D -prototype P with a small neighborhood U of $(X, [\omega])$ in the Dehn space chosen as above. We will assign a prototype $P(Y, \eta)$ to each Abelian differential $(Y, \eta) \in U \cap X_D$. For now, fix such an (Y, η) .

Recall that we identify two terminal prototypes if they differ by the involution (3.1). We temporarily don't want to make this identification, so we will call a *fine prototype* a prototype which is defined exactly as in §3.1, except without this identification.

The marking $\Sigma_2 \rightarrow Y$ is well-defined up to Dehn twist around the curves of $T_{2,0}$ or $T_{2,1}$. Choose a particular marking. This allows us to consider the symplectic basis $\{\alpha_i, \beta_i\}$ of $H_1(\Sigma_2; \mathbb{Z})$ as a basis of $H_1(Y; \mathbb{Z})$ as well.

By Propositions 3.4 and 7.18, there is a unique μ in K_D so that $\mu \cdot \alpha_1 = \alpha_2$, using the real multiplication on $\text{Jac}(Y)$. Since η is an eigenform, we know that $\mu^{(1)} = \lambda$ because

$$\lambda = \eta(\alpha_2) = \eta(\mu \cdot \alpha_1) = \mu^{(1)}\eta(\alpha_1) = \mu^{(1)}.$$

If D is not square, this means that μ doesn't depend on $(Y, \eta) \in U$ because μ is determined by its first embedding. If D is square, however, there can be Abelian differentials close to (X, ω) with different $\mu^{(2)}$.

Let a , b , and c be the integers given by Proposition 3.4. Since $\mathcal{O}_D = \mathbb{Z}[a\mu]$, the algebraic integer $a\mu$ acts on $H_1(Y; \mathbb{Z})$ by an integer matrix.

Proposition 7.20. *The matrix T of the action of $a\mu$ on $H_1(Y; \mathbb{Z})$ in the basis $\{\alpha_1, \alpha_2, \beta_1, \beta_2\}$ is*

$$T = \begin{pmatrix} 0 & -c & 0 & q \\ a & -b & -q & 0 \\ 0 & 0 & 0 & a \\ 0 & 0 & -c & -b \end{pmatrix} \quad (7.9)$$

with a , b , and c as above, and for some integer q .

Proof. The matrix T is self-adjoint with respect to the intersection pairing on $H_1(Y; \mathbb{Z})$, which is equivalent to

$$T^t J = J T,$$

where

$$J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}.$$

This easily implies that

$$T = \begin{pmatrix} A & Q \\ 0 & A^t \end{pmatrix}$$

with $Q = -Q^t$.

To calculate A , we know that

$$a\mu \cdot \alpha_1 = a\alpha_2,$$

and

$$a\mu\alpha_2 = a\mu^2\alpha_1 = (-b\mu - c)\alpha_1 = -c\alpha_1 - b\alpha_2,$$

so

$$A = \begin{pmatrix} 0 & -c \\ a & -b \end{pmatrix}.$$

■

Now let $P(Y, \eta) = (a, b, c, \bar{q})$, where q is as in the above Proposition, and $\bar{q}$ is the residue class of q , taken modulo $\gcd(a, b, c)$. Similarly, if (X, ω) has a double zero or a separating node, let $P'(Y, \eta) = (a, b, c, \bar{q})$, where $\bar{q}$ is the residue class of q , taken modulo $\gcd(a, c)$. Note that the definitions of $P(Y, \eta)$ and $P'(Y, \eta)$ involved some choices: there was a choice of which β_i to take as a basis dual to the α_i in $H_1(\Sigma_2; \mathbb{Z})$, and there was a choice of a marking $\Sigma_2 \rightarrow Y$ to take up to Dehn twist around $T_{2,0}$ or $T_{2,1}$ which we used to transport the basis of $H_1(\Sigma_2; \mathbb{Z})$ to $H_1(Y; \mathbb{Z})$.

Theorem 7.21. *The quadruple $P(Y, \eta)$ is a fine Y_D -prototype which does not depend on the choice of the β_i or the marking $\Sigma_2 \rightarrow Y$. If (X, ω) has a double zero or a separating node, then $P'(Y, \eta)$ is a fine W_D or P_D -prototype which is also independent of these choices.*

Proof. We first show that $P(Y, \eta)$ satisfies the required properties of a prototype. It follows directly from Proposition 3.4 that $b^2 - 4ac = D$ and $a > 0$. We know $N_{\mathbb{Q}}^{K_D}(\mu) < 0$ because $\{\alpha_1, \alpha_2\}$ is an admissible basis of the $\mathcal{O}_D$ -module $S = \langle \alpha_1, \alpha_2 \rangle$, so $c < 0$. Since $\mathbb{Z}[a\mu] = \mathcal{O}_D$ by Proposition 3.4, the matrix in (7.9) must be primitive because the action of $\mathcal{O}_D$ on $\text{Jac}(Y)$ is proper; thus $\gcd(a, b, c, \bar{q}) = 1$. Finally, $a + b + c \leq 1$ because $\mu^{(1)} \geq 1$.

Now we will show that $P(Y, \eta)$ is independent of the choices made. Replacing the β_i with a new pair which are dual to the α_i or changing the marking $\Sigma_2 \rightarrow Y$ by Dehn twists around $T_{2,0}$ or $T_{2,1}$ produces a new pair β'_i of the form,

$$\begin{aligned}\beta'_1 &= \beta_1 + r\alpha_1 + s\alpha_2 \\ \beta'_2 &= \beta_2 + s\alpha_1 + t\alpha_2\end{aligned}\tag{7.10}$$

for integers r , s , and t .

The matrix sending the basis $\{\alpha_1, \alpha_2, \beta_1, \beta_2\}$ to $\{\alpha_1, \alpha_2, \beta'_1, \beta'_2\}$ is

$$S = \begin{pmatrix} 1 & 0 & r & s \\ 0 & 1 & s & t \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

If T is the matrix of the endomorphism of $H_1(Y; \mathbb{Z})$ given by multiplication by $a\lambda$ in the old basis. and T' is the matrix of the same endomorphism in the new basis, then T' is

$$T' = S \cdot T \cdot S^{-1},$$

which is

$$\begin{pmatrix} 0 & -c & 0 & q' \\ a & -b & -q' & 0 \\ 0 & 0 & 0 & a \\ 0 & 0 & -c & -b \end{pmatrix},$$

where

$$q' = q + ar - bs + ct.\tag{7.11}$$

It follows that $q' \equiv q \pmod{\gcd(a, b, c)}$, so the prototype is independent of the choices.

Now suppose (X, ω) does have a separating node or a double zero. We must now show that q is well-defined modulo $\gcd(a, c)$. In this case, the choice of β_i is uniquely determined up to adding a multiple of α_i because we imposed additional requirements on the β_i . That means that different choices give a new pair $\{\beta'_i\}$ of the form (7.10) with $s = 0$. Then from (7.11), we see that $q' \equiv q \pmod{\gcd(a, c)}$.

Finally, if (X, ω) has a double zero, then $\lambda > 1$ by Proposition 6.10, so $a + b + c < 0$, and $P'(Y, \eta)$ is then a W_D -prototype. $\blacksquare$

Remark. If $\lambda = 1$, then there were also two possible ways to choose the α_i . In this case, one can check that swapping the α_i has the effect of applying the involution (3.1) to the prototypes $P(Y, \eta)$ and $P'(Y, \eta)$. Thus they are independent of this choice as ordinary prototypes but not as fine prototypes.

Equations for $\overline{X}_D$ in local coordinates. Given $(X, [\omega]) \in C_\lambda$ and a small neighborhood U of $(X, [\omega])$ in the Dehn space as above, we now give equations for $\overline{X}_D$ in U and show that $(X, [\omega])$ is indeed in $\overline{X}_D$. Since the prototype $P(Y, \eta)$ is locally constant on $U \cap X_D$, we can use these prototypes to label branches of $U \cap X_D$. Given a fine Y_D -prototype P , let $V_P \subset U$ be the closure in U of

$$\{(Y, \eta) \in U \cap X_D : P(Y, \eta) = P\},$$

which in fact lies in U_λ by Proposition 7.18. If (X, ω) has a double zero or a separating node, and P is a fine W_D or P_D -prototype (respectively), then let V'_P be the analogous locus where $P'(Y, \eta) = P$. In this case, given a fine W_D -prototype (or P_D -prototype) P ,

$$V_P = \bigcup_Q V'_Q,$$

where the union is over all fine W_D -prototypes (or P_D -prototypes) which map to P .

Let

$$(a', b', c', \bar{q}') = (a, b, c, \bar{q}) / \gcd(a, b, c),$$

and

$$(a'', b'', c'', \bar{q}'') = (a, b, c, \bar{q}) / \gcd(a, c)$$

(here we interpret $\bar{q}'$ and $\bar{q}''$ as elements of $\mathbb{Q}/\mathbb{Z}$).

Recall that we have coordinates (x, y, z) on U_λ defined above.

Theorem 7.22. *The locus V_P is cut out by the equation,*

$$y^{a'} = e^{-2\pi i \bar{q}'} z^{-c'}, \quad (7.12)$$

and C_λ is cut out by the equations $y = z = 0$.

If (X, ω) has a double zero or a separating node, then V'_P is cut out by the equation,

$$y^{a''} = e^{-2\pi i \bar{q}''} z^{-c''}, \quad (7.13)$$

and $\overline{W}_D \cap V_P$ or respectively $\overline{P}_D \cap V_P$ is cut out by the additional equation $x = 0$. The leaves of the foliation $\mathcal{A}_D$ of X_D are given in these coordinates by either of the equivalent equations $y = \text{const}$ or $z = \text{const}$.

Proof. The proofs of (7.12) and (7.13) essentially the same, so we will only give the proof of (7.12).

Choose some $(Y, \eta) \in V_P$ with a symplectic basis $\{\alpha_i, \beta_i\}$ and $\mu \in K_D$ as before. Equation (7.9) implies

$$a\mu \cdot \beta_1 = -q\alpha_2 - c\beta_2 \quad (7.14)$$

$$a\mu \cdot \beta_2 = q\alpha_1 + a\beta_1 - b\beta_2. \quad (7.15)$$

Since η is an eigenform, integrating η over both sides gives

$$a\lambda\eta(\beta_1) = -q\lambda - c\eta(\beta_2) \quad (7.16)$$

$$a\lambda\eta(\beta_2) = q + a\eta(\beta_1) - b\eta(\beta_2) \quad (7.17)$$

because $\mu^{(1)} = \lambda$. Dividing (7.16) by $\lambda \gcd(a, b, c)$ and exponentiating yields (7.12).

Now consider (Y, η) which lies in $U_\lambda \setminus V(y, z)$ and in the locus defined by (7.12) (using the notation $V(\cdot)$ for the zero locus of the enclosed functions). We must show that (Y, η) is an eigenform for real multiplication by $\mathcal{O}_D$ with prototype $P = (a, b, c, \bar{q})$.

Taking a logarithm of (7.12), we get

$$a\lambda\eta(\beta_1) \equiv -\lambda\bar{q} - c\eta(\beta_2) \pmod{\lambda \gcd(a, b, c)}.$$

We can then choose some $q \equiv \bar{q} \pmod{\gcd(a, b, c)}$ so that (7.16) holds. It then follows automatically from the relation $a\mu^2 + b\mu + c = 0$ that (7.17) holds. We can then define real multiplication of $\mathcal{O}_D$ on $\text{Jac}(Y)$, by defining $a\mu$ to act via the matrix T in (7.9). Equations (7.16) and (7.17) then imply (Y, η) is actually an eigenform for real multiplication with prototype P .

The equations for $\overline{W}_D$ and $\overline{P}_D$ are obvious from the definition of the coordinates. The leaves of $\mathcal{A}_D$ are as claimed because with our normalization the periods $\eta(\alpha_i)$ are constant, and either $y = \text{const}$ or $z = \text{const}$ implies that both of the $\eta(\beta_i)$ are constant by (7.12). $\blacksquare$

This also completes the proof of Theorem 7.17 because we explicitly constructed eigenforms near any point of $C_{\lambda(P)}$ for any nondegenerate Y_D -prototype P in Theorem 7.22 and showed that these are the only possible limits in Corollary 7.19.

Branches of $\overline{X}_D$ through C_λ . Let $p = (X, [\omega]) \in C_\lambda$, and let $q \in \mathbb{P}\Omega\mathcal{D}_2(S)$ (for the appropriate curve system S), with $\pi(q) = p$. We have seen that around q , the variety $\pi^{-1}(\overline{X}_D)$ is the union of the V_P over all fine Y_D -prototypes P such that $\lambda(P) = \lambda$.

To understand the local structure of $\overline{X}_D$ around p , we need only to understand how $\text{Aut}(X, [\omega])$ acts on this picture. If $\lambda > 1$, then $\text{Aut}(X, [\omega])$ consists only of the hyperelliptic involution which acts trivially on $\mathbb{P}\Omega\mathcal{D}_2(S)$. Thus π is a local isomorphism around q . Note also that there is no distinction between fine and ordinary prototypes when $\lambda > 1$.

When $\lambda = 1$, we have $\text{Aut}(X, [\omega]) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2$. It is generated by the hyperelliptic involution together with an involution t which interchanges the two nonseparating nodes of X and preserves ω . This involution t acts on $\mathbb{P}\Omega\mathcal{D}_2(S)$ by $t(x, y, z) = (x, z, y)$ in the coordinates around q defined above, so t identifies V_P with $V_{i(P)}$, where i is the involution of (3.1). Thus V_P and $V_{i(P)}$ have the same image in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$. We will abuse notation and denote the image of V_P in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$ by V_P as well, where now P is an ordinary Y_D -prototype. This makes sense because it defines the same object if P is represented by $i(P)$. We will think of this V_P as a union of branches of $\overline{X}_D$ through p . To summarize, whether or not $\lambda = 1$, the germ of $\overline{X}_D$ around P is the union of the V_P , where P ranges over all ordinary Y_D -prototypes such that $\lambda(P) = \lambda$.

If $p = (X, [\omega])$ has a double zero or a separating node, then in the same way – just replacing V_P with V'_P defined above – we define a branch V'_P of $\overline{X}_D$

through p , where now P is a W_D or P_D -prototype (respectively). This gives a finer classification of branches then we obtain for generic $p \in C_\lambda$

To summarize this discussion, we have the following Corollary of Theorem 7.22.

Corollary 7.23. *For any $p \in C_\lambda$, the germ of $\overline{X}_D$ through p is the union of germs,*

$$\bigcup_{\substack{P \in \mathcal{Y}_D \\ \lambda(P)=\lambda}} V_P.$$

Each germ V_P consists of $\text{mult}(P)$ branches of $\overline{X}_D$ through P .

If $p = w_\lambda$ or $p = p_\lambda$, then alternatively the germ of $\overline{X}_D$ through p is the union,

$$\bigcup_{\substack{P \in \mathcal{W}_D \\ \lambda(P)=\lambda}} V'_P,$$

or respectively,

$$\bigcup_{\substack{P \in \mathcal{P}_D \\ \lambda(P)=\lambda}} V'_P.$$

In these cases, each of these germs is actually an irreducible branch of $\overline{X}_D$ through p .

Proof. The only parts that don't follow directly from Theorem 7.22 and the discussion above are the statements that V_P consists of $\text{mult}(P)$ branches and that V'_P is irreducible. This will follow directly from Proposition 8.1 because

$$\gcd(a', c') = \frac{\gcd(a, c)}{\gcd(a, b, c)} = \text{mult}(P),$$

and $\gcd(a'', c'') = 1$. ■

A section of $\mathcal{O}(-1)$. We normalized each projective class $(Y, [\eta]) \in U_\lambda$ so that each $\eta(\alpha_1) = 1$. We can regard this as defining a section s of the canonical line bundle $\mathcal{O}(-1)$ over U_λ .

The following follows directly from (7.4) together with the definition of our coordinates.

Proposition 7.24. *The norm of this section s of $\mathcal{O}(-1)$ over U_λ is given by,*

$$h(s, s) = -\frac{1}{2\pi} \log |y| - \frac{\lambda^2}{2\pi} \log |z|. \quad (7.18)$$

7.7 The strata $X_D(T_{3,0})$ and $X_D(T_{3,0}, T_{3,0}^3)$

We now study Abelian differentials in $\overline{X}_D$ with three nonseparating nodes. These consist of the union of the stratum $X_D(T_{3,0})$, where all of the nodes are polar, and the stratum $X_D(T_{3,0}, T_{3,0}^3)$, where one of the nodes is holomorphic. We will prove the following classification of such differentials.

Theorem 7.25. *The stratum $X_D(T_{3,0})$ is the finite union of points,*

$$X_D(T_{3,0}) = \bigcup_P c_{\lambda(P)},$$

where the union is over all nonterminal Y_D prototypes P .

The stratum $X_D(T_{2,0}, T_{3,0}^3)$ is the finite union of points,

$$X_D(T_{3,0}, T_{3,0}^3) = \bigcup_P c_{\lambda(P)},$$

where the union is over all terminal Y_D -prototypes.

If P is neither degenerate nor terminal, then $c_{\lambda(P)}$ lies in the intersection of the closures of $C_{\lambda(P)}$ and $C_{\lambda(P+)}$. If P is a degenerate prototype, then c_P lies in the intersection of the closures of $X_D(T_{1,0})$ and $C_{\lambda(P+)}$. If P is a terminal prototype, then $c_{\lambda(P)}$ lies in the intersection of the closures of $C_{\lambda(P)}$ and $X_D(T_{1,0}, T_{1,0}^1)$. None of the other curves in ∂X_D pass through $c_{\lambda(P)}$.

Local coordinates. Fix an $(X, [\omega]) \in \mathbb{P}\Omega\mathcal{M}_2^0(T_{3,0})$ or $\mathbb{P}\Omega\mathcal{M}_2^0(T_{3,0}, T_{3,0}^3)$, and choose a marking $\Sigma_2 \rightarrow X$ so that we can regard $(X, [\omega])$ as lying in $\mathbb{P}\Omega\mathcal{T}_2^0(T_{3,0})$ or $\mathbb{P}\Omega\mathcal{T}_2^0(T_{3,0}, T_{3,0}^3)$.

Our first goal is to show that if $(X, [\omega]) \in \overline{X}_D$, then $(X, [\omega])$ is one of the points $c_{\lambda(P)}$ as in the statement of Theorem 7.25. If $(X, [\omega]) \in \overline{X}_D$, then ω is an eigenform for real multiplication of $\mathcal{O}_D$ on $\text{Jac}(X)$ by Theorem 5.10, so assume that it is such an eigenform. It follows that the ratios of the residues of any two poles of ω are real, so we can normalize the projective class $[\omega]$ so that all of these residues are real.

Suppose ω has no holomorphic node. In this case it also has two simple zeros; let $\{p_1, p_2\} \in \Sigma_2$ be the inverse image of the zeros. Let $\alpha_i \in H_1(\Sigma_2 \setminus \{p, q\})$ be homology classes representing the three curves $T_{3,0}^i$ such that $\omega(\alpha_i) \geq 0$, and choose $\gamma_i \in H_1(\Sigma_2, \{p_1, p_2\})$ so that $\alpha_i \cdot \gamma_j = \delta_{ij}$ as in §6.7. Then the classes,

$$\begin{aligned} \beta_1 &= \gamma_1 - \gamma_3 \\ \beta_2 &= \gamma_2 + \gamma_3, \end{aligned} \tag{7.19}$$

in $H_1(\Sigma_2; \mathbb{Z})$ are dual to α_1 and α_2 .

If ω does have a holomorphic node, then let $\alpha_i \in H_1(\Sigma_2)$ be homology classes representing the three curves $T_{3,0}^i$ such that $\omega(\alpha_i) \geq 0$, and let $\beta_i \in H_1(\Sigma_2; \mathbb{Z})$ be dual to α_1 and α_2 .

Choose a small neighborhood U of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{3,0})$ or $\mathbb{P}\Omega\mathcal{D}_2^0(T_{3,0}, T_{3,0}^3)$ such that the natural map $U/\text{Aut}(X, [\omega]) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$ is an isomorphism onto

its image and such that for each $(Y, [\eta]) \in U \cap \pi^{-1}(X_D)$, the subgroup $S = \langle \alpha_1, \alpha_2 \rangle$ of $H_1(\Sigma_2; \mathbb{Z})$ is preserved by the real multiplication, which is possible by Theorem 5.9.

For the rest of this section, we will adopt the convention that each projective class $(Y, [\eta]) \in U$ is normalized so that $\eta(\alpha_1) = 1$, so we can regard U as an open subset of a hypersurface in $\Omega\mathcal{D}_2^0(T_{3,0})$ or $\Omega\mathcal{D}_2^0(T_{3,0}, T_{3,0}^3)$.

Proposition 7.26. *If the neighborhood U of $(X, [\omega])$ is taken sufficiently small, then for each $(Y, \eta) \in U \cap \pi^{-1}(X_D)$, the $\mathcal{O}_D$ -submodule S of $H_1(\Sigma_2, \mathbb{Z})$ is a quasi-invertible $\mathcal{O}_D$ -module with $\{\alpha_1, \alpha_2, \alpha_3\}$ an admissible triple; furthermore, for $(Y, \eta) \in U$, the period maps P_η and $P_\omega: S \rightarrow \mathbb{R}$ are the same.*

Proof. Since the curves of $T_{3,0}$ bound a pair of pants, we get the relation

$$\pm \alpha_1 \pm \alpha_2 \pm \alpha_3. \quad (7.20)$$

To see that $\{\alpha_1, \alpha_2, \alpha_3\}$ is an admissible triple, it is enough to show that $\{\alpha_1, \alpha_2\}$ is an admissible basis. Just as in the proof of Proposition 7.18, it is enough to show that $\text{Im } \eta(\beta_i) > 0$ for (Y, η) sufficiently close to (X, ω) . This follows from Theorem 5.5, using the fact that the β_i pass through tall cylinders, and each cylinder that β_i passes through contributes positively to $\text{Im } \eta(\beta_i)$ because $\alpha_j \cdot \beta_i \geq 0$ for each j .

Since there are only finitely many admissible triples up to isomorphism by Proposition 3.6, the period $\eta(\alpha_2)$ must be constant on U if U is sufficiently small, and $\eta(\alpha_3)$ must be constant by (7.20). Thus the period maps P_ω and P_η are the same. $\blacksquare$

Now since $\{\alpha_1, \alpha_2, \alpha_3\}$ is an admissible triple, we can reorder them and replace ω by a constant multiple so that

$$\omega(\alpha_1) = 1, \omega(\alpha_2) = \lambda, \text{ and } \omega(\alpha_3) = \lambda - 1,$$

with $\lambda \geq 1$, and $\lambda = \lambda(P)$ for some Y_D -prototype P by Proposition 3.6. If ω has no holomorphic node, then there is a unique such choice of the α_i and ω (up to the hyperelliptic involution). Otherwise, there are two such choices because we can swap α_1 and α_2 . Also when $(X, [\omega])$ has no holomorphic nodes, reorder and possibly change the signs of the γ_i so that we still have $\alpha_i \cdot \gamma_j = \delta_{ij}$, and redefine the β_i as in (7.19) with the new γ_i . When $(X, [\omega])$ has a holomorphic node, reorder the β_i so that they are still dual to the α_i .

Thus we see that if $(X, [\omega])$ has three nonseparating nodes and $(X, [\omega]) \in \overline{X}_D$, then $(X, [\omega]) = c_{\lambda(P)}$ for some Y_D -prototype P as stated in Theorem 7.25. Also, Proposition 7.26 implies that if U is sufficiently small and if we define for $\mu \in \mathbb{R}$,

$$U_\mu = \{(Y, \eta) \in U : \eta(\alpha_1) = 1 \text{ and } \eta(\alpha_2) = \mu\},$$

then $U \cap \pi^{-1}(\overline{X}_D) \subset U_\lambda$, where $\lambda = \omega(\alpha_2)$, by Proposition 7.26.

Recall that in §6.7 we defined local coordinates (v, w, x, y, z) on $\Omega\mathcal{D}_2^0(T_{3,0})$. When $\lambda > 1$, in these local coordinates, we can identify U_λ with the subspace

defined by $v = 1$ and $w = \lambda$. Then on U_λ , the (x, y, z) become

$$\begin{aligned} x &= e^{2\pi i \eta(\gamma_1)} \\ y &= e^{2\pi i \eta(\gamma_2)/\lambda} \\ z &= e^{2\pi i \eta(\gamma_3)/(\lambda-1)}, \end{aligned}$$

and the (x, y, z) are local coordinates on U_λ .

In §6.8 we defined local coordinates (w, x, y, z) on $\Omega\mathcal{D}_2^0(T_{3,0}, T_{3,0}^3)$. In these local coordinates, we can identify U_1 with the subspace defined by $w = 1$. Then on U_1 , the (x, y, z) become

$$\begin{aligned} x &= e^{2\pi i \eta(\beta_1)} \\ y &= e^{2\pi i \eta(\beta_2)} \\ z &= \left(\int_I \eta \right)^2, \end{aligned}$$

and the (x, y, z) are local coordinates on U_1 .

Prototypes. Now assume that our $(X, [\omega]) = c_\lambda$, where $\lambda = \lambda(P)$ for some Y_D -prototype P with a small neighborhood U of $(X, [\omega])$ in the Dehn space chosen as above. We now assign a fine Y_D -prototype $P(Y, [\eta])$ to each $(Y, [\eta]) \in U \cap \pi^{-1}(X_D)$ as in §7.6.

Let $(Y, [\eta]) \in U \cap \pi^{-1}(X_D)$. We have a marking $\Sigma_2 \rightarrow Y$ which is defined up to Dehn twist around the curve system $T_{3,0}$. Choose a marking $\Sigma_2 \rightarrow Y$ and use this to transport the symplectic basis $\{\alpha_1, \alpha_2, \beta_1, \beta_2\}$ to Y .

There is some $\mu \in K_D$ with $\mu^{(1)} = \lambda$ such that $\mu \cdot \alpha_1 = \alpha_2$ and

$$a\mu^2 + b\mu + c = 0$$

for integers a, b , and c satisfying the properties of Proposition 3.6. The generator $a\mu$ of $\mathcal{O}_D$ defines by its action on $\text{Jac}(Y)$ an endomorphism T of $H_1(Y; \mathbb{Z})$ which is given by the matrix (7.9) for some integer q . Define the prototype $P(Y, \eta) = (a, b, c, \bar{q})$, where $\bar{q}$ is the reduction of q modulo $\gcd(a, b, c)$. By the same argument as in Theorem 7.21, $P(Y, [\eta])$ is a well-defined fine Y_D -prototype.

Local coordinates for $\overline{X}_D$. Given $(X, [\omega]) = c_\lambda$ and a small neighborhood U of $(X, [\omega])$ in the Dehn space as above, we now give equations for $\pi^{-1}(\overline{X}_D)$ in U and show that $(X, [\omega])$ is indeed in $\overline{X}_D$. For each fine Y_D -prototype P , let $\tilde{V}_P \subset U$ be the closure of

$$\{(Y, \eta) \in U \cap \pi^{-1}(X_D) : P(Y, \eta) = P\},$$

which actually lies in U_λ . The following theorem gives explicit equations for $\tilde{V}_P$ in the coordinates (x, y, z) on U_λ .

Theorem 7.27. *The locus $\tilde{V}_P \subset U_{\lambda(P)}$ is cut out by the equation,*

$$x^{a'} = e^{-2\pi i q'} y^{-c'} z^{-a'-b'-c'}. \quad (7.21)$$

If P is a nondegenerate and nonterminal Y_D -prototype, then the closure $\overline{C}_{\lambda(P)}$ is cut out by the equations $x = y = 0$, and $\overline{C}_{\lambda(P^+)}$ is cut out by the equations $x = z = 0$.

If P is a degenerate Y_D -prototype, then the additional equation $y = 0$ cuts out $U \cap X_D(T_{1,0})$, and the equations $x = z = 0$ cuts out $\overline{C}_{\lambda(P^+)}$.

If P is a terminal Y_D -prototype, then the additional equation $z = 0$ cuts out $U \cap X_D(T_{1,0}, T_{1,0}^1)$, and the equation $x = y = 0$ cuts out $\overline{C}_{\lambda(P)}$.

The leaves of the foliation $\mathcal{A}_D$ of X_D are given in these local coordinates by $y^\lambda z^{\lambda-1} = \text{const}$.

Proof. If the prototype P is terminal, then (7.21) reduces to (7.12), and the proof that $\tilde{V}_P$ is given by this equation is the same as in Theorem 7.22.

Now assume that P is not terminal. Using the same symplectic basis $\{\alpha_i, \beta_i\}$ and $\mu \in K_D$ such that $\mu \cdot \alpha_1 = \alpha_2$ as above, equation (7.9) again implies (7.14) and (7.15). Integrating η over both sides yields (7.16) and (7.17), and substituting in (7.19) gives

$$a\eta(\gamma_1) + \frac{c}{\lambda}\eta(\gamma_2) + \frac{a+b+c}{\lambda-1}\eta(\gamma_3) = -q \quad (7.22)$$

Dividing by $\gcd(a, b, c)$ and exponentiating yields (7.21).

Conversely if (Y, η) is in U_λ minus the coordinate axes and is in the locus defined by (7.21), then (7.22) holds modulo $\gcd(a, b, c)$, and by changing q modulo $\gcd(a, b, c)$, we can ensure that it holds exactly. It then follows that (7.16) and (7.17) hold, and so (Y, η) is an eigenform.

The equations for the C_λ and the other strata are straightforward.

The periods $\eta(\alpha_1)$ and $\eta(\alpha_2)$ are fixed because of our normalization for η , and the periods $\eta(\beta_1)$ and $\eta(\beta_2)$ are related by the real multiplication, so if one is constant, then the other is. Thus the leaves of the foliation $\mathcal{A}_D$ in these coordinates are given by $\eta(\beta_2) = \text{const}$, which is equivalent to $y^\lambda z^{\lambda-1} = \text{const}$. ■

Proof of Theorem 7.25. We saw as a consequence of Proposition 7.26 that each stable Abelian differential in $\overline{X}_D$ with three nonseparating nodes is one of the $c_{\lambda(P)}$ for some Y_D -prototype P , and it follows from Theorem 7.27 that these $c_{\lambda(P)}$ are in fact in $\overline{X}_D$. The other statements also follow directly from Theorem 7.27. ■

Branches of $\overline{X}_D$ through c_λ . The varieties cut out by (7.21) have a single branch through the origin, as we will see in Proposition 8.3, so we obtain a bijective correspondence between branches of $\pi^{-1}(\overline{X}_D)$ through $\pi^{-1}(c_\lambda)$ and fine Y_D -prototypes P such that $\lambda(P) = \lambda$ given by $P \mapsto \tilde{V}_P$. Just as in §7.6, the action of $\text{Aut}(X, [\omega])$ identifies branches corresponding to the same ordinary Y_D -prototype, so we obtain the first statement of the following theorem.

Theorem 7.28. *The germ of $\overline{X}_D$ through c_λ is the following union of irreducible branches:*

$$\bigcup_{\substack{P \in \mathcal{Y}_D \\ \lambda(P) = \lambda}} \tilde{V}_P.$$

If P is a nondegenerate Y_D -prototype such that $\lambda(P) = \lambda$, then the branch $\tilde{V}_P$ through c_λ intersects C_λ in the germ V_P of Corollary 7.23. If P is a nonterminal Y_D -prototype such that $\lambda(P) = \lambda$, then the branch $\tilde{V}_P$ through c_λ intersects $C_{\lambda(P+)}$ in the germ V_{P+} .

Proof. Suppose that $P = (a, b, c, \bar{q})$ is nonterminal. Let $(Y, \eta) \in \tilde{V}_P$ be a non-singular Abelian differential close enough to a point $p \in C_{\lambda(P+)}$ that it is contained in some well-defined subgerm $V_{P'}$ of $\overline{X}_D$ through p . We need to show that $P' = P^+$.

We will continue to use the notation that we used to define the local coordinates around c_λ , so we have classes α_1, α_2 , and $\alpha_3 \in H_1(Y; \mathbb{Z})$ representing the nodes of c_λ such that $\eta(\alpha_1) = 1$, $\eta(\alpha_2) = \lambda$, and $\eta(\alpha_3) = \lambda - 1$; we have β_1 , and $\beta_2 \in H_1(Y; \mathbb{Z})$ such that $\{\alpha_1, \alpha_2, \beta_1, \beta_2\}$ is a symplectic basis of $H_1(Y; \mathbb{Z})$; and we have a $\mu \in K_D$ such that $\mu \cdot \alpha_1 = \alpha_2$ and $a\mu^2 + b\mu + c = 0$.

In order to calculate the prototype P' , we must choose a symplectic basis $\{\alpha'_1, \alpha'_2, \beta'_1, \beta'_2\}$ of $H_1(Y; \mathbb{Z})$ such that the α'_i are homologous to cylinders on Y representing the nodes of $p \in C_{\lambda(P+)}$ and such that if we set

$$\tilde{\lambda} = \frac{\eta(\alpha'_2)}{\eta(\alpha'_1)},$$

then $\tilde{\lambda} \geq 1$ and $\tilde{\lambda}' < 0$. Given such a basis, let $\tilde{\mu} \in K_D$ be such that $\tilde{\mu} \cdot \alpha'_1 = \alpha'_2$. Let a', b' , and c' be the unique integers such that

$$a'\tilde{\mu}^2 + b'\tilde{\mu} + c' = 0,$$

$a' > 0$, and $(b')^2 - 4a'c' = D$. For $x \in K_D$, set T_x and T'_x to be the matrices of the action of x on $H_1(Y; \mathbb{Z})$ in the bases $\{\alpha_i, \beta_i\}$ and $\{\alpha'_i, \beta'_i\}$ respectively. Then the prototype P' is $P' = (a', b', c', \bar{q}')$, where q' is the entry in the upper right corner of $T'_{a'\tilde{\mu}}$.

There are two cases to consider, depending on whether or not $\lambda \geq 2$. First suppose that $\lambda \geq 2$, or equivalently $4a + 2b + c \leq 0$. Define the new symplectic basis of $H_1(Y; \mathbb{Z})$,

$$\begin{aligned} \alpha'_1 &= \alpha_1 & \beta'_1 &= \beta_1 + \beta_2 \\ \alpha'_2 &= \alpha_3 = \alpha_2 - \alpha_1 & \beta'_2 &= \beta_2. \end{aligned}$$

Then $\tilde{\mu} = \mu - 1$, and $\tilde{\lambda} = \lambda - 1$, which satisfies the required properties. We have

$$(a', b', c') = (a, 2a + b, a + b + c).$$

The matrix $T_{a\mu}$ is given by (7.9), and an easy calculation shows that the new matrix is

$$T'_{a'\tilde{\mu}} = \begin{pmatrix} 0 & -a-b-c & 0 & q \\ a & -2a-b & -q & 0 \\ 0 & 0 & 0 & a \\ 0 & 0 & -a-b-c & -2a-b \end{pmatrix},$$

so $P' = P^+$ is as claimed.

Now consider the case where $\lambda < 2$, or equivalently $4a + 2b + c > 0$. In this case, define a new symplectic basis of $H_1(Y; \mathbb{Z})$ by,

$$\begin{aligned} \alpha'_1 &= \alpha_2 - \alpha_1 & \beta'_1 &= \beta_2 \\ \alpha'_2 &= \alpha_1 & \beta'_2 &= \beta_1 + \beta_2. \end{aligned}$$

Then

$$\tilde{\mu} = \frac{1}{\mu - 1} = \frac{-a\mu - a - b}{a + b + c},$$

and $\tilde{\lambda} = 1/(\lambda - 1)$ satisfies the required properties that $\tilde{\lambda} \geq 1$ and $\tilde{\lambda}' < 0$, using the fact that $\lambda > 1$ because P is nonterminal. We have

$$(a', b', c') = (-a - b - c, -2a - b, -a),$$

and an easy calculation shows that the new matrix is,

$$T'_{a'\tilde{\mu}} = T'_{a\mu+a+b} = \begin{pmatrix} 0 & a & 0 & q \\ -a-b-c & 2a+b & -q & 0 \\ 0 & 0 & 0 & -a-b-c \\ 0 & 0 & a & 2a+b \end{pmatrix}.$$

Thus $P' = P^+$ as claimed.

Calculating the prototype of the intersection of $\tilde{V}_P$ with C_λ is much easier. Here we can use the old basis $\{\alpha_i, \beta_i\}$ of $H_1(Y; \mathbb{Z})$ to calculate the prototype, and thus it is just P . $\blacksquare$

A section of $\mathcal{O}(-1)$. We normalized each projective class $(Y, \eta) \in U_\lambda$ so that each $\eta(\alpha_1) = 1$. We can regard this as defining a section s of the canonical line bundle $\mathcal{O}(-1)$ over U_λ .

The following follows directly from (7.4) together with the definition of our coordinates.

Proposition 7.29. *The norm of this section s of $\mathcal{O}(-1)$ over U_λ is given by,*

$$h(s, s) = -\frac{1}{2\pi} \log |x| - \frac{\lambda^2}{2\pi} \log |y| - \frac{(\lambda - 1)^2}{2\pi} \log |z|. \quad (7.23)$$

8 Geometric compactification of X_D

We have compactified the Hilbert modular surface X_D by taking its closure $\overline{X}_D$ in $\mathbb{P}\Omega\overline{\mathcal{M}}_2$; however, this compactification is not suitable for our purposes. One problem is that $\overline{X}_D$ has non-normal singularities, and another is that several cusps of the curves W_D and P_D can come together to one point in $\overline{X}_D$. It turns out that taking the normalization of Y_D solves both of these problems and produces a useful compactification which has only quotient singularities.

Definition. The *geometric compactification* Y_D of the Hilbert modular surface X_D is the normalization of $\overline{X}_D$.

This defines Y_D as a variety. We will also give Y_D the structure of a complex orbifold and of a projective variety in §8.1.

In this section, we will study in detail the geometry of Y_D . In §8.1, we will study curves in $Y_D \setminus X_D$, and their intersections with the curves $\overline{W}_D$ and $\overline{P}_D$. We show in §8.2 that Y_D maps to the Baily-Borel compactification $\widehat{X}_D$ by a map which is the identity on X_D , and we draw conclusions about the cohomology of Y_D . In §8.3, we study the extension of involution τ of X_D to Y_D . We will assume that the reader is familiar with the operation of normalization. For a summary of the results about normalization which we use in this section, see §A.

8.1 Geometry of Y_D

The goal of this subsection is to classify the curves of $Y_D \setminus X_D$ and to understand the ways in which these curves intersect each other and the curves $\overline{W}_D, \overline{P}_D \subset Y_D$. This is basically a straightforward translation of the results of §7 on the local structure of $\overline{X}_D$ to the setting of the normalization.

Local normalization of $\overline{X}_D$. Normalization is a local operation: constructing the normalization of $\overline{X}_D$ just amounts to replacing each small open set with its normalization and then gluing together these normalizations. We saw in §7 that the singular points of $\overline{X}_D$ are locally modeled on the varieties $V(X^p - Y^q)$ and $V(X^p - Y^q Z^r)$ in $\mathbb{C}^3$, so we need to understand the normalizations of these varieties.

Let $\theta_m = e^{2\pi i/m}$. The proof of the following proposition is easy and will be left to the reader.

Proposition 8.1. *The normalization of the variety $V = V(X^p - Y^q) \subset \mathbb{C}^3$ is $\gcd(p, q)$ copies of $\mathbb{C}^2$ which map to V by*

$$f_r(x, y) = (x^q, \theta_{\gcd(p, q)}^r x^p, y).$$

It follows that V has $\gcd(p, q)$ branches through each singular point $(0, 0, Z)$.

Given p, q , and $r \in \mathbb{N}$ relatively prime, let

$$\begin{aligned} m &= \frac{p}{\gcd(p, q) \gcd(p, r)} \\ s &\equiv \left(\frac{r}{\gcd(p, r)} \right)^{-1} \pmod{\frac{p}{\gcd(p, r)}} \\ n &= \frac{-q}{\gcd(p, q)} s, \end{aligned} \tag{8.1}$$

and define $Q(p, q, r)$ to be the normal analytic space $\mathbb{C}^2/G$, where G is the order m cyclic group generated by the transformation,

$$(x, y) \mapsto (\theta_m x, \theta_m^n y).$$

Lemma 8.2. *If p, q and r are relatively prime integers, then the map*

$$f: \mathbb{Z}/\left(\frac{p}{\gcd(p, q)}\right) \times \mathbb{Z}/\left(\frac{p}{\gcd(p, r)}\right) \rightarrow \mathbb{Z}/p$$

defined by $f(\alpha, \beta) = q\alpha + r\beta$ is surjective, and its kernel is a cyclic group of order m which is generated by $(\gcd(p, r), -qs)$, where m and s are as defined in (8.1).

Proof. It follows from the fact that p, q , and r are relatively prime that f is surjective, so the kernel K has order m . The element $(\gcd(p, r), -qs)$ clearly belongs to K and has order at least m because $\gcd(p, r)$ has order m in

$$\mathbb{Z}/\left(\frac{p}{\gcd(p, q)}\right).$$

Thus it is a generator of K . ■

Proposition 8.3. *The normalization of the variety $V = V(X^p - Y^q Z^r) \subset \mathbb{C}^3$ is $p: Q(p, q, r) \rightarrow V$, where p is induced by the map $\tilde{p}: \mathbb{C}^2 \rightarrow V$ defined by*

$$\tilde{p}(x, y) = \begin{pmatrix} x^{q/\gcd(p, q)} y^{r/\gcd(q, r)} \\ x^{p/\gcd(p, q)} \\ y^{p/\gcd(p, r)} \end{pmatrix}.$$

Proof. For $i = 1$ or 2 , let (x_i, y_i) be a point in $\mathbb{C}^2$ minus its coordinate axes. We claim that

$$\tilde{p}(x_1, y_1) = \tilde{p}(x_2, y_2),$$

if and only if the (x_i, y_i) are related by the action of G . It would then follow that p is the normalization of V because p would be well-defined, finite-to-one, and biholomorphic on the complement of the coordinate axes.

It is easy to show that $\tilde{p}(x_1, y_1) = \tilde{p}(x_2, y_2)$ if the (x_i, y_i) are related by G . Conversely, suppose the (x_i, y_i) have the same image. Then we must have

$$(x_2, y_2) = \left(e^{2\pi i \alpha \gcd(p, q)/p} x_1, e^{2\pi i \beta \gcd(p, r)/p} x_2 \right)$$

for some pair,

$$(\alpha, \beta) \in \mathbb{Z}/\left(\frac{p}{\gcd(p, q)}\right) \times \mathbb{Z}/\left(\frac{p}{\gcd(p, r)}\right),$$

such that

$$q\alpha + r\beta \equiv 0 \pmod{p}.$$

By Lemma 8.2 there is some $k \in \mathbb{Z}$ such that

$$\begin{aligned} \alpha &\equiv k \gcd(p, r) \pmod{\frac{p}{\gcd(p, q)}} \\ \beta &\equiv -kqs \pmod{\frac{p}{\gcd(p, r)}}, \end{aligned}$$

and it follows that the (x_i, y_i) are related by an element of G . ■

Since the normalizations of Propositions 8.1 and 8.3 are either nonsingular or quotient singularities, we can regard Y_D as a complex orbifold.

One nonseparating polar node. We now begin to study the curves which make up $Y_D \setminus X_D$. Define $S_D^1 \subset Y_D$ by

$$S_D^1 = \pi^{-1}(X_D(T_{1,0})).$$

By Proposition 7.1, S_D^1 is empty unless D is square. By Corollary 7.7, points of $S_{d^2}^1$ correspond precisely to degree d cylinder covering differentials; $S_{d^2}^1$ is a suborbifold of Y_{d^2} by Theorem 7.4, and

$$S_{d^2}^1 \cong \mathbb{H}/\Gamma_1(d).$$

If $d > 3$, then $S_{d^2}^1$ is actually a submanifold of Y_{d^2} because $\Gamma_1(d)$ is torsion-free.

There are finitely many intersection points of $\overline{W}_{d^2}$ with $S_{d^2}^1$, and these intersections are transverse if $d > 3$ by Theorem 7.4. These intersection points are exactly the one-cylinder cusps of W_{d^2} .

One nonseparating holomorphic node. Define $S_D^2 \subset Y_D$ by

$$S_D^2 = \pi^{-1}(X_D(T_{1,0}, T_{1,0}^1)).$$

By Proposition 7.1, S_D^2 is empty unless D is square. $S_{d^2}^2$ is a suborbifold of Y_{d^2} isomorphic to $\mathbb{H}/\Gamma_1(d)$ by Theorem 7.9, and $S_{d^2}^2$ is a submanifold when $d > 3$. The points of $S_{d^2}^2$ correspond to genus one differentials with two marked points which differ by exactly d -torsion identified to form a node.

$\mathrm{SL}_2\mathbb{R}$ acts on $S_{d^2}^2$ with discrete stabilizer, so we can regard $S_{d^2}^2$ as a degenerate Teichmüller curve. Just as for W_{d^2} , cusps of $S_{d^2}^2$ correspond to periodic directions on a given $(X, \omega) \in S_{d^2}^2$, and we can divide the cusps of $S_{d^2}^2$ into one and two-cylinder cusps just as for W_{d^2} .

Given $q \in \mathbb{N}$ such that $0 < q \leq d/2$ and $\gcd(q, d) = 1$, let $s_{q/d} \in Y_{d^2}$ be the inverse image of $f_{q/d} \in \overline{X}_{d^2}$. By Theorem 7.13, these are transverse intersection points of $\overline{S}_{d^2}^1$ and $\overline{S}_{d^2}^2$, and these are all of their intersection points, because no other points in our classification of points in $\overline{X}_{d^2}$ lie in this intersection. These points $s_{q/d}$ are also exactly the one-cylinder cusps of $S_{d^2}^2$.

Two or more nodes. For each curve $C_\lambda \subset \overline{X}_D$, we associated a nondegenerate Y_D -prototype P to each branch of $\overline{X}_D$ through each point $p \in C_\lambda$. This means that we can associate a nondegenerate Y_D -prototype to each point in $\pi^{-1}(p)$ by Theorem A.8.

For any nondegenerate Y_D -prototype P , let $C_P^0 \subset Y_D$ be set of points in $\pi^{-1}(C_{\lambda(P)})$ associated to P . The projection $C_P^0 \rightarrow C_{\lambda(P)}$ is surjective and $\text{mult}(P)$ -to-one by Corollary 7.23. The union of the curves C_P parameterizes those points in Y_D representing differentials with two nonseparating nodes and possibly a separating node.

If $\lambda > 1$, then C_λ contains exactly one point $w_\lambda \in \overline{W}_D$ which represents an Abelian differential with a double zero by Proposition 6.10. We associated a W_D -prototype to each branch of $\overline{X}_D$ through w_λ , and we saw in Corollary 7.23 that W_D -prototypes P such that $\lambda(P) = \lambda$ correspond bijectively to such branches. Given any W_D -prototype P , let $w_P \in Y_D$ be the point in $\pi^{-1}(w_{\lambda(P)})$ corresponding to the branch associated to P . The point w_P is a two-cylinder cusp of W_D , and these are all of the two-cylinder cusps. This correspondence between W_D -prototypes and two-cylinder cusps of W_D is equivalent to the one described in [McM05a]; more precisely, the point w_P described here is the cusp w_P of Theorem 4.9.

Similarly, for any P_D -prototype P , define $p_P \in Y_D$ to be the unique point in $\pi^{-1}(p_{\lambda(P)})$ coming from the branch associated to P . These points p_P are all of the cusps of P_D .

We know that W_D and P_D are disjoint since they are $\text{SL}_2\mathbb{R}$ orbits. Their intersections with $Y_D \setminus X_D$ are also disjoint because we have accounted for all of these intersection points above, so it follows that $\overline{P}_D \cap \overline{W}_D = \emptyset$ in Y_D .

Suppose P is a W_D or a P_D -prototype that maps to the Y_D -prototype Q . Then the point w_P or p_P is contained in C_Q^0 . This means that if Q is a nonterminal prototype, then C_Q contains $\text{mult}(P)$ points of $\overline{W}_D$ and the same number of points of $\overline{P}_D$. If Q is a terminal prototype, then C_Q contains a single point of $\overline{P}_D$ (because $\text{mult}(Q) = 1$) and no point of $\overline{W}_D$ because no W_D -prototype maps to a terminal Y_D -prototype.

Proposition 8.4. *The locus C_P^0 is a nonsingular curve in Y_D . The restriction $\pi: C_P^0 \rightarrow C_{\lambda(P)}$ is an unramified cover of degree $\text{mult}(P)$, and C_P^0 is a twice-punctured sphere.*

Furthermore, $\overline{W}_D$ and $\overline{P}_D$ are smooth suborbifolds of Y_D which intersect C_P^0 transversely.

Proof. Let $p \in C_P^0$, and $q = \pi(p)$. By Theorem 7.22, the branch V of $\overline{X}_D$ through q corresponding to p is isomorphic to the germ at the origin of the

variety in $\mathbb{C}^3$ cut out by

$$x^{a''} = \theta y^{-c''}$$

for some root of unity θ . In these coordinates, $C_{\lambda(P)}^0$ is cut out by $x = y = 0$. If $q = w_{\lambda(P)}$ or $q = P_{\lambda(P)}$, then $\overline{W}_D$ or $\overline{P}_D$ is cut out by $z = 0$.

In coordinates (u, v) on a neighborhood of p , the map $\pi: Y_D \rightarrow X_D$ is given by

$$\pi(u, v) = (u^{-c''}, \theta' u^{a''}, v),$$

for some root of unity θ' . In these coordinates, C_P is cut out by $u = 0$, and if $p \in \overline{W}_D$ or $\overline{P}_D$, then $\overline{W}_D$ or $\overline{P}_D$ is cut out by $v = 0$. The smoothness and transversality statements follow immediately.

Since every point in $C_{\lambda(P)}^0$ has $\text{mult}(P)$ preimages in C_P , the projection is unramified of that degree. Thus C_P^0 is a twice-punctured sphere because it is an unramified cover of one.

Finally, $\overline{W}_D$ and $\overline{P}_D$ are suborbifolds because the coordinates above show that $\overline{W}_D$ and $\overline{P}_D$ are nonsingular where they cross the C_P^0 . These are the only points where $\overline{P}_D$ and $\overline{W}_D$ meet $Y_D \setminus X_D$ except for the intersections of $\overline{W}_D$ with S_D^1 in the case when D is square, and we have already seen that $\overline{W}_D$ is nonsingular there in Theorem 7.4. $\blacksquare$

Remark. The curves C_P^0 contain no orbifold points of Y_D . This is clear when $\lambda(P) > 1$ because the image $C_{\lambda(P)}^0$ of C_P^0 in $\mathbb{P}\Omega\mathcal{M}_2$ contains no orbifold points. When $\lambda(P) = 1$, the image C_1 is an orbifold locus of order two, so *a priori* C_P could be also. This doesn't happen because for any $(X, [\omega]) \in C_1$, the group $\text{Aut}(X, [\omega])$ fixes none of the branches of the inverse image of $\overline{X}_D$ through the inverse image of $(X, [\omega])$ in $\mathbb{P}\Omega\mathcal{D}_2(S)$.

We now study the closure of C_P^0 , which we call C_P . By Proposition 6.9, $C_P \setminus C_P^0$ consists of points representing differentials c_λ with three nonseparating nodes, defined in §6.6.

By Theorems 7.25 and 7.28, for every Y_D -prototype P , there is a point $c_{\lambda(P)} \in \overline{X}_D$ and a branch of $\overline{X}_D$ through $c_{\lambda(P)}$ labeled by P . Let $c_P \in Y_D$ be the point corresponding to the branch labeled with P . This gives a bijective correspondence between the points of Y_D representing Abelian differentials with three nonseparating nodes and Y_D -prototypes. Unless P is terminal, all three nodes of c_P are polar.

Recall that for a prototype $P = (a, b, c, \bar{q})$, we set

$$(a', b', c') = (a, b, c) / \gcd(a, b, c).$$

The following proposition describes the structure of Y_D in a neighborhood of c_P .

Proposition 8.5. *A neighborhood of the point c_P , where $P = (a, b, c, \bar{q})$ is a nondegenerate, nonterminal Y_D -prototype, is isomorphic to a neighborhood U/G*

of the origin in the quotient singularity $Q(a', -c', -a' - b' - c')$. In coordinates (u, v) on U , we have

$$\begin{aligned} V(uv) &= (Y_D \setminus X_D) \cap U \\ V(u) &= C_P \cap U \\ V(v) &= C_{P^+} \cap U. \end{aligned}$$

If P is a degenerate prototype, then c_P is a smooth point of Y_D . In a neighborhood of c_P , the complement $Y_D \setminus X_D$ is contained in $\overline{S}_D^1 \cup C_{P^+}$, and these curves meet transversely at c_P .

If P is a terminal prototype, then c_P is also a smooth point of Y_D . In a neighborhood of c_P , the complement $Y_D \setminus X_D$ is contained in $\overline{S}_D^2 \cup C_P$, and these curves meet transversely at c_P .

Proof. These statements all follow directly from Theorems 7.27 and 7.28, using Proposition 8.3 to translate these theorems into local coordinates on Y_D . ■

It follows from this proposition that the order of the orbifold point c_P is

$$m_P = \frac{a'}{\gcd(a', c') \gcd(a', b' + c')} = \frac{a}{\gcd(a, c) \gcd(a, b + c) \gcd(a, b, c)}.$$

Proposition 8.6. *The curve C_P is a connected rational curve which meets the points c_P and c_{P^-} and no other point c_Q . If C_P is given the structure of an orbifold with orbifold points of order m_P at c_P and m_{P^-} at c_{P^-} , then C_P is a suborbifold of Y_D .*

Proof. The complement $C_P \setminus C_P^0$ is contained in the union of the points c_Q , and by Proposition 8.5 it consists of exactly the points c_P and c_{P^-} . It follows from the coordinates in Proposition 8.5 that C_P is a suborbifold at these two points.

The curve C_P is rational because it is the closure of a twice-punctured sphere. ■

When D is not square, the operations P^+ and P^- are defined for all P , so the curves C_P are divided into finitely many closed chains of rational curves. When $D = d^2$, the curves C_P are divided into finitely many chains of rational curves joining $\overline{S}_{d^2}^1$ to $\overline{S}_{d^2}^2$.

To summarize, we have established the following:

Theorem 8.7. *Y_D has the following properties:*

1. Y_D is a compact, complex orbifold. Its orbifold points are located at the elliptic points of X_D , the elliptic points of $S_{d^2}^i \cong \mathbb{H}/\Gamma_1(d)$, and the points c_P for which $m_P > 1$.
2. The curves C_P , $\overline{W}_D$, $\overline{P}_D$, and $\overline{S}_D^i$ are all suborbifolds of Y_D .
3. The curves $\overline{S}_D^2$, $\overline{W}_D$, and $\overline{P}_D$ are pairwise disjoint, and $\overline{S}_D^1$ is disjoint from $\overline{P}_D$.

4. The curves $\overline{S}_D^1, \overline{S}_D^2, \overline{W}_D$, and $\overline{P}_D$ meet the curves C_P transversely. These intersections correspond to two cylinder cusps of S_D^2 and W_D . If $D = d^2$ with $d > 3$, then $\overline{W}_{d^2}$ intersects $S_{d^2}^1$ transversely.
5. The point $w_P \in \overline{W}_D$ lies on the curve C_Q , where Q is the Y_D -prototype associated to P , and w_P lies on no other curve $C_{Q'}$. This gives a bijective correspondence between the intersection points of $\overline{W}_D$ with C_Q and the set of W_D -prototypes associated to the Y_D -prototype Q . Similarly, the intersection points of $\overline{P}_D$ with C_P correspond bijectively in the same way to the set of P_D -prototypes associated to the Y_D -prototype P .

If P is neither initial nor terminal, then C_P meets W_D and P_D in $\text{mult}(P)$ points.

If P is terminal, then C_P meets $\overline{S}_{d^2}^2$ and $\overline{P}_D$ once each and is disjoint from $\overline{W}_D$.

If P is initial, then C_P meets $\overline{S}_{d^2}^1, \overline{P}_D$, and $\overline{W}_D$ once each.

6. The curves $\overline{S}_{d^2}^1$ and $\overline{S}_{d^2}^2$ meet transversely in the N points $s_{q/d}$, where

$$N = \begin{cases} 1, & \text{if } d = 2; \\ \frac{1}{2}\phi(d), & \text{if } d > 2 \end{cases}$$

(here ϕ is the Euler phi-function).

Corollary 8.8. *If D is not square, then W_D and P_D have the same number of cusps. The number of cusps of P_{d^2} is equal to the number of two-cylinder cusps of W_{d^2} plus the number of two-cylinder cusps of $S_{d^2}^2$.*

Example: Y_{17} . We will now illustrate these results in some specific cases. Most of the prototypes which arise will have $\bar{q} = 0$, and we will abbreviate those prototypes by omitting q .

There are five Y_{17} prototypes:

$$(1, 1, -4), \quad (2, -3, -1), \quad (2, -1, -2), \quad (1, -3, -2), \quad (1, -1, -4).$$

They form one orbit under the operation $P \mapsto P^+$, with each prototype being sent to the next on the list. We have $\text{mult}(2, -1, -2) = 2$, so $C_{(2, -1, -2)}$ meets $\overline{W}_{17}$ in the points $w_{(2, -1, -2, 0)}$ and $w_{(2, -1, 2, 1)}$ and $\overline{P}_{17}$ in the points $p_{(2, -1, -2, 0)}$ and $p_{(2, -1, 2, 1)}$. The other prototypes have multiplicity one, so the other curves C_P meet $\overline{W}_{17}$ and $\overline{P}_{17}$ once each. Since $m_P = 1$ for each P , none of the points c_P are orbifold points, and the C_P meet each other transversely.

This configuration is shown in Figure 13. The curves C_P form the pentagon and are marked by their prototypes. The curves representing $\overline{W}_{17}$ and $\overline{P}_{17}$ in this figure accurately represent the connected components of these curves and their intersections with the C_P .




Figure 13: Y_{17}

Example: Y_{12} . There are three Y_{12} prototypes:

$$(2, -2, -1), \quad (1, -2, -2), \quad (1, 0, -3).$$

They form one orbit under the operation $P \mapsto P^+$. We have $m_{(2,-2,-1)} = 2$, so the point $c_{(2,-2,-1)}$ is an orbifold point of order two in the intersection of $C_{(1,-2,-2)}$ and $C_{(2,-2,-1)}$. The other two points c_P are nonsingular. Each of these prototypes has multiplicity one, so W_{12} and P_{12} each have three cusps and intersect each C_P once.

Example: Y_{25} . There are nine Y_{25} -prototypes, which are divided into two orbits under $P \mapsto P^+$:

$$(1, -5, 0), \quad (1, -3, -4), \quad (1, -1, -6), \quad (1, 1, -6), \quad (1, 3, -4);$$

and

$$(2, -5, 0), \quad (2, -1, -3), \quad (2, -3, -2), \quad (2, 1, -3).$$

The prototypes after the first one on each list are nondegenerate and correspond to seven curves C_P in Y_{25} . Each of these prototypes has $m_P = 1$, so none of the c_P are orbifold points. The curve W_{25} has six two-cylinder cusps; it meets $C_{(2,-3,-2)}$ twice and the other four curves C_P corresponding to nonterminal prototypes once each. W_{25} also has two one-cylinder cusps (see §13, [McM05a], or [LR06] for formulas for the number of one-cylinder cusps) corresponding to two intersections with S_{25}^1 . The curve $\overline{P}_{25}$ also meets $C_{(2,-3,-2)}$ twice and each of the other C_P once each.

Each of the two-cylinder cusps of W_{25} can be represented by a square-tiled surface with five squares having periodic horizontal direction. Figure 14 depicts the curves C_P and S_{25}^i in Y_{25} . Next to each C_P are diagrams representing square tiled surfaces associated to each cusp of W_{25} which meets C_P .

Projective structure of Y_D . So far, we have given Y_D the structure of an algebraic variety and a complex orbifold. We can also give Y_D the structure of a projective variety:

Proposition 8.9. *Y_D is a projective variety.*

Proof. Let $Z \subset \overline{\mathcal{M}}_2$ be the closure of the image of the natural map $X_D \rightarrow \overline{\mathcal{M}}_2$. The image of $\pi: Y_D \rightarrow \overline{\mathcal{M}}_2$ is contained in Z , so we have a finite surjective morphism $Y_D \rightarrow Z$. This means that Y_D is the normalization of Z in $K(Y_D) = K(X_D)$. By Theorem A.3, Y_D is a projective variety because $\overline{\mathcal{M}}_2$ is projective. ■

8.2 Y_D maps to $\widehat{X}_D$

Recall that we introduced the Baily-Borel compactification $\widehat{X}_D$ of X_D in §2. If D is not square, $\widehat{X}_D$ consists of X_D together with a finite set C of cusps. If $D = d^2$, then $\widehat{X}_{d^2}$ consists of X_D together with the curves $R_{d^2}^1$ and $R_{d^2}^2$




Figure 14: Y_{25}

isomorphic to $\mathbb{H}/\Gamma_1(d)$ and the finite set C of cusps. Let $i: X_D \rightarrow Y_D$ be the natural inclusion.

Theorem 8.10. *There is a unique morphism $p: Y_D \rightarrow \widehat{X}_D$ such that $p \circ i = \text{id}_{X_D}$. This map p has the following properties:*

- p maps each curve C_P onto a cusp of $\widehat{X}_D$, and this induces a bijection between the connected components of $\bigcup_P C_P$ and the set of cusps $C \subset \widehat{X}_D$.

- The restriction,

$$p: Y_D \setminus \bigcup_P C_P \rightarrow X_D \setminus C$$

is an isomorphism.

- p maps $S_{d^2}^i$ isomorphically onto $R_{d^2}^i$.

Proof. Recall from §2 that there is the natural morphism $j: \widehat{X}_D \rightarrow \widehat{\mathcal{A}}_2$ with image Z such that $j: \widehat{X}_D \rightarrow Z$ is the normalization of Z in $K(X_D)$.

The varieties Y_D and $\widehat{X}_D$ both contain X_D as an open dense set, so we can identify both of their function fields with $K(X_D)$. The morphism $q: Y_D \rightarrow \widehat{\mathcal{A}}_2$ which is the composition of the maps

$$Y_D \rightarrow \overline{\mathcal{M}}_2 \rightarrow \widehat{\mathcal{A}}_2$$

has image Z . By the universal property of normalization, Theorem A.2, there is a map $p: Y_D \rightarrow \widehat{X}_D$ such that p^* is the identity on $K(X_D)$ and $j \circ p = q$. Since p^* is the identity on $K(X_D)$, we must have $p \circ i = \text{id}_{X_D}$.

The locus of points in Y_D representing stable Riemann surfaces whose Jacobian is $(\mathbb{C}^*)^2$ is the union $\bigcup C_P$. Thus $q^{-1}(\mathcal{A}_0) = \bigcup C_P$. We have $j^{-1}(\mathcal{A}_0) = C$, so $p^{-1}(C) = \bigcup C_P$. By Theorem A.4, Zariski's Main Theorem, the fiber of p over each point in C is connected, so p induces a bijection between the components of $\bigcup C_P$ and the points of C .

For each point $t \in \widehat{X}_D \setminus C$, the image $j(t)$ represents either an Abelian surface in $\mathcal{A}_2$ or an elliptic curve in $\mathcal{A}_1$. In the either case, there are only finitely many points in Y_D which map to $j(t)$, so the fiber $p^{-1}(t)$ is finite. This fiber is also connected by Zariski's Main Theorem, so it is a single point. This means that the restriction,

$$p: Y_D \setminus \bigcup_P C_P \rightarrow X_D \setminus C,$$

is a bijection, and it is thus an isomorphism because these are normal varieties.

The locus of points t in $\widehat{X}_{d^2}$ such that $j(t) \in \mathcal{A}_1$ is the union of $R_{d^2}^1$ and $R_{d^2}^2$, and the locus of points in Y_{d^2} whose Jacobians are extensions of elliptic curves is the union of $S_{d^2}^1$ and $S_{d^2}^2$. This means that p must take the curves $S_{d^2}^i$ to $R_{d^2}^i$, and the restriction of p to the $S_{d^2}^i$ is an isomorphism onto the $R_{d^2}^i$ because $S_{d^2}^i$ is disjoint from each C_P . $\blacksquare$

Cohomology of Y_D . We can use Theorem 8.10 to get information about the cohomology of Y_D . In $H^2(Y_D; \mathbb{Q})$, let B be the subspace generated by the fundamental classes of the rational curves C_P .

Theorem 8.11. *The intersection form on B is negative definite.*

Proof. Consider the map $q: Y'_D \rightarrow Y_D$ obtained by resolving the singularities at the points c_P of Y_D . Define $r: Y'_D \rightarrow \hat{X}_D$ by $r = p \circ q$, and let $E = r^{-1}(C)$. The locus E is the union of irreducible curves E_i . By possibly taking some blow-ups of Y'_D , we can assume that the curves E_i are nonsingular and transverse (actually, performing these blow-ups is not necessary; see the remark below).

Let $B' \subset H^2(Y'_D; \mathbb{Q})$ be the subspace spanned by the fundamental classes of the E_i . By Theorem A.10, the intersection form on B' is negative definite.

The map $q^*: B \rightarrow B'$ preserves the intersection forms on B and B' because q is degree one. We claim that $q^*: B \rightarrow B'$ is injective. To see this, define a linear map $r: B' \rightarrow B$ as follows. If $E_i \subset E$, and $q(E_i)$ is one of the curves C_P , let $r([E_i])$ be the fundamental class of that curve. Otherwise, $q(E_i)$ is a point; in this case let $r([E_i]) = 0$. We have $r \circ q^* = \text{id}_B$ on B , so q^* is injective on B .

Thus q^* embeds B in B' , preserving the intersection forms. Since the intersection form on B' is negative definite, the intersection form on B must be as well. $\blacksquare$

Remark. We can actually describe the variety Y'_D above very explicitly. The exceptional fiber of a resolution of a cyclic quotient singularity consists of a chain of rational curves $\{E_i\}_{i=1}^n$ with E_i having one transverse intersection with E_{i+1} if $i \leq i \leq n-1$. This is described in [vdG88].

Thus resolving each $c_P \in Y_D$ which is singular replaces each chain of curves C_P with a longer chain of rational curves with transverse crossings.

Since Y_D is an orbifold, Poincaré duality holds for rational cohomology, as Satake showed in [Sat56a]. This together with Theorem 8.11 means that we can define the orthogonal projection $\pi_B: H^2(Y_D; \mathbb{Q}) \rightarrow B$.

Corollary 8.12. *If D is not square, then*

$$\pi_B[\overline{P}_D] = \pi_B[\overline{W}_D].$$

as classes in $H^2(Y_D; \mathbb{Q})$. If $D = d^2$, then

$$\pi_B[\overline{P}_{d^2}] = \pi_B[\overline{W}_{d^2}] + \pi_B[\overline{S}_{d^2}^2]$$

Proof. For each curve C_P , the number of intersection points of C_P with $\overline{P}_D$ is equal to the number of intersection points of C_P with $\overline{W}_D$ and $\overline{S}_D^2$ by Part 4 of Theorem 8.7 (using the convention that S_D^2 is empty if D is not square). Since these intersection points are transverse, this implies

$$[C_P] \cdot [\overline{P}_D] = [C_P] \cdot ([\overline{W}_D] + [\overline{S}_D^2]).$$

The desired equations then follow because the intersection matrix of the C_P is nondegenerate. $\blacksquare$

8.3 Involution of Y_D

We now study the extension of the involution τ of X_D to Y_D . We then use this involution to deduce information about the fundamental classes of the curves $\overline{W}_D^\epsilon$.

Lemma 8.13. *Let $\Gamma \subset X_D \times \overline{\mathcal{M}}_2$ be the graph of the natural map $X_D \rightarrow \overline{\mathcal{M}}_2$, and let $\overline{\Gamma}$ be the closure of Γ in $\widehat{X}_D \times \overline{\mathcal{M}}_2$. Then there is a morphism $q: Y_D \rightarrow \overline{\Gamma}$ which is the normalization of $\overline{\Gamma}$.*

Proof. We have a map $q = p \times \pi: Y_D \rightarrow \widehat{X}_D \times \overline{\mathcal{M}}_2$ by Theorem 8.10, and the image of q is in $\overline{\Gamma}$. To show that $q: Y_D \rightarrow \overline{\Gamma}$ is the normalization of $\overline{\Gamma}$, we need only to show that q is finite and birational. The map q is birational because $p: Y_D \rightarrow \widehat{X}_D$ is birational, as is the projection $\overline{\Gamma} \rightarrow \widehat{X}_D$. The map q is finite because the map $\pi: Y_D \rightarrow \overline{\mathcal{M}}_2$ is finite. $\blacksquare$

Theorem 8.14. *The involution τ of X_D extends to an involution τ of Y_D . This involution satisfies*

$$\tau(S_D^1) = S_D^2, \quad (8.2)$$

and

$$\tau(C_P) = C_{t(P)}, \quad (8.3)$$

where t is the involution on the set of nondegenerate Y_D -prototypes defined in §3.1.

Proof. Let $\overline{\Gamma} \subset \widehat{X}_D \times \overline{\mathcal{M}}_2$ as in Lemma 8.13. The involution,

$$\tilde{\tau} = \tau \times \text{id}_{\overline{\mathcal{M}}_2}: \widehat{X}_D \times \overline{\mathcal{M}}_2 \rightarrow \widehat{X}_D \times \overline{\mathcal{M}}_2,$$

restricts to an involution $\tilde{\tau}$ of $\overline{\Gamma}$. By the universal property, Theorem A.2, the composition $\tilde{\tau} \circ q: Y_D \rightarrow \overline{\Gamma}$ lifts to a map $\tau: Y_D \rightarrow Y_D$ which extends τ on X_D . Since τ^2 is the identity on the open, dense subset X_D of Y_D , the map τ is an involution of Y_D .

The involution τ of $\widehat{X}_D$ sends R_D^1 to R_D^2 . Since the map $p: Y_D \rightarrow X_D$ commutes with the involutions, we must have $\tau(S_D^1) = S_D^2$ by Theorem 8.10.

We are regarding $X_D \subset \mathbb{P}\Omega\overline{\mathcal{M}}_2$ as the set of stable Abelian differentials $(X, [\omega])$, where X has a compact Jacobian, and $\text{Jac}(X)$ has real multiplication $\rho: \mathcal{O}_D \rightarrow \text{End}(\text{Jac}(X))$ with ω as an ι_1 -eigenform defined up to constant multiple. In these terms, $\tau(X, [\omega])$ is the pair $(X, [\omega'])$, where ω' is an ι_2 -eigenform for the same real multiplication on $\text{Jac}(X)$ (ω' is then an ι_1 -eigenform for the Galois conjugate real multiplication ρ').

Now suppose that $(X, [\omega]) \in X_D$ is close to a point in the curve C_P^0 , and $(X, [\omega']) = \tau(X, [\omega])$ is close to a point in $C_{P'}^0$. We must show that $P' = t(P)$.

Recall the definition of the prototype P from §7.6. On (X, ω) , there are two tall cylinders C_1 and C_2 . Let $\{\alpha_i, \beta_i\}_{i=1}^2$ be a symplectic basis of $H_1(X; \mathbb{Z})$ such that α_i represents the cylinder C_i and $\omega(\alpha_2)/\omega(\alpha_1) \geq 1$. Let $\mu \in K_D$ be such that $\mu \cdot \alpha_1 = \alpha_2$, and let $\psi(t) = at^2 + bt + c$ be as in Proposition 3.4, in particular $\psi(\mu) = 0$. For $\lambda \in K_D$, let T_λ be the matrix of the action of λ on $H_1(X; \mathbb{Q})$

in the symplectic basis $\{\alpha_i, \beta_i\}$. Then $T_{a\mu}$ is as in (7.9). The prototype is then $P = (a, b, c, \bar{q})$, where a , b , and c are as above, and $\bar{q}$ is the reduction of the upper right entry of $T_{a\mu}$ modulo $\gcd(a, b, c)$.

There are two cases to consider. First suppose that $\mu^{(2)} \leq -1$, which is equivalent to $a - b + c \leq 0$. Let $\{\alpha'_i, \beta'_i\}_{i=1}^2$ be the symplectic basis,

$$\{\alpha'_1, \alpha'_2, \beta'_1, \beta'_2\} = \{\alpha_1, -\alpha_2, \beta_1, -\beta_2\},$$

of $H_1(X; \mathbb{Z})$. Let $\tilde{\mu} = -\mu$, and normalize $[\omega']$ so that $\omega'(\alpha'_1) = 1$. Then $\tilde{\mu} \cdot \alpha'_1 = \alpha'_2$, and

$$\omega'(\alpha'_2) = \omega'(\tilde{\mu} \cdot \alpha'_1) = \tilde{\mu}^{(2)} \omega'(\alpha'_1) = -\mu^{(2)} \geq 1.$$

This means that we can use the symplectic basis $\{\alpha'_i, \beta'_i\}$ to calculate P' . If $\psi'(t) = a't^2 + b't + c'$ is the polynomial from Proposition 3.4 with $\psi'(\tilde{\mu}) = 0$, then

$$\psi'(t) = \psi(-t) = at^2 - bt + c,$$

so $(a', b', c') = (a, -b, c)$. For $\lambda \in K_D$, let T'_λ be the matrix T_λ in the new basis $\{\alpha_i, \beta_i\}$. Then a simple calculation shows,

$$T'_{a'\tilde{\mu}} = T'_{-a\mu} = \begin{pmatrix} 0 & -c & 0 & q \\ a & b & -q & 0 \\ 0 & 0 & 0 & a \\ 0 & 0 & -c & b \end{pmatrix}.$$

Thus $P' = (a, -b, c, \bar{q}) = t(P)$.

Now suppose that $-1 < \mu^{(2)} < 0$, which is equivalent to $a - b + c > 0$. Let $\{\alpha'_i, \beta'_i\}_{i=1}^2$ be the symplectic basis,

$$\{\alpha'_1, \alpha'_2, \beta'_1, \beta'_2\} = \{\alpha_2, -\alpha_1, \beta_2, -\beta_1\},$$

of $H_1(X; \mathbb{Z})$, let $\tilde{\mu} = -1/\mu$, and normalize ω' so that $\omega'(\alpha'_1) = 1$. Then $\tilde{\mu} \cdot \alpha'_1 = \alpha'_2$, and

$$\omega'(\alpha'_2) = -\mu^{(2)} > 1$$

as before. If $\psi'(t) = a't^2 + b't + c'$ is the polynomial from Proposition 3.4 with $\psi'(\tilde{\mu}) = 0$, then

$$\psi'(t) = -t^2 \psi(-1/t) = -ct^2 + bt - a,$$

so $(a', b', c') = (-c, b, -a)$. For $\lambda \in K_D$, let T'_λ be the matrix T_λ in the new basis $\{\alpha_i, \beta_i\}$. Then,

$$T'_{a'\tilde{\mu}} = T_{c/\mu} = ac(T_{a\mu})^{-1} = \begin{pmatrix} -b & c & 0 & q \\ -a & 0 & -q & 0 \\ 0 & 0 & -b & -a \\ 0 & 0 & c & 0 \end{pmatrix},$$

and in the new basis,

$$T'_{a'\bar{\mu}} = \begin{pmatrix} 0 & a & 0 & q \\ -c & -b & -q & 0 \\ 0 & 0 & 0 & -c \\ 0 & 0 & a & -b \end{pmatrix}.$$

Thus $P' = (-c, b, -a, \bar{q}) = t(P)$. ■

It is not true that $\tau(W_D^0) = W_D^1$, but we will prove that something like this is true on the level of cusps.

Theorem 8.15. *Suppose $D \equiv 1 \pmod{8}$. If D is not square, then for any nondegenerate Y_D -prototype,*

$$\#(\overline{W}_D^1 \cap C_P) = \#(\overline{W}_D^0 \cap \tau(C_P)). \quad (8.4)$$

If $D = d^2$, then

$$\#(\overline{W}_{d^2}^1 \cap C_P) + \#(\overline{S}_{d^2}^2 \cap C_P) = \#(\overline{W}_{d^2}^0 \cap \tau(C_P)). \quad (8.5)$$

Proof. Suppose P is a nondegenerate, nonterminal, noninitial Y_D -prototype. Recall that the cusps of W_D lying on C_P , where $P = (a, b, c, \bar{q})$, are the points w_Q , where $Q = (a, b, c, \bar{r})$ is a W_D -prototype with $\bar{r} \in \mathbb{Z}/\gcd(a, c)$, and $\bar{r} \equiv q \pmod{\gcd(a, b, c)}$. The cusp w_Q lies on $W_D^{\epsilon(Q)}$, where $\epsilon(Q)$ is given by (4.6).

There are two cases to consider, depending on whether or not $a - b + c \leq 0$. First suppose $a - b + c \leq 0$. In this case,

$$\epsilon(t(Q)) \equiv \epsilon(Q) + b \equiv \epsilon(Q) + 1 \pmod{2}$$

for any W_D -prototype Q associated to P , using the fact that b is odd because $b^2 - 4ac = D \equiv 1 \pmod{8}$. It follows that C_P has as many cusps of W_D^1 as $C_{t(P)}$ has cusps of W_D^0 .

Now suppose that $a - b + c > 0$. Here there are four cases to consider, depending on the parity of a and c . First suppose $a \equiv 1 \pmod{2}$, and $c \equiv 0 \pmod{2}$. In this case, for any W_D -prototype Q associated to P , we have $\epsilon(Q) \equiv 0 \pmod{2}$ and $\epsilon(t(Q)) \equiv 1 \pmod{2}$. Thus, every cusp of W_D on C_P is in W_D^0 , and $C_{t(P)}$ has the same number of cusps, all in W_D^1 .

Suppose $a \equiv 0 \pmod{2}$, and $c \equiv 1 \pmod{2}$. In this case, for any W_D -prototype Q associated to P , we have $\epsilon(Q) \equiv 1 \pmod{2}$ and $\epsilon(t(Q)) \equiv 0 \pmod{2}$. Thus, every cusp of W_D on C_P is in W_D^1 , and $C_{t(P)}$ has the same number of cusps, all in W_D^0 .

Suppose $a \equiv c \equiv 0 \pmod{2}$. In this case, for any W_D -prototype $Q = (a, b, c, \bar{r})$ associated to P , we have $\epsilon(Q) \equiv \epsilon(t(Q))\bar{r} \pmod{2}$. Since $\gcd(a, b, c) \equiv 1 \pmod{2}$, this means that exactly half of the cusps of W_D on C_P are on W_D^1 , and the same is true for $C_{t(P)}$.

The last case, $a \equiv c \equiv 1 \pmod{2}$ doesn't occur because $b^2 - 4ac \equiv 1 \pmod{8}$. Thus we have shown (8.4).

It remains to prove (8.5) when P is an initial or terminal prototype. We claim that if P is an initial Y_{d^2} -prototype, then C_P intersects $\overline{W}_{d^2}^0$ once and is disjoint from $\overline{W}_{d^2}^1$. Let $P = (a, b, c, \bar{q})$ be an initial prototype. Since $a - b + c = 0$,

$$\gcd(a, c) = \gcd(a, b, c),$$

so $\text{mult}(P) = 1$, and $\overline{W}_{d^2}$ intersects C_P once. We have

$$\begin{aligned} b &= a + c \\ \Rightarrow (a + c)^2 - 4ac &= d^2 \\ \Rightarrow a - c &= d \\ \Rightarrow a &\equiv c + 1 \pmod{2}, \end{aligned}$$

and furthermore $f = d$. It follows that $\epsilon(P) = 0$, thus the intersection point of C_P with $\overline{W}_{d^2}$ is in $\overline{W}_{d^2}^0$.

We saw in Theorem 8.7 that if P is a terminal Y_D -prototype, then C_P intersects $\overline{S}_{d^2}^2$ once and is disjoint from $\overline{W}_{d^2}$. Thus if P is a terminal prototype, then both sides of (8.5) are one, and if P is an initial prototype, then both sides of (8.5) are zero. $\blacksquare$

Corollary 8.16. *If D is not square, then*

$$\tau_* \pi_B[\overline{W}_D^0] = \pi_B[\overline{W}_D^1] \quad (8.6)$$

in $H^2(Y_D; \mathbb{Q})$. If $D = d^2$, then

$$\tau_* \pi_B[\overline{W}_{d^2}^0] = \pi_B[\overline{W}_{d^2}^1] + \pi_B[\overline{S}_{d^2}^2] \quad (8.7)$$

in $H^2(Y_{d^2}; \mathbb{Q})$.

Proof. Since the intersection pairing on $B \subset H^2(Y_D; \mathbb{Q})$ is nondegenerate by Theorem 8.11, it suffices to show that

$$[C_P] \cdot (\tau_* \pi_B[\overline{W}_D^0]) = [C_P] \cdot \pi_B[\overline{W}_D^1],$$

and

$$[C_P] \cdot (\tau_* \pi_B[\overline{W}_{d^2}^0]) = [C_P] \cdot (\pi_B[\overline{W}_{d^2}^1] + \pi_B[\overline{S}_{d^2}^2])$$

for each C_P . Since the intersections of $\overline{W}_D$ and $\overline{S}_{d^2}^2$ with each C_P are transverse, this follows directly from Theorem 8.15. $\blacksquare$

9 Line bundles over Y_D

In this section, we will define extensions of some line bundles over X_D to Y_D , and we will calculate their Chern classes. In §9.1, we discuss background material on vector bundles over orbifolds. In §9.2, we recall Mumford's notion of a good Hermitian metric, which allows one to express the Chern classes of a vector bundle in terms of curvatures of singular Hermitian metrics. In §9.3, we discuss the extension of ΩX_D to Y_D , and we discuss the extension of $T^* \mathcal{A}_D$ to Y_D in §9.4.

9.1 Orbifold vector bundles

Most of the theory of vector bundles over manifolds works for vector bundles over orbifolds. Here following [CR02] we discuss the facts that we need and refer the reader to the Appendix to [CR02] for details.

Let X be a complex orbifold with an atlas $\{U_\alpha/G_\alpha \rightarrow X\}$, where $U_\alpha \subset \mathbb{C}^n$, and G_α is a finite automorphism group of U_α (we allow G_α to have elements which act trivially on U_α , and X is said to be *reduced* if no such element exists). A *rank r holomorphic orbifold bundle over X* is an orbifold E with a map $\pi: E \rightarrow X$ such that for each α ,

$$\pi^{-1}(U_\alpha) \cong (U_\alpha \times \mathbb{C}^r)/G_\alpha,$$

where G_α acts on $U_\alpha \times \mathbb{C}^r$ by

$$g \cdot (z, v) = (g \cdot z, \rho(z, g)(v))$$

for some $\rho: U_\alpha \times G_\alpha \rightarrow \mathrm{GL}_r \mathbb{C}$ which is a homomorphism if we fix z and is holomorphic if we fix g .

Nonreduced orbifold structures occur naturally. For example, $\mathcal{M}_2$ has a nonreduced orbifold structure, where the local group G at a point X is $\mathrm{Aut}(X)$. This is nonreduced because the hyperelliptic involution acts trivially. The associated reduced orbifold structure has local group $G = \mathrm{Aut}(X)/J$ at X . With this definition of an orbifold vector bundle, $\Omega \mathcal{M}_2$ is only an orbifold vector bundle if we take the nonreduced orbifold structure on $\mathcal{M}_2$. When necessary, we will implicitly use this orbifold structure. Similarly, X_D has a nonreduced orbifold structure with local group $G = \mathrm{Aut}(A, \rho)$ at (A, ρ) .

We want to be able to pull back bundles along maps between orbifolds. The operation of pulling back a bundle is much more complicated in the category of orbifolds than in the category of topological spaces. In particular, given a map $f: X \rightarrow Y$ of orbifolds, it is not always possible to pull back a bundle over Y by f ; however, this is possible if f is what is called a *good* map. We will informally discuss pullbacks of orbifold bundles without actually defining a good map.

Given a map $f: X \rightarrow Y$ of orbifolds and charts $V/H \subset Y$ and $U/G \subset X$ such that $f(U/G) \subset V/H$, there is a lift $\tilde{f}: U \rightarrow V$ such that for each $g \in G$ there is some $h \in H$ such that

$$\tilde{f}(g \cdot z) = h \cdot \tilde{f}(z).$$

Note that there may not be a group homomorphism $\sigma: G \rightarrow H$ such that

$$\tilde{f}(g \cdot z) = \sigma(g)\tilde{f}(z).$$

If f is good in the sense of [CR02], then such a homomorphism does exist. The actual definition of a good map is some complicated global condition which locally amounts to the existence of such a homomorphism; we will not get into this here.

In the notation of the previous paragraph, suppose that $f: U/G \rightarrow V/H$ is good and we have over V/H a vector bundle $E = (V \times \mathbb{C}^r)/H \rightarrow V/H$, with H acting on $V \times \mathbb{C}^r$ by $\rho: V \times H \rightarrow \mathrm{GL}_r(\mathbb{C})$. We define f^*E over U by

$$f^*E = (U \times \mathbb{C}^r)/G, \tag{9.1}$$

where G acts on $U \times \mathbb{C}^r$ by $g \cdot (z, v) = (g \cdot z, \tau(z, g)(v))$, with

$$\tau(z, g) = \rho(\tilde{f}(z), \sigma(g)).$$

Globally, the pullback of a bundle $E \rightarrow Y$ along a map $f: X \rightarrow Y$ is an orbifold vector bundle f^*E which locally satisfies (9.1).

Given an orbifold X , write X_{reg} for the set of p points of X such that for a chart $U/G \ni p$, no element of G fixes p . The following theorem follows from Lemmas 4.4.3 and 4.4.11 of [CR02].

Theorem 9.1. *Let $f: X \rightarrow Y$ be a map of orbifolds such that $f^{-1}(Y_{\mathrm{reg}})$ is an open, dense, and connected subset of X . Then f is a good map of orbifolds, and for any orbifold vector bundle $E \rightarrow Y$, there is a well-defined pullback bundle $f^*E \rightarrow X$. The pullback bundle satisfies the naturality property,*

$$c_1(f^*E) = f^*c_1(E).$$

Remark. Chern classes for orbifold vector bundles are defined using the Chern-Weil construction, which associates a de Rham cohomology class to a metric on an orbifold bundle. This theory is worked out in [CR02].

9.2 Good metrics

It is well known that the Chern classes of a complex vector bundle can be given in terms of the curvature of a Hermitian metric. In [Mum77], Mumford showed that for a Hermitian metric with certain mild singularities, called a *good metric*, the Chern classes of the vector bundle can still be given in terms of the curvature of the metric. Mumford's results are in terms of nonsingular complex projective varieties, but his results and proofs work just as well in the setting of complex orbifolds. In this section, following [Mum77], we will summarize what we need about good metrics, translating to the setting of orbifolds.

Let $X \subset \overline{X}$ be complex n -dimensional orbifolds with $D = \overline{X} - X$ a divisor and $\overline{X}$ compact. Suppose that we can cover D with coordinate charts Δ^n/G , where $\Delta^n \subset \mathbb{C}^n$ is a polydisk, and G is a finite group of automorphisms of Δ^n such that:

- Each transformation $g \in G$ is of the form,

$$g(z_1, \dots, z_n) = (\theta_1 z_1, \dots, \theta_n z_n),$$

for some roots of unity θ_i .

- $D \cap \Delta$ is a union of the coordinate axes $z_1 = 0, \dots, z_r = 0$ for $1 \leq r \leq n$.

In such a neighborhood Δ^n/G , we have $\Delta^n \cap X = (\Delta^*)^r \times \Delta^{n-r}$. We give $\Delta^n \cap X$ a metric by putting the Poincaré metric,

$$ds^2 = \frac{|dz|^2}{|z|^2(\log|z|)^2},$$

on the Δ^* factors and putting the Euclidean metric $|dz|^2$ on the Δ factors. Call the product metric ω . This metric is G -invariant, so it is a metric on Δ^n/G in the orbifold sense.

Definition. A p -form η on X has *Poincaré growth* if there is a cover of $\overline{X} \setminus X$ by polycylinders Δ_α^n/G_α as above such that in each Δ_α^n we have some constant C_α such that for any p vectors t_i at any point $x \in \Delta_\alpha^n \cap X$,

$$|\eta(t_1, \dots, t_p)|^2 < C_\alpha \omega_\alpha(t_1, t_1) \cdots \omega_\alpha(t_p, t_p), \quad (9.2)$$

where ω_α is the metric on Δ_α^n/G_α defined above.

In this paper, given a differential form η , we will write $\langle \eta \rangle$ for the current defined by η , and $[\eta]$ for the de Rham cohomology class defined by η .

Proposition 9.2. *A p -form η with Poincaré growth satisfies,*

$$\int_X |\eta \wedge \zeta| < \infty$$

for any smooth form ζ on $\overline{X}$ of complementary dimension. Thus, η defines a p -current $\langle \eta \rangle$.

Definition. A p -form η in X is *good* if η and $d\eta$ both have Poincaré growth.

Proposition 9.3. *If η is a good p -form, then*

$$d\langle \eta \rangle = \langle d\eta \rangle.$$

Note that if η is a closed good form, it is not necessarily true that $\eta = d\zeta$ for a good form ζ ; however, this will be the case if η is the Chern form of a *good* Hermitian metric.

Let $\overline{L}$ be a holomorphic orbifold line bundle over $\overline{X}$ with L the restriction to X , and let H be a Hermitian metric on E .

Definition. H is a good metric if for every point $x \in D$, and every polycylindrical neighborhood Δ^n/G as described above with D given by

$$\prod_{i=1}^k z_i,$$

and for every holomorphic section e of $\overline{E}$ over Δ^n/G , setting $h = H(e, e)$, we have

•

$$|h|, |h|^{-1} < C \left(\sum_{i=1}^k \log |z_i| \right)^{2m}$$

for some $C > 0$ and $m > 1$.

- The 1-forms $(\partial h)/h$ are good on $\Delta^n/G \cap X$.

Theorem 9.4. *If H is a good metric, then the Chern form, $c_1(E, H)$ is good and the current $\langle c_1(E, H) \rangle$ represents the Chern class $c_1(\overline{E})$ in $H^2(\overline{X}; \mathbb{Q})$. Furthermore,*

$$c_1(E, H) - d\eta$$

is a smooth 2-form on $\overline{X}$ for some good 1-form η .

The last statement follows from the proof of Theorem 1.4 in [Mum77].

We now show that Chern forms of good metrics behave as they should with respect to the cup product pairing and integrating over curves.

Proposition 9.5. *Suppose $\dim_{\mathbb{C}} \overline{X} = 2$. If ω and η are Chern forms of two good metrics, then*

$$\langle [\omega], [\eta] \rangle = \int_X \omega \wedge \eta. \quad (9.3)$$

Proof. By Theorem 9.4, we can write

$$\begin{aligned} \omega &= \omega' + d\omega'' \\ \eta &= \eta' + d\eta'', \end{aligned}$$

where ω' and η' are smooth, closed 2-forms on $\overline{X}$, and ω'' and η'' are good 1-forms. We have,

$$\begin{aligned} \langle [\omega], [\eta] \rangle &= \int_{\overline{X}} \omega' \wedge \eta' \\ &= \int_X (\omega \wedge \eta - \omega \wedge d\eta'' - d\omega'' \wedge \eta + d\omega'' \wedge d\eta'') \\ &= \int_X (\omega \wedge \eta + d(\omega \wedge \eta'') - d(\omega'' \wedge \eta) + d(\omega'' \wedge d\eta'')). \end{aligned} \quad (9.4)$$

We claim that $\int_X d\alpha = 0$ for any good 3-form α . The last three terms in (9.4) are of this form, so (9.3) follows from this claim. By Proposition 9.3,

$$\int_X d\alpha = \langle d\alpha \rangle(1) = d\langle \alpha \rangle(1) = -\langle \alpha \rangle(d1) = 0, \quad (9.5)$$

so the claim follows. $\blacksquare$

Proposition 9.6. *Suppose $C \subset \overline{X}$ is a curve with no irreducible component contained in D , and suppose ω is the Chern form of a good Hermitian metric. Then*

$$\langle [C], [\omega] \rangle = \int_C \omega.$$

Sketch of proof. Let $\omega = \omega' + d\omega''$ with ω' a smooth 2-form on $\overline{X}$ and ω'' a good 1-form. We have

$$\langle [C], [\omega] \rangle = \int_C \omega' = \int_C \omega - \int_C d\omega''.$$

We need to show that

$$\int_C |\omega| < \infty \quad (9.6)$$

and

$$\int_C d\omega'' = 0. \quad (9.7)$$

Let $C_0 = C \setminus C \cap D$. We can speak of Poincaré growth for forms on C_0 using the compactification of C_0 obtained by adding a point to each cusp of C_0 . It follows from the Schwartz Lemma that any good p -form on X restricts to a good p -form on C_0 , so $\omega|_{C_0}$ and $\omega''|_{C_0}$ are good. Then (9.6) follows from Proposition 9.2, and (9.7) follows from (9.5). $\blacksquare$

9.3 Extension of the bundle ΩX_D to Y_D

Extension of ΩX_D . Recall that we have the map $\pi: Y_D \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$ from Theorem 8.10. Let

$$\pi_1: \mathbb{P}\Omega\overline{\mathcal{M}}_2 \rightarrow \overline{\mathcal{M}}_2$$

be the natural projection, and let

$$\pi_2 = \pi_1 \circ \pi: Y_D \rightarrow \overline{\mathcal{M}}_2.$$

Let $\Omega^0 Y_D = \pi_2^*(\Omega\overline{\mathcal{M}}_2)$. It does not follow directly from Theorem 9.1 that this pullback is well defined because we must take the nonreduced orbifold structure on $\overline{\mathcal{M}}_2$ for $\Omega\overline{\mathcal{M}}_2$ to be a bundle, and then $\pi_2^{-1}(\overline{\mathcal{M}}_2)_{\text{reg}} = Y_D$. Instead, first pull back the square $(\Omega\overline{\mathcal{M}}_2)^{\otimes 2}$. This is a bundle over $\overline{\mathcal{M}}_2$ with the usual reduced orbifold structure, so $\pi_2^{-1}(\overline{\mathcal{M}}_2)_{\text{reg}}$ consists of the elliptic points of X_D together with the elliptic points of $S_{d^2}^i$ and the curves C_P . This is an open, dense, and connected subset of Y_D , thus the pullback of the square is well defined by Theorem 9.1. Then define $\Omega^0 Y_D$ to be the quotient of this pullback by ± 1 .

Given $p \in Y_D$ with $\pi_2(p) = X \in \overline{\mathcal{M}}_2$, the fiber of $\Omega^0 Y_D$ over p is isomorphic to $\Omega(X)/G_p$ for some finite group G_p . If p is not an orbifold point of Y_D , then $G = \{\pm 1\}$. If p is an elliptic point of X_D or S_D^i , then $G_p = \text{Aut}(X)$. If $p = c_P$ with P a nondegenerate, nonterminal Y_D -prototype, then *a priori* G_p could be bigger than $\{\pm 1\}$ because c_P is a singular point of Y_D . In fact, $\pi(c_P)$ is a nonsingular point of $\mathbb{P}\Omega\overline{\mathcal{M}}_2$ because the residues at the nodes of $\pi(c_P)$ are all different, so by (9.1), $G_p \cong \{\pm 1\}$. Thus we have shown that the fiber of $\Omega^0 Y_D$ over p is $\Omega(X)/\{\pm 1\}$ unless p is an elliptic point of X_D or S_D^i .

Over $\mathbb{P}\Omega\overline{\mathcal{M}}_2$, there is the canonical orbifold line bundle $\mathcal{O}(-1) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_2$ whose fiber over $(X, [\omega])$ is the subspace of $\Omega(X)$ spanned by ω . Define

$$\begin{aligned}\Omega^1 Y_D &= \pi^* \mathcal{O}(-1) \\ \Omega^2 Y_D &= \tau^* \Omega^1 Y_D.\end{aligned}$$

Since $\mathcal{O}(1) \subset \pi_1^* \Omega\overline{\mathcal{M}}_2$ as a sub line bundle, $\Omega^1 Y_D$ is a sub line bundle of $\Omega^0 Y_D$. The involution τ lifts to an involution of $\Omega^0 Y_D$, so we can also regard $\Omega^2 Y_D$ as a sub line bundle of $\Omega^0 Y_D$. Given $p \in Y_D$ with $\pi_2(p) = X$, the Jacobian $\text{Jac}(X)$ comes with a choice of real multiplication by $\mathcal{O}_D$, and we can regard the fiber of $\Omega^i Y_D$ over p as $\Omega^i(X)$, the space of ι_i -eigenforms. Since

$$\Omega(X) = \Omega^1(X) \oplus \Omega^2(X),$$

we obtain

$$\Omega^0 Y_D = \Omega^1 Y_D \oplus \Omega^2 Y_D. \quad (9.8)$$

Define

$$Q^i Y_D = (\Omega^i Y_D)^2,$$

which parameterizes quadratic differentials which are squares of ι_i -eigenforms. In what follows, we will sometimes abbreviate $\Omega^1 Y_D$ and $Q^1 Y_D$ by ΩY_D and $Q Y_D$.

Proposition 9.7. *The restriction of $Q Y_D$ to each curve $C_P \subset Y_D$ is trivial.*

Sketch of proof. Define a global nonzero section of $Q Y_D$ over C_P as follows. If $X \in C_P^0$, then let $q_X \in Q(X)$ be the unique quadratic differential which is the square of the Abelian differential $\omega_X \in \Omega(X)$ which has residue 1 at both nonseparating nodes of X . Otherwise, X has three nonseparating nodes. In this case, define q_X in the same way, using the two nodes of X which are limits of nodes of surfaces in C_P^0 . ■

Chern class of ΩY_D . In §2.3, we gave ΩX_D a Hermitian metric h_Ω . In terms of Riemann surfaces, the metric is given on the fiber over X by

$$h_\Omega(\omega, \omega) = \int_X |\omega|^2.$$

The induced metric h_Q on $Q X_D$ is given in the fiber over X by

$$h_Q(q, q) = \left(\int_X |q| \right)^2.$$

On Y_D , these metrics become singular along the curves C_P and $\bar{S}_D^1$ because stable Abelian differentials representing points on these curves have infinite area.

Theorem 9.8. *The metric h_Q on QY_D is a good metric.*

Lemma 9.9. *Let H be a singular Hermitian metric on the trivial bundle $L = \Delta^n \times \mathbb{C}$ over the polydisk $\Delta^n \in \mathbb{C}^n$, and let s be a holomorphic, nonzero section with*

$$H(s, s) = \left(\sum_{i=1}^r c_i \log |z_i| + c \right)^2,$$

for some constants c_i and c with $c_i < 0$. Then H is a good metric on L .

Proof. Let $h = H(s, s)$. Clearly,

$$h, h^{-1} < O \left(\sum_{i=1}^r \log |z_i| \right)^2,$$

so it remains to show that $\alpha = \partial h / h$ is a good 1-form.

We have,

$$\alpha = \frac{\sum_i \frac{c_i dz_i}{z_i}}{\sum_i c_i \log |z_i| + c},$$

so

$$\begin{aligned} \left| \alpha \left(\frac{\partial}{\partial z_j} \right) \right| &= \frac{c_j}{|z_j| |\sum_i c_i \log |z_i| + c|} \\ &\leq \frac{1}{|z_j| \log |z_j|} \\ &= \omega \left(\frac{\partial}{\partial z_j}, \frac{\partial}{\partial z_j} \right)^{1/2}. \end{aligned}$$

Thus α has Poincaré growth. We have,

$$d\alpha = \frac{1}{2} \frac{\sum_{ij} c_i c_j \frac{dz_i \wedge d\bar{z}_j}{z_i \bar{z}_j}}{(\sum_i c_i \log |z_i| + c)^2},$$

so

$$\begin{aligned} \left| d\alpha \left(\frac{\partial}{\partial z_k}, \frac{\partial}{\partial \bar{z}_l} \right) \right| &= \frac{1}{2} \frac{c_k c_l}{|z_k z_l| (\sum_i c_i \log |z_i| + c)^2} \\ &\leq \frac{1}{4} \frac{1}{|z_k| \log |z_k|} \frac{1}{|z_l| \log |z_l|} \\ &= \frac{1}{4} \omega \left(\frac{\partial}{\partial z_k}, \frac{\partial}{\partial z_k} \right)^{1/2} \omega \left(\frac{\partial}{\partial \bar{z}_l}, \frac{\partial}{\partial \bar{z}_l} \right)^{1/2}. \end{aligned}$$

Thus, $d\alpha$ also has Poincaré growth, so α is a good 1-form, and H is a good metric. ■

Proof of Theorem 9.8. We need to give holomorphic sections of QY_D around every point of $\overline{S}_D^1$ and C_P and show that the norms of these sections satisfy the required bounds.

If $p \in \overline{S}_{d^2}^1$, then we saw in §7.3 and §7.5 that $\overline{X}_{d^2}$ is normal around p , so we can work in $\overline{X}_{d^2}$. We gave a section s of $\mathcal{O}(-1)$ around p whose norm is given by (7.3) or (7.7). The norm of s^2 as a section of $\mathcal{O}(-2)$ is the square of the norm of s , and s^2 can be regarded as a section of QY_D around p . Thus h_Q is good around each $p \in \overline{S}_{d^2}^1$ by Lemma 9.9.

If $p \in C_P^0$, then we can consider $\pi(p)$ as a point in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,0})$ or $\mathbb{P}\Omega\mathcal{D}_2^0(T_{2,1})$. As we saw in §7.6, $\pi(p)$ is contained in a hypersurface U_λ with coordinates (x, y, z) on U_λ . The variety $\overline{X}_D$ is contained in U_λ near $\pi(p)$ and is cut out by the equation, (7.12). In a neighborhood of p in Y_D , there are coordinates (u, v) with

$$x = u, \ y = v^q, \text{ and } z = \theta v^r,$$

for some $p, q \in \mathbb{N}$ and root of unity θ by Proposition 8.1. We defined a section s of $\mathcal{O}(-1)$ over U_λ with norm given by (7.18). The pullback of s^2 to Y_D gives a section t of QY_D with norm,

$$h_Q(t, t) = (c_1 \log |v| + c)^2,$$

for some constants c_1 and c with $c_1 < 0$. Thus by Lemma 9.9, h_Q is good around every $p \in C_P^0$.

Now suppose $p = c_P$ for some prototype P . We can consider $\pi(p)$ as a point in $\mathbb{P}\Omega\mathcal{D}_2^0(T_{3,0})$ or $\mathbb{P}\Omega\mathcal{D}_2^0(T_{3,0}^3)$. As we saw in §7.7, $\pi(p)$ is contained in a hypersurface U_λ with coordinates (x, y, z) on U_λ . The variety $\overline{X}_D$ is contained in U_λ near $\pi(p)$ and is cut out by the equation, (7.21). In a neighborhood of p in Y_D , there are coordinates (u, v) with

$$x = u^p v^q, \ y = u^r, \text{ and } z = v^s,$$

for some $p, q, r, s \in \mathbb{N}$ by Proposition 8.3. We defined a section s of $\mathcal{O}(-1)$ over U_λ with norm given by (7.23). The pullback of s^2 to Y_D gives a section t of QY_D with norm,

$$h_Q(t, t) = (c_1 \log |u| + c_2 \log |v|)^2,$$

for some negative constants c_1 and c_2 . Thus by Lemma 9.9, h_Q is good around every c_P . ■

Corollary 9.10. *The first Chern class of $Q^i Y_D$ is*

$$c_1(Q^i Y_D) = [\omega_i]. \tag{9.9}$$

Proof. Since h_Q is a good metric, $c_1(QY_D, h_Q)$ represents $c_1(QY_D)$ by Theorem 9.4. We showed in Proposition 2.8 that $c_1(QY_D, h_Q) = \omega_1$. This shows that $c_1(Q^1 Y_D) = [\omega_1]$. Also,

$$c_1(Q^2 Y_D) = c_1(\tau^* Q^1 Y_D) = \tau^* c_1(Q^1 Y_D) = [\omega_2].$$

■

Corollary 9.11. *For $i = 1, 2$, and for any nondegenerate Y_D -prototype,*

$$[\omega_i] \cdot [C_P] = 0.$$

Proof. This pairing is equal to the degree of QY_D restricted to C_P . Since this restriction is trivial by Proposition 9.7, the degree is zero. ■

9.4 Extension of the foliation $\mathcal{A}_D$ to Y_D

Proposition 9.12. *The foliation $\mathcal{A}_D$ of X_D extends to a foliation (which we will continue to call $\mathcal{A}_D$) of*

$$Y_D \setminus \bigcup_P C_P,$$

where the union is over all nonterminal Y_D -prototypes. The curves C_P^0 and S_D^1 are leaves of $\mathcal{A}_D$, and $\mathcal{A}_D$ is transverse to $\overline{S}_D^2$.

Proof. It follows directly from the equations for $\mathcal{A}_D$ given in Corollaries 7.7, and 7.12 that $\mathcal{A}_D$ extends over S_D^1 and S_D^2 to a foliation which contains S_D^1 as a leaf and is transverse to S_D^2 . The equations for $\mathcal{A}_D$ in Corollary 7.15 show that $\mathcal{A}_D$ extends over the intersection points of S_D^1 and S_D^2 and is transverse to S_D^2 there. For $p \in C_P^0$, equations for $\mathcal{A}_D$ near $\pi(p) \in \overline{X}_D$ are given in Theorem 7.22. There are local coordinates (u, v) in a neighborhood of p in Y_D such that $x = u$, $y = v^p$, and $z = \theta v^q$ for some $p, q \in \mathbb{N}$ and root of unity q . In these coordinates, C_P is cut out by $v = 0$, and $\mathcal{A}_D$ is given by $v = \text{const}$. Thus $\mathcal{A}_D$ extends over C_P^0 to a foliation which contains C_P^0 as a leaf. ■

We will now study $T^*\mathcal{A}_D$, the cotangent bundle to the leaves of $\mathcal{A}_D$. Since every orbifold line bundle on a complex orbifold minus a subvariety of codimension at least two extends to an orbifold line bundle over the entire orbifold, we can regard $T^*\mathcal{A}_D$ as an orbifold bundle over all of Y_D , even though the foliation is singular at the points c_P . The holomorphic sections of $\mathcal{A}_D$ over an open set U are exactly the holomorphic sections over U minus the singular points of $\mathcal{A}_D$.

Proposition 9.13. *The only singular fibers of $T^*\mathcal{A}_D$ are over the elliptic points of X_D and the elliptic points of $S_{d^2}^i$. For each nonterminal Y_D -prototype P , the restriction of $T^*\mathcal{A}_D$ to the curve C_P is trivial. For each terminal Y_D -prototype, the restriction of $T^*\mathcal{A}_D$ to the curve C_P is isomorphic to $T^*C_P(c_P-)$.*

Proof. Besides the elliptic points, the only points over which $T^*\mathcal{A}_D$ could have singular fibers are the singular points c_P of $\mathcal{A}_D$. Every section of an orbifold line bundle vanishes on a singular fiber, so to show that the fiber over c_P is nonsingular, it suffices to show that $T^*\mathcal{A}_D$ has a nonzero section over c_P .

Let (x, y, z) be the coordinates on in a neighborhood of $\pi(c_P)$ in the hypersurface $U_\lambda \subset \mathbb{P}\Omega\mathcal{D}_2(T_{3,0})$ as on p. 108. By Theorem 7.27 and Proposition 8.3, a neighborhood of c_P in Y_D is of the form Δ^2/G , where Δ^2 is a polydisk with coordinates (u, v) such that

$$x = u^p v^q, \quad y = u^r, \quad z = v^s$$

for some $p, q, r, s \in \mathbb{N}$, and G is a cyclic group whose action on Δ^2 is generated by

$$(u, v) = (\theta u, \theta^n v)$$

for some root of unity θ . By the equation,

$$y^\lambda z^{\lambda-1},$$

for the foliation $\mathcal{A}_D$ from Theorem 7.27, $\mathcal{A}_D$ is given on Δ^2 by the equation

$$u^a v^b = \text{const}$$

for some $a, b \geq 0$. If P is nonterminal, then $\lambda > 1$, and $a, b > 0$. Otherwise $\lambda = 1$, and $b = 0$, so the foliation is nonsingular at c_P . Define a vector field X on Δ^2 by

$$X = bu \frac{\partial}{\partial u} - av \frac{\partial}{\partial v}.$$

X is tangent to $\mathcal{A}_D$ and is invariant under G , so it gives a nonzero section of $T^*\mathcal{A}_D$ in a neighborhood of c_P . Thus the fiber of $T^*\mathcal{A}_D$ over c_P is not singular.

The restriction of X to C_P is a vector field on C_P in a neighborhood of c_P which vanishes at c_P . That means that a vector field Y on C_P defines a nonzero section of $T^*\mathcal{A}_D|_{C_P}$ if and only if Y vanishes at c_P . If P is nonterminal, then C_P has two points c_P and c_{P-} which pass through singular points of $\mathcal{A}_D$. A holomorphic vector field on C_P which has a zero at each of these points and no other zeros determines a nonzero holomorphic section of $T^*\mathcal{A}_D|_{C_P}$. Thus this restriction is trivial.

If P is terminal, then c_P is a nonsingular point of $\mathcal{A}_D$. Thus, the only singular point of $\mathcal{A}_D$ which C_P intersects is c_{P-} , so

$$T^*\mathcal{A}_D|_{C_P} \cong T^*C_P(c_{P-}).$$

■

Corollary 9.14. *The pairings of $c_1(Y^*\mathcal{A}_D)$ with the fundamental classes $[C_P]$ are*

$$c_1(T^*\mathcal{A}_D) \cdot [C_P] = \begin{cases} 0 & \text{if } P \text{ is not terminal;} \\ -1 & \text{if } P \text{ is terminal.} \end{cases}$$

Proof. The pairing $c_1(T^*\mathcal{A}_D) \cdot [C_P]$ is equal to the degree of the restriction of $T^*\mathcal{A}_D$ to C_P . If P is nonterminal, then this restriction is trivial, so the pairing is zero. If P is terminal, then this restriction has degree -1 by Proposition 9.13. ■

Chern class of $T^*\mathcal{A}_D$ when D is not square. We now calculate the first Chern class of $T^*\mathcal{A}_D$ by relating it to the bundle Q^2Y_D .

Lemma 9.15. *If D is not square, and $\mathcal{L} \rightarrow Y_D$ is an orbifold line bundle with a nonzero section $s \in \Gamma(X_D, \mathcal{L})$ defined over X_D , then S is a meromorphic section of $\mathcal{L}$ over Y_D .*

Proof. Let $\mathcal{O}(\mathcal{L})$ be the sheaf of sections of $\mathcal{L}$. By [Ser56], $\mathcal{O}(\mathcal{L})$ is a coherent algebraic sheaf on Y_D . We claim that s is defined algebraically over X . It would follow from this because every algebraic section of a line bundle over a dense, Zariski-open subset is meromorphic by [Har77, Lemma II.5.3].

The direct image $\pi_*\mathcal{O}(\mathcal{L})$ is a coherent algebraic sheaf on $\widehat{X}_D$. Given an open set $U \subset \widehat{X}_D$ and $t \in \Gamma(U, \pi_*\mathcal{O}(\mathcal{L}))$, let $r = t/s$, a holomorphic function on $U \setminus C$, where $C = \widehat{X}_D \setminus X_D$ is the set of cusps of X_D . By Koecher's principle (see [vdG88]), r extends uniquely to a holomorphic function $\tilde{r}$ on U . This defines an injective analytic map of sheaves,

$$i: \pi_*\mathcal{O}(\mathcal{L}) \rightarrow \mathcal{O}_{\widehat{X}_D},$$

with $i(t) = \tilde{r}$. By [Ser56], i is actually defined algebraically. Since $i(s) = 1$, and 1 is algebraic, it follows that s is an algebraic section of $\mathcal{L}$ over X_D . $\blacksquare$

Theorem 9.16. *When D is not square,*

$$Q^2Y_D \cong T^*\mathcal{A}_D.$$

Proof. Define

$$\mathcal{L} = Q^2Y_D \otimes (T^*\mathcal{A}_D)^{-1}.$$

By Propositions 2.8 and 2.9,

$$Q^2Y_D|_{X_D} \cong L_2 \cong T^*\mathcal{A}_D|_{X_D},$$

so there is a nonzero, holomorphic section s of $\mathcal{L}$ over X_D , which is a meromorphic section of $\mathcal{L}$ over Y_D by Lemma 9.15.

The divisor of s is

$$(s) = \sum e_P C_P,$$

so

$$c_1(\mathcal{L}) = \sum e_P [C_P].$$

We need to show that the e_P are all zero, for then s would be nonzero and holomorphic, so $\mathcal{L}$ would be trivial. By Propositions 9.7 and 9.13, the restriction of $\mathcal{L}$ to each C_P is trivial, so

$$[C_Q] \cdot \sum e_P [C_P] = [C_Q] \cdot c_1(\mathcal{L}) = 0,$$

for each Y_D -prototype Q . It then follows from Theorem 8.11 that $e_P = 0$ for each P . $\blacksquare$

Corollary 9.17. *If D is not square, then the first Chern class of $T^*\mathcal{A}_D$ is*

$$c_1(T^*\mathcal{A}_D) = [\omega_2]. \quad (9.10)$$

Chern class of $T^*\mathcal{A}_{d^2}$. The foliation $\mathcal{A}_{d^2}$ of X_{d^2} extends to a foliation $\widehat{\mathcal{A}}_{d^2}$ of the orbifold $\widehat{X}_{d^2}$. Give $T\widehat{\mathcal{A}}_{d^2}$ the metric ρ induced by the hyperbolic metric along the leaves of $\mathcal{A}_D$. This metric is singular along $\overline{R}_{d^2}^2$.

Theorem 9.18. *The metric ρ is a good metric for $T\widehat{\mathcal{A}}_{d^2}(-\overline{R}_{d^2}^2)$.*

Proof. Let $G \subset \mathrm{SL}_2\mathcal{O}_{d^2}$ be the cyclic subgroup generated by

$$\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & d \\ 0 & 1 \end{pmatrix} \right).$$

We then have a map,

$$\mathbb{H} \times \mathbb{H}/G \cong \mathbb{H} \times \Delta^* \rightarrow X_{d^2}$$

(where Δ^* is the punctured unit disk), which extends to an unramified map of orbifolds,

$$p: \mathbb{H} \times \Delta \rightarrow \widehat{X}_{d^2}$$

sending $\mathbb{H} \times \{0\}$ to $R_{d^2}^2$, and sending leaves of the vertical foliation by disks to leaves of $\widehat{\mathcal{A}}_{d^2}$. The vector field,

$$X = z_2 \frac{\partial}{\partial z_2},$$

on $\mathbb{H} \times \Delta^*$ locally defines a nonzero, holomorphic section of $T\widehat{\mathcal{A}}_{d^2}(-\overline{R}_{d^2}^2)$ around points in $R_{d^2}^2$. The norm of this section is

$$\rho(X, X) = (\log |z_2|)^{-2},$$

so by Lemma 9.9, ρ is good near points of $R_{d^2}^2$ (note that the exponent of the metric in Lemma 9.9 is irrelevant because changing the exponent only changes $\partial h/h$ by a constant).

The proof that ρ is good near the cusps of $\widehat{X}_D$ is the same, except these points are covered by $\Delta \times \Delta$, rather than $\mathbb{H} \times \Delta$. $\blacksquare$

Corollary 9.19. *The first Chern class of $T^*\mathcal{A}_{d^2}$ is*

$$c_1(T^*\mathcal{A}_{d^2}) = [\omega_2] - [\overline{S}_{d^2}^2] \quad (9.11)$$

Proof. By Theorem 9.18,

$$c_1(T^*\widehat{\mathcal{A}}_{d^2}) = [\omega_2] - [\overline{R}_{d^2}^2].$$

Since $\pi^*(T^*\widehat{\mathcal{A}}_{d^2})$ is isomorphic to $T^*\mathcal{A}_{d^2}$ over X_{d^2} ,

$$\pi^*(T^*\widehat{\mathcal{A}}_{d^2}) = T^*\mathcal{A}_{d^2} \left(\sum_P e_P C_P \right),$$

where the sum is over all nondegenerate Y_D -prototypes P . Thus

$$\sum_P e_P[C_P] = [\omega_2] - [\overline{S}_{d^2}^1] - c_1(T^*\mathcal{A}_{d^2}) \quad (9.12)$$

because $\pi^*[\overline{R}_{d^2}^2] = [\overline{S}_{d^2}^2]$.

The pairing of $[C_P]$ with the right hand side of (9.12) is trivial for all P by Corollary 9.14 together with the fact that from Theorem 8.7 that C_P is disjoint from $\overline{S}_{d^2}^2$ if P is nonterminal, and C_P intersects $\overline{S}_{d^2}^2$ in one transverse intersection if P is terminal. Theorem 8.11 then implies that the e_P are all zero. Thus $c_1(T^*\mathcal{A}_{d^2})$ is as claimed. $\blacksquare$

10 Euler characteristic of W_D

In this section, we calculate $\chi(W_D)$. We construct a meromorphic section of a line bundle over Y_D which vanishes along $\overline{W}_D$ and has simple poles along $\overline{P}_D$ and $\overline{S}_D^2$. This allows us to relate $\chi(W_D)$ to $\chi(P_D)$, $\chi(X_D)$, and $\chi(S_D^2)$.

In [McMb], McMullen defined a quadratic differential on the leaves of $\mathcal{A}_D$ in X_D , and used this to construct a transverse measure for the foliation of X_D by $\mathrm{SL}_2\mathbb{R}$ orbits. This quadratic differential was also studied in [Sch05] in the case when D is square. Here we recall the construction of this quadratic differential from [McMb] and define it on all of Y_D .

Theorem 10.1. *On each leaf L of $\mathcal{A}_D$, there is locally a quadratic differential q which has a simple zero on each point of $\overline{W}_D \cap L$, has a simple pole on each point of $(\overline{P}_D \cup \overline{S}_D^2) \cap L$, and is elsewhere holomorphic and nonzero.*

Proof. For each $z \in L$, let X_z be the corresponding Riemann surface with real multiplication. Choose a basepoint $z_0 \in L$, and choose some eigenform $\omega_{z_0} \in \Omega^1(X_{z_0})$. There is a unique section $z \mapsto \omega_z$ of ΩY_D over L such that the absolute periods of the ω_z are locally constant. When $L \subset X_D$, this follows from Proposition 2.10. If $L = C_P$, this is clear because the periods of ω_z are determined by the residues at the nonseparating nodes of X_z , and along C_P these residues have constant ratio, and if $L = S_{d^2}^1$, this is also clear because if (X_z, ω_z) is a cylinder covering differential, then the periods of ω_z are all rational multiples of the period around the node of X_z , so if this period is constant along L , then all of the periods are constant.

If $z \in L \setminus (\overline{W}_D \cup \overline{P}_D \cup \overline{S}_D^2)$, then (X_z, ω_z) has two distinct simple zeros. Let

$$f(z) = \int_{\gamma} \omega_z, \quad (10.1)$$

where γ is some path joining the zeros of ω_z . This $f(z)$ is a multivalued holomorphic function on $L \setminus (\overline{W}_D \cup \overline{P}_D \cup \overline{S}_D^2)$ because the value of $f(z)$ depends on a choice of an oriented path joining the zeros of ω_z . Define

$$q = (\partial f)^2.$$

We claim that q is a well-defined quadratic differential on L . To see this, suppose we replace an oriented path γ joining the zeros p and q of ω_z with a new path γ' joining p to q . Since $\gamma - \gamma'$ is a closed path, this changes f by an absolute period of ω_z . Since the absolute periods of ω_z are constant along L , replacing γ with γ' does not change ∂f . If we replace γ with $-\gamma$, the same path with the opposite orientation, this changes f to $-f$, which does not affect q because of the exponent. Thus q is well defined.

We now identify the zeros and poles of q . Since the absolute and relative periods give a system of local coordinates on the strata in $\Omega\overline{\mathcal{M}}_2$, the relative periods give local coordinates on L because the absolute periods are constant, so q is holomorphic and nonzero on the complement of $\overline{P}_D$, $\overline{W}_D$, and $\overline{S}_D^2$.

Suppose $z \in L \cap \overline{W}_D$. For $z \in \mathbb{C}$, let $I(z) \in \mathbb{C}$ be the segment joining 0 to z . For $w \in \Delta_\epsilon$, where Δ_ϵ is some small ϵ -ball around 0 in $\mathbb{C}$, let

$$\phi(w) = (X_w, \omega_w) \#_{I(w^{3/2})},$$

(this is the operation of splitting a double zero defined in §4.1). As in §6.2, ϕ is a conformal isomorphism onto some neighborhood of z in $\mathbb{C}$. In these coordinate, we have $f(w) = w^{3/2}$, so

$$q = \frac{9}{4}w dw^2.$$

Thus q has a simple zero at z .

Now suppose $z \in L \cap \overline{P}_D$. Then

$$(X_z, \omega_z) = (X_1, \omega_1) \# (X_2, \omega_2),$$

the one-point connected sum of two genus one differentials. Define a conformal mapping $\phi: \Delta_\epsilon \rightarrow L$ by

$$\phi(w) = (X_1, \omega_1) \#_{I(w^{1/2})} (X_2, \omega_2),$$

taking a connected sum along $I(w^{1/2})$. Similarly, if $z \in L \cap \overline{S}_D^2$, then we can regard (X_z, ω_z) as an genus one differential with two points identified to a node. Define a conformal mapping $\phi: \Delta_\epsilon \rightarrow L$ by

$$\phi(w) = (X_z, \omega_z) \#_{I(w^{1/2})},$$

taking a self-connected sum along $I(w^{1/2})$ as described in §4.1. In either case, we have in these coordinates $f(w) = w^{1/2}$, so

$$q = \frac{1}{2}w^{-1}dw^2.$$

Thus q has a simple pole at z . ■

This construction locally defines meromorphic sections of $(T^*\mathcal{A}_D)^2$, but does not give a global section. The problem is that the definition of the quadratic differential q on a leaf L depended on a choice of $\omega_{z_0} \in \Omega(X_{z_0})$ for some base-point $z_0 \in L$. There is no obvious way to choose these quadratic differentials coherently to get a global section. To get a global section of a bundle, we must twist by some power of ΩY_D .

Theorem 10.2. *There is a meromorphic section of the line bundle*

$$\mathcal{L} = (QY_D)^* \otimes (T^*\mathcal{A}_D)^2$$

on Y_D which has a simple zero along $\overline{W}_D$, has a simple pole along $\overline{P}_D$ and $\overline{S}_D^2$ (which is empty if D is nonsquare), and is elsewhere nonzero and finite.

Proof. The construction in the proof of Theorem 10.1 defined for each point,

$$z \in Y_D \setminus \bigcup c_P,$$

together with a choice of $\omega \in \Omega^1(X_z)$ a quadratic differential q on the leaf of $\mathcal{A}_D$ through z . Thus, we have a map $h: \Omega^1(X_z) \rightarrow (T^*\mathcal{A}_D)^2|_z$. With f as in (10.1), if we replace $\omega \in \Omega^1(X_z)$ with $a\omega$, then f becomes af , and q becomes a^2q . Thus h satisfies,

$$h(a\omega) = a^2h(\omega).$$

We can thus regard h as a linear map,

$$\Omega^1(X_z)^{\otimes 2} \rightarrow (T^*\mathcal{A}_D)|_z.$$

This defines a meromorphic section s of $\mathcal{L}$ over $Y_D \setminus \bigcup c_P$ which has the same zeros and poles as the quadratic differentials on the leaves of $\mathcal{A}_D$. This section extends holomorphically over the points c_P to give a section over all of Y_D because any holomorphic section of a line bundle defined on a neighborhood of a normal point extends over that point. ■

Corollary 10.3. *If D is not square, then the fundamental class $[\overline{W}_D]$ of $\overline{W}_D$ in $H^2(Y_D; \mathbb{Q})$ is given by*

$$[\overline{W}_D] = [\overline{P}_D] - [\omega_1] + 2[\omega_2]. \quad (10.2)$$

The fundamental class $[\overline{W}_{d^2}]$ of $\overline{W}_{d^2}$ in $H^2(Y_{d^2}; \mathbb{Q})$ is given by

$$[\overline{W}_{d^2}] = [\overline{P}_{d^2}] - [\overline{S}_{d^2}^2] - [\omega_1] + 2[\omega_2]. \quad (10.3)$$

Proof. By [GH78, p. 141], for any line bundle L over a compact, complex manifold M with a meromorphic section s of L having divisor D ,

$$c_1(L) = [D],$$

This is proved for complex manifolds, but this is still true and the proof works just as well for orbifolds.

In our situation, the section s of $\mathcal{L}$ from Theorem 10.2 implies

$$[\overline{W}_D] - [\overline{P}_D] - [\overline{S}_D^2] = c_1(\mathcal{L}) = -c_1(QY_D) + 2c_1(T^*\mathcal{A}_D).$$

This, together with (9.9), (9.10), and (9.11) yield the desired formulas. ■

Corollary 10.4. *If $D \neq 1$ is a fundamental discriminant, then*

$$\chi(W_{f^2D}) = \chi(P_{f^2D}) - 2\chi(X_{f^2D}) \quad (10.4)$$

$$= -9\zeta_{K_D}(-1)f^3 \sum_{r|f} \left(\frac{D}{r}\right) \frac{\mu(r)}{r^2}. \quad (10.5)$$

If $D = d^2$, then

$$\chi(W_{d^2}) = \chi(P_{d^2}) - \chi(S_{d^2}^2) - 2\chi(X_{d^2}) \quad (10.6)$$

$$= -\frac{1}{16}d^2(d-2) \sum_{r|d} \frac{\mu(r)}{r^2}. \quad (10.7)$$

Proof. If we pair $-\omega_1$ with both sides of (10.2) and (10.3), then by Propositions 9.5 and 9.6, we get

$$-\int_{W_D} \omega_1 = -\int_{P_D} \omega_1 - 2 \int_{X_D} \omega_1 \wedge \omega_2, \quad \text{and} \quad (10.8)$$

$$-\int_{W_{d^2}} \omega_1 = -\int_{P_{d^2}} \omega_1 - 2 \int_{X_{d^2}} \omega_1 \wedge \omega_2 + \int_{S_{d^2}^2} \omega_1. \quad (10.9)$$

Since W_D , P_D , and $S_{d^2}^2$ are transverse to $\mathcal{A}_D$, the form $-\omega_1$ restricts to the Chern form of the hyperbolic metric on these Riemann surfaces. Thus by the Gauss-Bonnet theorem we get (10.4) and (10.6). We calculated $\chi(X_D)$ and $\chi(P_D)$ in Theorems 2.12 and 2.22. We will calculate $\chi(S_{d^2}^2)$ in the following proposition. Putting all of this together yields (10.5) and (10.7). ■

Proposition 10.5. *We have,*

$$\chi(S_4^i) = -\frac{1}{2}, \quad (10.10)$$

and

$$\chi(S_{d^2}^i) = -\frac{1}{12}d^2 \sum_{r|d} \frac{\mu(r)}{r^2} \quad (10.11)$$

when $d > 2$.

Proof. We know that

$$S_{d^2}^i \cong \mathbb{H}/\Gamma_1(d).$$

It follows from [Miy89, Theorem 4.2.5] that,

$$\chi(\mathbb{H}/\Gamma_1(d)) = \begin{cases} -\frac{1}{2} & \text{if } d = 2; \\ -\frac{1}{12}\phi(d)d \prod_{p|d} \left(1 + \frac{1}{p}\right) & \text{when } d > 2, \end{cases}$$

where ϕ is the Euler ϕ -function, and the product is over all primes dividing d . By [IR90, Proposition 2.2.5],

$$\phi(d) = d \prod_{p|d} \left(1 - \frac{1}{p}\right),$$

so when $d > 2$,

$$\chi(\mathbb{H}/\Gamma_1(d)) = -\frac{1}{12}d^2 \prod_{p|d} \left(1 - \frac{1}{p^2}\right) = -\frac{1}{12}d^2 \sum_{r|d} \frac{\mu(r)}{r^2},$$

as claimed. ■

11 Fundamental class of $\overline{W}_D$

In this section, we will calculate the fundamental class of $\overline{W}_D$. By Corollary 10.3, we just need to know the fundamental classes of $\overline{S}_D^i$ and $\overline{P}_D$.

Theorem 11.1. *The fundamental class of $\overline{S}_{d^2}^i$ in $H^2(Y_{d^2}; \mathbb{Q})$ is*

$$[\overline{S}_{d^2}^i] = \frac{6}{d}[\omega_i] + \pi_B[\overline{S}_{d^2}^i] \quad (11.1)$$

Proof. From Theorem 8.10, we have a map $p: Y_{d^2} \rightarrow \widehat{X}_{d^2}$ which collapses the curves C_P to cusps of $\widehat{X}_{d^2}$. Let

$$q: \widehat{X}_{d^2} \rightarrow \overline{\mathbb{H}/\mathrm{SL}_2\mathbb{Z}} = \overline{\mathcal{M}}_{1,1}$$

be the map induced by the projection of $\mathbb{H} \times \mathbb{H}$, the universal cover of X_{d^2} , onto its second factor. Let $r = q \circ p$, and let $\infty \in \overline{\mathcal{M}}_{1,1}$ be the single point added to $\mathbb{H}/\mathrm{SL}_2\mathbb{Z}$. Then

$$r^{-1}(\infty) = S_{d^2}^1 \cup \bigcup_P C_P.$$

Let f be a holomorphic function defined on a neighborhood of ∞ in $\overline{\mathcal{M}}_{1,1}$ which has a simple zero at ∞ . We claim that $f \circ r$ vanishes to order d along $S_{d^2}^1$.

It is enough to show that $f \circ q$ vanishes to order d along $R_{d^2}^1$. Let $G \subset \mathrm{SL}_2\mathcal{O}_{d^2}$ be the cyclic group generated by

$$\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & d \\ 0 & 1 \end{pmatrix} \right),$$

and let $N \subset \mathrm{SL}_2\mathbb{Z}$ be the subgroup of upper-triangular matrices. We have the following commutative diagram:

$$\begin{array}{ccc} \mathbb{H} \times \mathbb{H}/G & \longrightarrow & \mathbb{H}/N \\ \downarrow & & \downarrow \\ \mathbb{H} \times \Delta & \xrightarrow{t} & \Delta \\ \downarrow i & & \downarrow j \\ \widehat{X}_{d^2} & \xrightarrow{q} & \overline{\mathcal{M}}_{1,1} \end{array}$$

where $\Delta \subset \mathbb{C}$ is the unit disk. The two topmost vertical maps are isomorphisms onto $\mathbb{H} \times \Delta^*$ and Δ^* respectively, where Δ^* is the punctured disk. The maps i and j are unramified maps of orbifolds satisfying $i^{-1}(R_{d^2}^1) = \mathbb{H} \times \{0\}$ and $j^{-1}(\infty) = \{0\}$. The map t is given by $t(z, w) = w^d$, and it follows that $f \circ q$ vanishes to order d along $R_{d^2}^1$ as claimed because $f \circ j \circ t$ vanishes to order d along $\mathbb{H} \times \{0\}$.

It follows from this claim that

$$r^*[\infty] = d[\overline{S}_{d^2}^1] + \sum_P e_P[C_P], \quad (11.2)$$

where e_P is the order of vanishing of $f \circ r$ along C_P . Let η be the 2-form on $\mathbb{H}/\mathrm{SL}_2\mathbb{Z}$ induced by

$$\frac{1}{2\pi} \frac{dx \wedge dy}{y^2}$$

on $\mathbb{H}$. Since $\chi(\mathbb{H}/\mathrm{SL}_2\mathbb{Z}) = -1/6$,

$$\int_{\overline{\mathcal{M}}_{1,1}} \eta = \frac{1}{6},$$

by the Gauss-Bonnet Theorem, so the cohomology class defined by the closed current $\langle \eta \rangle$ on $\overline{\mathcal{M}}_{1,1}$ satisfies

$$6[\eta] = [\infty]$$

in $H^2(\overline{\mathcal{M}}_{1,1}; \mathbb{Q})$. We have

$$r^*\eta = \omega_1,$$

so

$$r^*[\infty] = 6[\omega_1]. \quad (11.3)$$

Equations (11.2) and (11.3) imply (11.1) for $i = 1$. The case $i = 2$ follows from the same argument or by applying τ^* to (11.1). $\blacksquare$

Theorem 11.2. *If D is not square, then the fundamental class of $\overline{P}_D$ in $H^2(Y_D; \mathbb{Q})$ is*

$$[\overline{P}_D] = \frac{5}{2}([\omega_1] + [\omega_2]) + \pi_B[\overline{P}_D]. \quad (11.4)$$

The fundamental class of $\overline{P}_{d^2}$ in $H^2(Y_{d^2}; \mathbb{Q})$ is

$$[\overline{P}_{d^2}] = \left(\frac{5}{2} - \frac{3}{d} \right) ([\omega_1] + [\omega_2]) + \pi_B[\overline{P}_{d^2}] \quad (11.5)$$

Proof. In $\overline{\mathcal{M}}_2$, let Δ_0 be the divisor which is the closure of the locus of stable Riemann surfaces with one nonseparating node, and let Δ_1 be the divisor which is the closure of the locus of stable Riemann surfaces with one separating node. Let $\delta_i = [\Delta_i]$, the fundamental class of Δ_i . Define

$$\lambda_1 = c_1(\Omega \overline{\mathcal{M}}_2).$$

These cohomology classes satisfy the well-known relation,

$$\delta_1 = 5\lambda_1 - \frac{1}{2}\delta_0, \quad (11.6)$$

proved in [Mum83].

Since $\pi_2^*(\Omega\overline{\mathcal{M}}_2) = \Omega^1 Y_D \oplus \Omega^2 Y_D$, where $\pi_2: Y_D \rightarrow \overline{\mathcal{M}}_2$ is the natural map, we have

$$\pi_2^*(\lambda_1) = \frac{1}{2}([\omega_1] + [\omega_2]) \quad (11.7)$$

by Corollary 9.10.

We claim that

$$\pi_2^*(\delta_1) = [\overline{P}_D]. \quad (11.8)$$

Since $\pi_2^{-1}(\Delta_1) = \overline{P}_D$, it suffices to show that Δ_1 is generically transverse to $\pi_2(Y_D)$. In $\mathcal{A}_2$, the divisor Δ_1 corresponds to the surface X_1 parameterizing products of elliptic curves. In $\mathcal{A}_2$, the intersection of X_1 with X_D is transverse because these are both linear subspaces of $\mathcal{A}_2$, so if their intersection was not transverse, one would be contained in the other. Equation (11.8) follows.

We now claim that

$$\pi_2^*(\delta_0) = [\overline{S}_D^1] + [\overline{S}_D^2] + \sum_P e_P [C_P] \quad (11.9)$$

for some integers e_P . Since

$$\pi_2^{-1}(\Delta_0) = \bigcup_{i=1}^2 \overline{S}_D^i \cup \bigcup_P C_P,$$

$\pi_2 \circ \tau = \pi_2$, and $\tau^* S_D^1 = S_D^2$, it suffices to show that Δ_0 meets $\pi_2(Y_D)$ transversely along $\pi_2(S_D^1)$.

Let $p \in S_{d^2}^1$, and let $(X, \omega) \in \Omega\overline{\mathcal{M}}_2$ be a corresponding eigenform. We claim that $\pi_2(Y_D)$ meets Δ_0 transversely at $\pi_2(p)$. Let $\{\alpha_i, \beta_i\}$ be a symplectic basis for $H_1(X; \mathbb{Z})$ as in §6.3. Then we get coordinates (v, w, x, y, z) on a neighborhood U of (X, ω) in $\Omega\overline{\mathcal{M}}_2$ as in §6.3. The subspace H of U defined by the equations,

$$v = \omega(\alpha_1) \quad w = \omega(\alpha_2),$$

maps locally biholomorphically to $\overline{\mathcal{M}}_2$, so the coordinates (x, y, z) on H induce coordinates on $\overline{\mathcal{M}}_2$ on a neighborhood W of X . In these coordinates,

$$\Delta_0 = V(z).$$

By Corollary 7.7,

$$\pi_2(Y_D) \cap W = V(x - \omega(\beta_2)).$$

Thus $\pi_2(Y_D)$ and Δ_0 meet transversely at p as claimed, and (11.9) follows.

Equations (11.4) and (11.5) then follow from (11.6), (11.7), (11.8), and (11.9). ■

Remark. I am grateful to Gerard van der Geer for providing the idea of the proof of this theorem.

Remark. We can also use (11.4) and (11.5) to get a new proof of Theorem 2.22. Since this proof doesn't use Siegel's formula, Theorem 2.16, this together with the previous proof of Theorem 2.22 can be used to give a proof of Siegel's formula.

Corollary 11.3. *If D is not square, then the fundamental class of $\overline{W}_D$ in $H^2(Y_D; \mathbb{Q})$ is given by*

$$[\overline{W}_D] = \frac{3}{2}[\omega_1] + \frac{9}{2}[\omega_2] + \pi_B[\overline{W}_D]. \quad (11.10)$$

The fundamental class of $\overline{W}_{d^2}$ in $H^2(Y_{d^2}; \mathbb{Q})$ is given by

$$[\overline{W}_{d^2}] = \frac{3}{2} \left(1 - \frac{2}{d}\right) [\omega_1] + \frac{9}{2} \left(1 - \frac{2}{d}\right) [\omega_2] + \pi_B[\overline{W}_{d^2}]. \quad (11.11)$$

Proof. This follows from plugging the formulas from Theorems 11.1 and 11.2 into the formulas from Corollary 10.3. ■

12 Normal Bundles

We now study the normal bundles of the curves $\overline{W}_D$, $\overline{P}_D$, and $\overline{S}_D^2$ with the goal of calculating the self-intersection numbers of these curves. For any curve $C \subset Y_D$, we will write $N(C)$ for its normal bundle.

Proposition 12.1. *For any connected component C of $\overline{W}_D$, $\overline{P}_D$, or $\overline{S}_D^2$,*

$$N(C) \cong T\mathcal{A}_D|_C \quad (12.1)$$

as holomorphic line bundles over C .

Proof. We claim that the foliation $\mathcal{A}_D$ is transverse to $\overline{W}_D$, $\overline{P}_D$, and $\overline{S}_D^2$. Equation (12.1) follows directly from this claim.

The curves W_D and P_D are transverse to $\mathcal{A}_D$ because the inverse images of these curves in $\mathbb{H} \times \mathbb{H}$ are the unions of graphs of holomorphic functions $\mathbb{H} \rightarrow \mathbb{H}$. The curve S_D^2 is also transverse to $\mathcal{A}_D$ by Proposition 9.12. The closures of these curves are then transverse to $\mathcal{A}_D$ because they intersect $Y_D \setminus X_D$ in the curves C_P , these intersections are transverse by Theorem 8.7, and the curves C_P are leaves of $\mathcal{A}_D$ by Proposition 9.12. ■

Theorem 12.2. *For any connected component C of P_D or S_D^2 ,*

$$[\overline{C}]^2 = \chi(C).$$

For any component C of W_D ,

$$[\overline{C}]^2 = \frac{1}{3}\chi(C).$$

Proof. Let C be a connected component of P_D , S_D^2 , or W_D , and choose a tubular neighborhood $U \subset Y_D$ of $\overline{C}$ which is small enough that each point in $U \setminus \overline{C}$ represents a stable Abelian differential which has a unique shortest saddle connection joining distinct zeros. We wish to define a map,

$$\Phi: U \rightarrow (Q^1\overline{C})^*,$$

where $Q^1\overline{C}$ is the restriction of Q^1Y_D to $\overline{C}$.

Consider $p \in U$ representing the projective class of a stable Abelian differential $(X, [\omega])$. Let $I \subset X$ be the unique shortest saddle connection connecting distinct zeros, and let (Y, η) be the stable Abelian differential obtained by collapsing I as in §4.1. The projective class $(Y, [\eta])$ then represents a point of $\overline{C}$. Since ω and η are both eigenforms for real multiplication, we have an isomorphism,

$$T: \Omega^1(Y) \rightarrow \Omega^1(X),$$

defined by $T(\eta) = \omega$. Define $S \in Q^1(Y)^* = \Omega^1(Y)^{-2}$ by

$$S(\nu) = \left(\int_I T(\sqrt{\nu}) \right)^2,$$

where the integral along I is with respect to some choice of orientation of I . It doesn't matter which orientation we take for I or which square root of ν we take, so S is well-defined. Now define Φ by

$$\Phi(p) = ((Y, [\eta]), S).$$

For $p \in \overline{C}$, we define $\Phi(p) = (p, 0)$. Note that Φ takes leaves of $\mathcal{A}_D$ to fibers of $(Q^1\overline{C})^*$.

Suppose that C is a connected component of P_D . We claim that in this case, Φ is injective. To see this, let $q \in (Q^1\overline{C})^*$ be represented by $(Y, [\eta]) \in \mathbb{P}\Omega\overline{\mathcal{M}}_2$ and $S \in Q^1(Y)^*$. This (Y, η) is the one point union of two genus one differentials or cylinders:

$$(Y, \eta) = (Y_1, \eta_1) \# (Y_2, \eta_2).$$

Normalize η so that $S(\eta^2) = 1$. If $\Phi(p) = q$, then p is represented by the connected sum

$$(X, \omega) = (Y_1, \eta_1) \#_I (Y_2, \eta_2),$$

where $I \subset \mathbb{C}$ is the segment joining 0 to 1. Thus p is determined uniquely by q , so Φ is injective as claimed. When $C = S_D^2$, then Φ is injective by the same argument, using a self-connected sum in place of the connected sum operation above.

Recall that in §4.1, we defined the operation of splitting a double zero, which is inverse to the operation of collapsing a saddle connection, and associates to a sufficiently small segment $I \subset (Y, \eta)$ starting at the zero of η the Abelian differential $(Y, \eta) \#_I$.

Suppose now that C is a connected component of W_D . We claim that in this case, Φ is branched of order three along $\overline{C}$. Again, let $q \in (Q^1\overline{C})^*$ be represented by $(Y, \eta) \in \mathbb{P}\Omega\overline{\mathcal{M}}_2$ and $S \in Q^1(Y)^*$, and normalize η so that $S(\eta^2) = 1$. This (Y, η) is a stable Abelian differential with a double zero z . If $\Phi(p) = q$, then p is represented by

$$(X, \omega) = (Y, \eta) \#_I$$

for some oriented segment I starting at z such that

$$\int_I \eta = 1. \tag{12.2}$$

There are at most three such segments because there are three positively oriented horizontal directions at the zero of η ; therefore, q has at most three preimages. If S is small, then there is an embedded ball around z with large radius. This means that there are three embedded segments starting at z satisfying (12.2), and we can split along each of these segments. Thus any point in $Q^1\overline{C}$ sufficiently close to the zero section has exactly three preimages under Φ , and the claim follows.

Now if C is a connected component of P_D or S_D^2 , then we have seen that U is homeomorphic to a neighborhood of the zero section in $Q^1\overline{C}$. Since U is a

tubular neighborhood of $\overline{C}$, U is also homeomorphic to a neighborhood of the zero section in $N(\overline{C})$. Therefore,

$$[\overline{C}]^2 = \deg N(\overline{C}) = \deg Q^1\overline{C}.$$

Since $c_1(Q^1Y_D) = [\omega_1]$ by Corollary 9.10,

$$[\overline{C}]^2 = \deg Q^1\overline{C} = - \int_C \omega_1 = \chi(C),$$

as claimed.

Now suppose C is a connected component of W_D . Regard U as a neighborhood of the zero section in $N(\overline{C})$. By the above claim, we can choose U so that $\Phi: U \rightarrow (Q^1\overline{C})^*$ preserves fibers and is exactly three-to-one onto its image. For any line bundle $B \rightarrow \overline{C}$, let $\tau_B \in H^2(B, B \setminus \overline{C}; \mathbb{R})$ be its Thom class. The Thom class $\tau_{(Q^1\overline{C})^*}$ is represented by a 2-form which is supported in $\Phi(U)$ which satisfies

$$\int_F \tau_{(Q^1\overline{C})^*} = 1$$

for each fiber F of $(Q^1\overline{C})^*$. For each fiber F of $N(\overline{C})$, we have

$$\int_F \Phi^* \tau_{(Q^1\overline{C})^*} = 3,$$

thus

$$\Phi^* \tau_{(Q^1\overline{C})^*} = 3\tau_{N(\overline{C})}.$$

It follows that

$$\begin{aligned} [\overline{C}]^2 &= \tau_{N(\overline{C})} \cdot [\overline{C}] \\ &= \frac{1}{3} \tau_{(Q^1\overline{C})^*} \cdot [\overline{C}] \\ &= \frac{1}{3} \deg(Q^1\overline{C})^* \\ &= \frac{1}{3} \chi(C), \end{aligned}$$

as claimed. ■

Corollary 12.3. *For any connected component C of W_D , we have*

$$\int_C \omega_2 = \frac{1}{3} \int_C \omega_1.$$

Proof. By Proposition 12.1, we have

$$[C]^2 = \deg T\mathcal{A}_D|_C = c_1(T\mathcal{A}_D) \cdot [C] = - \int_C \omega_2.$$

By Theorem 12.2,

$$[C]^2 = -\frac{1}{3} \int_C \omega_1,$$

and the claim follows. ■

13 Euler characteristic of W_D^ϵ

Cohomology of Y_D . We saw in Theorem 8.11 that the intersection pairing on $H^2(Y_D; \mathbb{Q})$ is negative definite on the subspace B generated by the fundamental classes of the curves C_P . By Corollary 9.11, the subspace $\langle [\omega_1], [\omega_2] \rangle \subset H^2(Y_D; \mathbb{Q})$ is orthogonal to B . Thus, if we let $J \subset H^2(Y_D; \mathbb{Q})$ be the orthogonal complement to $B \oplus \langle [\omega_1], [\omega_2] \rangle$, then we have the orthogonal direct sum,

$$H^2(Y_D; \mathbb{Q}) = B \oplus \langle [\omega_1], [\omega_2] \rangle \oplus J. \quad (13.1)$$

Since

$$B \oplus \langle [\omega_1], [\omega_2] \rangle \subset H^{1,1}(Y_D; \mathbb{Q}),$$

J contains all of $H^{2,0}(Y_D; \mathbb{Q})$ and $H^{0,2}(Y_D; \mathbb{Q})$.

$\chi(W_D^\epsilon)$ when D is not square. We now calculate $\chi(W_D^\epsilon)$ when D is not square. Until further notice, we will assume that D is not square.

According to Corollary 8.12,

$$\pi_B[\overline{W}_D] = \pi_B[\overline{P}_D].$$

Let $B_D = \pi_B[\overline{P}_D]$, and let $B_D^\epsilon = \pi_B[\overline{W}_D^\epsilon]$ for $\epsilon = 1, 2$. Since

$$[\overline{W}_D] = [\overline{W}_D^0] + [\overline{W}_D^1],$$

we have

$$B_D^0 + B_D^1 = B_D.$$

Lemma 13.1. *For any nonsquare D , we have*

$$(B_D)^2 = -15\chi(X_D). \quad (13.2)$$

Proof. Since W_D and P_D are disjoint,

$$[\overline{W}_D] \cdot [\overline{P}_D] = 0.$$

Equation (13.2) follows directly from this together with the equations (11.4) and (11.10) for these fundamental classes. $\blacksquare$

Theorem 13.2. *If D is not square, then the fundamental class of $\overline{W}_D^\epsilon$ in $H^2(Y_D; \mathbb{Q})$ is*

$$[\overline{W}_D^\epsilon] = \frac{3}{4}[\omega_1] + \frac{9}{4}[\omega_2] + B_D^\epsilon + j \quad (13.3)$$

for some $j \in J$.

Proof. Since W_D^0 and P_D are disjoint, we have

$$[\overline{W}_D^0] \cdot [P_D] = 0, \quad (13.4)$$

and by (8.6), we have

$$(B_D^0)^2 = (B_D^1)^2. \quad (13.5)$$

By (13.1) and Corollary 12.3, the fundamental classes of the $\overline{W}_D^\epsilon$ are of the form,

$$\begin{aligned} [\overline{W}_D^0] &= a[\omega_1] + 3a[\omega_2] + B_D^0 + j \\ [\overline{W}_D^1] &= \left(\frac{3}{2} - a\right)[\omega_1] + \left(\frac{9}{2} - 3a\right)[\omega_2] + B_D^1 - j, \end{aligned}$$

for some $a \in \mathbb{Q}$ and $j \in J$. In terms of a , (13.4) becomes

$$10a\chi(X_D) + (B_D^0)^2 + B_D^0 \cdot B_D^1 = 0. \quad (13.6)$$

From (13.2) and (13.5), we obtain

$$(B_D^0)^2 + B_D^0 \cdot B_D^1 = \frac{1}{2}(B_D^0 + B_D^1)^2 = \frac{1}{2}(B_D)^2 = -\frac{15}{2}\chi(X_D). \quad (13.7)$$

Combining (13.6) and (13.7) yields

$$a = \frac{3}{4},$$

as desired. ■

Remark. It seems likely that $j = 0$, but we don't know how to prove this.

Corollary 13.3. *If D is not square, then*

$$\chi(W_D^0) = \chi(W_D^1).$$

$\chi(W_D^\epsilon)$ when D is square. We now turn to the calculation of $\chi(W_{d^2}^\epsilon)$. The idea is the same as the proof of Theorem 13.2, but the calculation is more complicated because of the presence of the curves $S_{d^2}^i$. We will restrict to the case $d > 2$ because $W_4 = \emptyset$. We start by calculating the intersections of various classes in $H^2(Y_{d^2}; \mathbb{Q})$.

Lemma 13.4. *For any $d > 2$, we have the following intersection numbers:*

$$[\omega_1] \cdot [\omega_2] = \frac{1}{72}d^3 \sum_{r|d} \frac{\mu(r)}{r^2} \quad (13.8)$$

$$\pi_B[\overline{S}_{d^2}^i] \cdot \pi_B[\overline{P}_{d^2}] = \left(-\frac{5}{24}d^2 + \frac{1}{4}d\right) \sum_{r|d} \frac{\mu(r)}{r^2} \quad (13.9)$$

$$(\pi_B[\overline{S}_{d^2}^i])^2 = -\frac{1}{12}d^2 \sum_{r|d} \frac{\mu(r)}{r^2} \quad (13.10)$$

$$\pi_B[\overline{S}_{d^2}^1] \cdot \pi_B[\overline{S}_{d^2}^2] = -\frac{1}{2}d \sum_{r|d} \frac{\mu(r)}{r^2} + \frac{1}{2}\phi(d) \quad (13.11)$$

$$\pi_B[\overline{S}_{d^2}^2] \cdot \pi_B[\overline{W}_{d^2}] = \left(-\frac{1}{8}d^2 + \frac{1}{4}d\right) \sum_{r|d} \frac{\mu(r)}{r^2} \quad (13.12)$$

$$(\pi_B[\overline{W}_{d^2}])^2 = \left(-\frac{5}{24}d^3 + \frac{19}{24}d^2 - \frac{3}{4}d\right) \sum_{r|d} \frac{\mu(r)}{r^2} \quad (13.13)$$

$$(\pi_B[\overline{P}_{d^2}])^2 = \left(-\frac{5}{24}d^3 + \frac{11}{24}d^2 - \frac{1}{4}d\right) \sum_{r|d} \frac{\mu(r)}{r^2} \quad (13.14)$$

$$\pi_B[\overline{S}_{d^2}^1] \cdot \pi_B[\overline{W}_{d^2}] = \left(-\frac{5}{24}d^2 + \frac{3}{4}d\right) \sum_{r|d} \frac{\mu(r)}{r^2} - \frac{1}{2}\phi(d) \quad (13.15)$$

Proof. Equation (13.8) is $\chi(X_{d^2})$, which is given in Theorem 2.12.

By Theorem 8.7, $\overline{P}_{d^2} \cap \overline{S}_{d^2}^i = \emptyset$. Thus $[\overline{P}_{d^2}] \cdot [\overline{S}_{d^2}^i] = 0$, from which (13.9) follows.

By Theorem 12.2 and (10.11),

$$[\overline{S}_{d^2}^i]^2 = \chi(S_{d^2}^i) = -\frac{1}{12}d^2 \sum_{r|d} \frac{\mu(r)}{r^2},$$

from which (13.10) follows.

By Theorem 8.7,

$$[\overline{S}_{d^2}^1] \cdot [\overline{S}_{d^2}^2] = -\frac{1}{2}\phi(d).$$

Equation (13.11) follows.

By Theorem 8.7,

$$[\overline{S}_{d^2}^2] \cdot [\overline{W}_{d^2}] = 0.$$

Equation (13.12) follows.

By Theorem 12.2 and (10.6),

$$[\overline{W}_{d^2}]^2 = \frac{1}{3}\chi(W_{d^2}) = -\frac{1}{48}d^2(d-2) \sum_{r|d} \frac{\mu(r)}{r^2},$$

from which (13.13) follows. Equation (13.14) is proved similarly.

By Corollary 8.12,

$$\pi_B[\overline{S}_{d^2}^1] \cdot \pi_B[\overline{W}_{d^2}] = \pi_B[\overline{S}_{d^2}^1] \cdot \pi_B[\overline{P}_{d^2}] - \pi_B[\overline{S}_{d^2}^1] \cdot \pi_B[\overline{S}_{d^2}^2].$$

Then (13.15) follows from (13.9) and (13.11). ■

Theorem 13.5. *For any $d > 2$, the fundamental class of $\overline{W}_{d^2}^\epsilon$ in $H^2(Y_{d^2}; \mathbb{Q})$ is given by*

$$\begin{aligned} [\overline{W}_{d^2}^0] &= \frac{3}{4} \left(1 - \frac{1}{d}\right) [\omega_1] + \frac{9}{4} \left(1 - \frac{1}{d}\right) [\omega_2] + \pi_B[\overline{W}_{d^2}^0] + j \\ [\overline{W}_{d^2}^1] &= \frac{3}{4} \left(1 - \frac{3}{d}\right) [\omega_1] + \frac{9}{4} \left(1 - \frac{3}{d}\right) [\omega_2] + \pi_B[\overline{W}_{d^2}^1] - j. \end{aligned}$$

for some $j \in J$.

Proof. By (13.1), Corollary 11.3, and Corollary 12.3, the fundamental classes of the $\overline{W}_{d^2}^\epsilon$ are given by

$$\begin{aligned} [\overline{W}_{d^2}^0] &= a[\omega_1] + 3a[\omega_2] + \pi_B[\overline{W}_{d^2}^0] + j \\ [\overline{W}_{d^2}^1] &= \left(\frac{3}{2} - \frac{3}{d} - a\right)[\omega_1] + \left(\frac{9}{2} - \frac{9}{d} - 3a\right)[\omega_2] + \pi_B[\overline{W}_{d^2}^1] - j \end{aligned}$$

for some $a \in \mathbb{Q}$ and $j \in J$.

From Corollary 8.16, we have

$$\begin{aligned} (\pi_B[\overline{W}_{d^2}^0])^2 &= (\pi_B[\overline{W}_{d^2}^1] + \pi_B[\overline{S}_{d^2}^2])^2 \\ &= (\pi_B[\overline{W}_{d^2}^1] - \pi_B[\overline{W}_{d^2}^0] + \pi_B[\overline{S}_{d^2}^2])^2. \end{aligned}$$

Using (13.10), (13.12), and (13.13), this simplifies to

$$\begin{aligned} \pi_B[\overline{W}_{d^2}] \cdot \pi_B[\overline{W}_{d^2}^0] + \pi_B[\overline{W}_{d^2}^0] \cdot \pi_B[\overline{S}_{d^2}^2] \\ = \left(-\frac{5}{48}d^3 + \frac{11}{48}d^2 - \frac{1}{8}d\right) \sum_{r|d} \frac{\mu(r)}{r^2}. \end{aligned} \quad (13.16)$$

From Corollary 8.12, we have

$$\pi_B[W_{d^2}] + \pi_B[S_{d^2}^2] = \pi_B[P_{d^2}]. \quad (13.17)$$

Multiplying (13.17) by $\pi_B[\overline{W}_{d^2}^0]$ and subtracting the result from (13.16), we obtain

$$\pi_B[\overline{W}_{d^2}^0] \cdot \pi_B[\overline{P}_{d^2}] = \left(-\frac{5}{48}d^3 + \frac{11}{48}d^2 - \frac{1}{8}d\right) \sum_{r|d} \frac{\mu(r)}{r^2}. \quad (13.18)$$

Since $\overline{W}_{d^2}^0$ and $\overline{P}_{d^2}$ are disjoint, we have

$$[\overline{W}_{d^2}^0] \cdot [\overline{P}_{d^2}] = 0.$$

Expanding, this becomes

$$\left(\frac{5}{32}d^3 - \frac{1}{6}d^2\right) a \sum_{r|d} \frac{\mu(r)}{r^2} + \pi_B[\overline{W}_{d^2}^0] \cdot \pi_B[\overline{P}_{d^2}] = 0,$$

which with (13.18) yields

$$a = \frac{3}{4} \left(1 - \frac{1}{d}\right).$$

■

Corollary 13.6. *For any $d > 2$ with $d \equiv 1 \pmod{2}$,*

$$\begin{aligned}\chi(W_{d^2}^0) &= -\frac{1}{32}d^2(d-1) \sum_{r|d} \frac{\mu(r)}{r^2} \\ \chi(W_{d^2}^1) &= -\frac{1}{32}d^2(d-3) \sum_{r|d} \frac{\mu(r)}{r^2}.\end{aligned}$$

Once cylinder cusps. As an application of the calculation of $[\overline{W}_{d^2}^\epsilon]$, we give formulas for the number of one-cylinder cusps of $W_{d^2}^\epsilon$. These formulas were established independently by Lelièvre and Royer in [LR06].

Theorem 13.7. *For any $d > 3$, the number of one-cylinder cusps of W_{d^2} is*

$$\frac{1}{6}d^2 \sum_{r|d} \frac{\mu(r)}{r^2} - \frac{1}{2}\phi(d); \quad (13.19)$$

the number of one-cylinder cusps of $W_{d^2}^0$ is

$$\frac{1}{24}d^2 \sum_{r|d} \frac{\mu(r)}{r^2}; \quad (13.20)$$

and the number of one-cylinder cusps of $W_{d^2}^1$ is

$$\frac{1}{8}d^2 \sum_{r|d} \frac{\mu(r)}{r^2} - \frac{1}{2}\phi(d). \quad (13.21)$$

Proof. The one-cylinder cusps of W_{d^2} are the points of the intersection,

$$S_{d^2}^1 \cap \overline{W}_{d^2}.$$

When $d > 3$ this intersection is transverse by Theorem 8.7, so $[\overline{W}_{d^2}] \cdot [\overline{S}_{d^2}^1]$ is equal to the number of one-cylinder cusps of W_{d^2} . Similarly, $[\overline{W}_{d^2}^\epsilon] \cdot [\overline{S}_{d^2}^1]$ is equal to the number of one-cylinder cusps of $W_{d^2}^\epsilon$.

Using (13.15), we obtain

$$[\overline{W}_{d^2}] \cdot [\overline{S}_{d^2}^1] = \frac{1}{6}d^2 \sum_{r|d} \frac{\mu(r)}{r^2} - \frac{1}{2}\phi(d),$$

which implies (13.19).

Now let's calculate $[\overline{S}_{d^2}^1] \cdot [\overline{W}_{d^2}^0]$. By Corollary 8.16, we have

$$\begin{aligned}\pi_B[\overline{S}_{d^2}^1] \cdot \pi_B[\overline{W}_{d^2}^0] &= \tau^* \left(\pi_B[\overline{S}_{d^2}^1] \right) \cdot \tau^* \left(\pi_B[\overline{W}_{d^2}^0] \right) \\ &= \pi_B[\overline{S}_{d^2}^2] \cdot \pi_B[\overline{W}_{d^2}^1] + (\pi_B[\overline{S}_{d^2}^2])^2 \\ &= \pi_B[\overline{S}_{d^2}^2] \cdot \pi_B[\overline{W}_{d^2}] + \pi_B[\overline{S}_{d^2}^2] \cdot \pi_B[\overline{W}_{d^2}^0] + (\pi_B[\overline{S}_{d^2}^2])^2\end{aligned} \quad (13.22)$$

Since $\overline{W}_{d^2}^0$ and $\overline{S}_{d^2}^2$ are disjoint, we have $[\overline{W}_{d^2}^0] \cdot [\overline{S}_{d^2}^2] = 0$, and it follows that

$$\pi_B[\overline{S}_{d^2}^2] \cdot \pi_B[\overline{W}_{d^2}^0] = \left(-\frac{1}{16}d^2 + \frac{1}{16}d\right) \sum_{r|d} \frac{\mu(r)}{r^2}. \quad (13.23)$$

Substituting (13.10), (13.12), and (13.23) into (13.22), we obtain

$$\pi_B[\overline{S}_{d^2}^1] \cdot \pi_B[\overline{W}_{d^2}^0] = \left(-\frac{7}{48}d^2 + \frac{3}{16}d\right) \sum_{r|d} \frac{\mu(r)}{r^2}. \quad (13.24)$$

Equation (13.24) yields

$$[\overline{S}_{d^2}^1] \cdot [\overline{W}_{d^2}^0] = \frac{1}{24}d^2 \sum_{r|d} \frac{\mu(r)}{r^2},$$

which implies (13.20). Equation (13.21) follows from (13.19) and (13.20). ■

14 Siegel-Veech Constants

As an application of our results, we record the Siegel-Veech constants counting cylinders on translation surfaces on the Teichmüller curves W_D and W_D^ϵ . This is basically a matter of plugging in our results into known formulas for these constants.

Given a translation surface (X, ω) , let

$$N((X, \omega), L) = \#\{\text{maximal cylinders of length at most } L \text{ on } (X, \omega)\}.$$

If (X, ω) lies on a Teichmüller curve, then Veech [Vee89] showed

$$N((X, \omega), L) \sim \frac{c}{\pi \text{Area}(X, \omega)} L^2. \quad (14.1)$$

The constant c , known as a Siegel-Veech constant, only depends on the Teichmüller curve on which (X, ω) lies. Let c_D be and c_D^ϵ be the Siegel-Veech constants associated to W_D and W_D^ϵ respectively.

Given a W_D -prototype $P = (a, b, c, \bar{q})$, define

$$v(P) = \frac{c}{\gcd(a, c)} \left(1 - \frac{a}{c} \lambda^2\right) \left(1 + \frac{1}{\lambda^2}\right),$$

where $\lambda = \lambda(P)$, the positive root of $ax^2 + bx + c = 0$.

Theorem 14.1. *If D is not square, then*

$$c_D = \frac{\sum_{P \in \mathcal{W}_D} v(P)}{-2\chi(W_D)}, \quad \text{and} \quad c_D^\epsilon = \frac{\sum_{P \in \mathcal{W}_D^\epsilon} v(P)}{-2\chi(W_D^\epsilon)},$$

where $\mathcal{W}_D^\epsilon$ is the set of W_D -prototypes of spin invariant ϵ .

Sketch of proof. To fix notation, assume W_D is connected as the proof is the same otherwise. Each cusp of W_D corresponds to a W_D -prototype P , and we associated to this cusp on p. 52 a surface (X_P, ω_P) on W_D with a decomposition into two cylinders C_1 and C_2 (say C_1 is the short cylinder). The subgroup of the Veech group of (X_P, ω_P) (the stabilizer of this surface in $\text{SL}_2\mathbb{R}$) which preserves the horizontal direction is generated by

$$g = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix},$$

where

$$t = \frac{c}{\gcd(a, c)}.$$

Let $i(C_i)$ be the order of the Dehn twist which g induces on C_i . We have,

$$\begin{aligned} i(C_1) &= -\frac{c}{\gcd(a, c)}, \quad \text{and} \\ i(C_2) &= \frac{a}{\gcd(a, c)}. \end{aligned}$$

We have

$$v(P) = \text{Area}(X_P, \omega_P) \sum_i \frac{i(C_i)}{\text{Area}(C_i)}.$$

It follows from Theorem 6.5 of [GJ00] that for any $(X, \omega) \in \Omega_1 W_D$,

$$N((X, \omega), L) \sim \frac{\sum_{P \in \mathcal{W}_D} v(P)}{\text{Area } W_D} L^2,$$

as desired. ■

We list the Siegel-Veech constants for $D < 100$ in Table 2, using the convention $c_D = c_D^0$ if $D \equiv 1 \pmod{8}$ (c_D^1 is the Galois conjugate of c_D^0 as we prove below). From numerical calculations, as $D \rightarrow \infty$, the constants appear to converge to 10, which by [EMZ03] is the Siegel-Veech constant for counting cylinders on a generic $(X, \omega) \in \Omega \mathcal{M}_2(2)$. It would be interesting to find a closed formula for c_D .

Arithmetic of Siegel-Veech constants. According to [GJ00], the Siegel-Veech constant c in (14.1) lies in the trace field of the Veech group of (X, ω) . We get more precise information in the case of the W_D :

Theorem 14.2. *Suppose D is not square. If $D \not\equiv 1 \pmod{8}$, then $c_D \in \mathbb{Q}$. Otherwise, c_D^0 and c_D^1 are Galois conjugate elements of $\mathbb{Q}(\sqrt{D})$.*

Proof. First assume $D \not\equiv 1 \pmod{8}$. We have the involution t on the set of W_D -prototypes, defined in §3. Actually, it was defined on Y_D -prototypes, but the definition works just as well in this case. This involution satisfies

$$v(t(P)) = v(P)',$$

it follows that $c_D' = c_D$ as desired.

Now suppose $D \equiv 1 \pmod{8}$. If it were true that $\epsilon(t(P)) = \epsilon(P) + 1$ – where ϵ is the spin invariant – then we would be done. This is not true, but we can modify t so that it is. Define a bijection on the set of W_D -prototypes,

$$s(P) = \begin{cases} P, & \text{if } a - b + c < 0 \text{ or } a \not\equiv b \pmod{2}; \\ P + (0, 0, 0, \gcd(a.b.c)), & \text{otherwise.} \end{cases}$$

Then $t' = s \circ t$ is also a bijection. One can check that it satisfies

$$\begin{aligned} \epsilon(t'(P)) &= \epsilon(P) + 1, \quad \text{and} \\ v(t'(P)) &= v(P)', \end{aligned}$$

from which the second claim follows. ■

Applications to billiards. Given a L-shaped polygon P (or more generally a rational angled polygon), there is a construction called *unfolding* which produced a translation surface from P . To construct the unfolding of P , join four copies of P to form a cross and glue opposite sides as in Figure 15. This yields a genus two translation surface (X, ω) with a single double zero. Billiards paths on P unfold to closed geodesics on (X, ω) , and we obtain a one-to-one correspondence between closed paths of length L on P and on the unfolding. Recall that we defined in §1 a L-shaped polygon $P(D)$ for each real quadratic discriminant D , and we defined $N(P(D), L)$ to be the number of families of closed billiards paths of length at least L .




Figure 15: Unfolding a L-shaped polygon

Theorem 14.3. *If D is not square, then*

$$N(P(D), L) \sim c(D) \frac{\pi}{4 \text{Area}(P(D))} L^2,$$

where $c(D) = c_D$, if $D \not\equiv 1 \pmod{8}$, and $c(D) = c_D^{(1+f)/2}$ if $D \equiv 1 \pmod{8}$, where f is the conductor of D .

Proof. This follows directly from (14.1) applied to the unfolding of $P(D)$. The factor of 4 is because the unfolding has 4 times the area. To see that the unfolding has discriminant D and spin invariant $\epsilon = (1 + f)/2$, note that the unfolding is $\text{GL}_2\mathbb{R}$ -equivalent to the surface (X_P, ω_P) associated to the W_D -prototype,

$$P = \begin{cases} (1, -1, \frac{1-D}{4}, 0); & \text{if } D \text{ is odd;} \\ (1, 0, -\frac{D}{4}, 0); & \text{if } D \text{ is even,} \end{cases}$$

defined on p. 52. The spin invariant is then given by Theorem 4.10. ■

D	c_D	D	c_D
5	$\frac{25}{3}$	56	$\frac{1796}{195}$
8	$\frac{28}{3}$	57	$\frac{23693}{2352} + \frac{29}{784}\sqrt{57}$
12	$\frac{26}{3}$	60	$\frac{2158}{231}$
13	$\frac{91}{9}$	61	$\frac{194651}{19305}$
17	$\frac{221}{24} + \frac{1}{8}\sqrt{17}$	65	$\frac{52429}{5376} + \frac{113}{1792}\sqrt{65}$
20	$\frac{31}{3}$	68	$\frac{413}{39}$
21	$\frac{133}{15}$	69	$\frac{26611}{2805}$
24	$\frac{148}{15}$	72	$\frac{18868}{1785}$
28	$\frac{82}{9}$	73	$\frac{3285}{352} + \frac{23}{864}\sqrt{73}$
29	$\frac{377}{35}$	76	$\frac{2822}{285}$
32	$\frac{190}{21}$	77	$\frac{116699}{12597}$
33	$\frac{473}{48} - \frac{11}{144}\sqrt{33}$	80	$\frac{12631}{1254}$
37	$\frac{9139}{945}$	84	$\frac{487}{51}$
40	$\frac{1924}{189}$	85	$\frac{336821}{32319}$
41	$\frac{8897}{960} - \frac{23}{320}\sqrt{41}$	88	$\frac{182236}{18837}$
44	$\frac{7682}{735}$	89	$\frac{702833}{68640} - \frac{831}{22880}\sqrt{89}$
45	$\frac{299}{33}$	92	$\frac{204178}{21945}$
48	$\frac{325}{33}$	93	$\frac{2823449}{270963}$
52	$\frac{1283}{135}$	96	$\frac{3194}{345}$
53	$\frac{228695}{21021}$	97	$\frac{44329}{4488} - \frac{1145}{40392}\sqrt{97}$

Table 2: Siegel-Veech constants for W_D

15 Lyapunov Exponents

Consider the rank $2g$ bundle $\mathcal{H}(\mathbb{R}) \rightarrow \Omega_1\mathcal{M}_g$ whose fiber over the surface (X, ω) is $H_1(X; \mathbb{R})$. The action of the diagonal subgroup of $\mathrm{SL}_2\mathbb{R}$ lifts to a linear action on $\mathcal{H}_1(\mathbb{R})$. This action is called the Kontsevich-Zorich cocycle. The $2g$ Lyapunov exponents of the Kontsevich-Zorich cocycle with respect to a finite, ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measure μ are of the form,

$$1 = \lambda_1(\mu) > \cdots > \lambda_g(\mu) > 0 > -\lambda_g(\mu) > \cdots > -\lambda_1(\mu) = -1.$$

The goal of this section is to prove the following unpublished theorem of Kontsevich and Zorich as an application of our previous results.

Theorem 15.1. *If μ is any finite, ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measure on $\Omega_1\mathcal{M}_2$, then*

$$\lambda_2(\mu) = \begin{cases} 1/3, & \text{if } \mu \text{ is supported on } \Omega_1\mathcal{M}_2(2); \\ 1/2, & \text{if } \mu \text{ is supported on } \Omega_1\mathcal{M}_2(1, 1). \end{cases}$$

Remark. When μ is period measure on E_D , with D nonsquare, the proof will require Theorems 15.10 and 15.11 below. These results will also be used in a future paper, and we will defer the proofs until then.

The proof of Theorem 15.1 is based on a formula due to Kontsevich [Kon97] for the sums of Lyapunov exponents in terms of some integrals over $\mathbb{P}\Omega\mathcal{M}_g$. Kontsevich states his formula for the Lyapunov exponents on the entire strata $\Omega_1\mathcal{M}_g(\mathbf{n})$, but his formula is equally valid for all ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant probability measures on $\Omega_1\mathcal{M}_2$ and we will state it for these.

Let E be the bundle over $\mathbb{P}\Omega\mathcal{M}_2$ obtained by pulling back $\Omega\mathcal{M}_2$ by $\mathbb{P}\Omega\mathcal{M}_2 \rightarrow \mathcal{M}_2$, and let L be the canonical sub-line-bundle whose fiber over $(X, [\omega])$ is $\mathbb{C}\omega$. Give E the Hodge metric,

$$h(\omega, \eta) = \int_X \omega \wedge \bar{\eta},$$

and define two-forms on $\mathbb{P}\Omega\mathcal{M}_2$,

$$\gamma_1 = c_1(E, h) \quad \text{and} \quad \gamma_2 = c_1(L, h).$$

For any $\mathrm{SL}_2\mathbb{R}$ -invariant measure μ on $\Omega_1\mathcal{M}_g$, let $\pi_*\mu$ be the pushforward to $\mathbb{P}\Omega\mathcal{M}_g$, and let $\mathbb{P}\mu$ be the disintegration of $\pi_*\mu$ with respect to the foliation $\mathcal{F}$ of $\mathbb{P}\Omega\mathcal{M}_g$ by images of $\mathrm{SL}_2\mathbb{R}$ -orbits. That is, $\mathbb{P}\mu$ is the unique transverse invariant measure such that the product of $\mathbb{P}\mu$ with the hyperbolic area measure on the leaves of $\mathcal{F}$ is $\pi_*\mu$. See [Bou69] for a discussion of disintegration.

Theorem 15.2 (Kontsevich). *For any finite ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measure μ on $\Omega_1\mathcal{M}_2$,*

$$\lambda_1(\mu) + \lambda_2(\mu) = \frac{\int_{\mathbb{P}\mu} \gamma_1}{\int_{\mathbb{P}\mu} \gamma_2}. \quad (15.1)$$

Uniform distribution. In order to evaluate (15.1), we will need the following weak uniform distribution result for Teichmüller curves. Given a space X with an $\mathrm{SL}_2\mathbb{R}$ -action, let $M(X)$ be the space of finite $\mathrm{SL}_2\mathbb{R}$ -invariant measures on X with the weak* topology. Let $C(\mathbf{n}) \subset M(\Omega_1\mathcal{M}_g(\mathbf{n}))$ be the convex cone spanned by the measures defined by hyperbolic area on those Teichmüller curves in $\Omega_1\mathcal{M}_g(\mathbf{n})$ which are generated by square-tiled surfaces. Similarly, let $C_{d^2} \subset M(\Omega_1E_{d^2}(1,1))$ be the convex cone spanned by those Teichmüller curves in $\Omega_1E_{d^2}(1,1)$ generated by square-tiled surfaces.

Theorem 15.3. *We have*

$$\begin{aligned}\mu'(\mathbf{n}) &\in C(\mathbf{n}), \quad \text{and} \\ \mu'_{d^2} &\in C_{d^2},\end{aligned}$$

where the measures on the left are the period measures on these strata defined in §4.

Let $\eta_m(\mathbf{n})$ and $\eta_m[d]$ be the measures on $\Omega\mathcal{M}_g(\mathbf{n})$ and $\Omega E_{d^2}(1,1)$ defined by putting δ -masses of equal weight on the square-tiled surfaces with at most m squares, normalized to have the same volumes as $\mu'(\mathbf{n})$ and μ'_{d^2} respectively. Let $\eta'_m(\mathbf{n})$ and $\eta'_m[d]$ be the projections to $\Omega_1\mathcal{M}_g(\mathbf{n})$ and $\Omega_1E_{d^2}(1,1)$. The following lemma is well-known.

Lemma 15.4. *We have*

$$\begin{aligned}\lim_{m \rightarrow \infty} \eta'_m(\mathbf{n}) &= \mu'(\mathbf{n}), \\ \lim_{m \rightarrow \infty} \eta'_m[d] &= \mu'_{d^2}\end{aligned}$$

in the weak* topology on measures.

Proof. We will prove the first statement, the other having the same proof. Let $\mu''(\mathbf{n})$ be the restriction of $\mu'(\mathbf{n})$ to $\Omega_{\leq 1}\mathcal{M}_g(\mathbf{n})$, and let

$$\eta''_m(\mathbf{n}) = (t_{1/\sqrt{m}})_* \eta_m(\mathbf{n}),$$

where t_r is the multiplication-by- r map on $\Omega\mathcal{M}_g$. Since $\mu''(\mathbf{n})$ and $\eta''_m(\mathbf{n})$ project to $\mu'(\mathbf{n})$ and $\eta'_m(\mathbf{n})$ respectively, it is enough to show that

$$\lim_{m \rightarrow \infty} \eta''_m(\mathbf{n}) = \mu''(\mathbf{n}).$$

This is easy to see in the period coordinates on $\Omega\mathcal{M}_g(\mathbf{n})$ (described in §4). In period coordinates, the square-tiled surfaces are exactly the points on the integer lattice, so $\eta''_m(\mathbf{n})$ consists of δ -masses of uniform weight on points of norm less than one on the $\mathbb{Z}/\sqrt{m}$ lattice. Since these lattices have mesh approaching zero, the measures converge to the uniform measure which is just $\mu''(\mathbf{n})$ in period coordinates. $\blacksquare$

Proof of Theorem 15.3. The set of square-tiled surfaces is $\mathrm{SL}_2\mathbb{Z}$ -invariant, so the measure $\eta'_m(\mathbf{n})$ is also $\mathrm{SL}_2\mathbb{Z}$ -invariant. Thus we can define

$$\nu_m(\mathbf{n}) = \int_{\mathrm{SL}_2\mathbb{R}/\mathrm{SL}_2\mathbb{Z}} \eta'_m(\mathbf{n}) d\rho \in M(\Omega_1\mathcal{M}_g(\mathbf{n})),$$

where ρ is Haar measure on $\mathrm{SL}_2\mathbb{R}$. The measure $\nu_m(\mathbf{n})$ is $\mathrm{SL}_2\mathbb{R}$ -invariant and supported on finitely many Teichmüller curves generated by square-tiled surfaces, so $\nu_m(\mathbf{n})$ lies in the cone $C(\mathbf{n})$.

Let $\Delta \subset \mathrm{SL}_2\mathbb{R}$ be a fundamental domain for $\mathrm{SL}_2\mathbb{Z}$. Given a compactly supported, continuous function f on $\Omega_1\mathcal{M}_g(\mathbf{n})$, we have,

$$\begin{aligned} \lim_{m \rightarrow \infty} \int_{\Omega_1\mathcal{M}_g(\mathbf{n})} f d\nu_m(\mathbf{n}) &= \lim_{m \rightarrow \infty} \int_{A \in \Delta} \int_{\Omega_1\mathcal{M}_g(\mathbf{n})} f \circ A d\eta'_m(\mathbf{n}) d\rho \\ &= \int_{A \in \Delta} \left(\lim_{m \rightarrow \infty} \int_{\Omega_1\mathcal{M}_g(\mathbf{n})} f \circ A d\eta'_m(\mathbf{n}) \right) d\rho \\ &= \int_{A \in \Delta} \int_{\Omega_1\mathcal{M}_g(\mathbf{n})} f \circ A d\mu'(\mathbf{n}) d\rho \\ &= \int_{A \in \Delta} \int_{\Omega_1\mathcal{M}_g(\mathbf{n})} f d\mu'(\mathbf{n}) d\rho \\ &= \mathrm{vol}(\mathrm{SL}_2\mathbb{R}/\mathrm{SL}_2\mathbb{Z}) \int_{\Omega_1\mathcal{M}_g(\mathbf{n})} f d\mu'(\mathbf{n}), \end{aligned}$$

where the second equality follows from the Dominated Convergence Theorem; the third equality follows from Lemma 15.4; and the fourth equality uses the $\mathrm{SL}_2\mathbb{R}$ -invariance of $\mu'(\mathbf{n})$. It follows that $\mu'(\mathbf{n}) \in C(\mathbf{n})$. The proof for μ'_{d^2} is the same. $\blacksquare$

The Siegel-Veech transform. Given any translation surface (X, ω) with an oriented saddle connection I , we associate the complex number $v(I) = \int_I \omega$. Let $V(X, \omega)$ be the collection of vectors in $\mathbb{C}$ associated with saddle connections on (X, ω) , together with the multiplicities with which they appear.

For any continuous, compactly supported function $f: \mathbb{C} \rightarrow \mathbb{R}$, the Siegel-Veech transform $\hat{f}: \Omega_1\mathcal{M}_g(\mathbf{n}) \rightarrow \mathbb{R}$ is defined by

$$\hat{f}(X, \omega) = \sum_{v \in V(X, \omega)} f(v).$$

Theorem 15.5 ([Vee98]). *Given any ergodic, $\mathrm{SL}_2\mathbb{R}$ -invariant measure μ on $\Omega_1\mathcal{M}_g(\mathbf{n})$, we have*

$$\frac{1}{\mathrm{vol} \mu} \int_{\Omega_1\mathcal{M}_g(\mathbf{n})} \hat{f} d\mu = c_{\mathrm{sc}}(\mu) \int_{\mathbb{R}^2} f. \quad (15.2)$$

The Siegel-Veech constant $c_{\text{sc}}(\mu)$ depends only on the measure μ .

If μ is supported on a Teichmüller curve C , we will also write $c_{\text{sc}}(C)$ for $c_{\text{sc}}(\mu)$. It follows from [Vee98] that for any (X, ω) on a Teichmüller curve C ,

$$N_{\text{sc}}((X, \omega), L) \sim c_{\text{sc}}(C) \frac{\pi}{\text{Area}(X, \omega)} L^2,$$

where N_{sc} is the counting function for saddle connections.

Bounded Siegel-Veech constants. We now show that Siegel-Veech constants are uniformly bounded over Teichmüller curves of a given genus.

Theorem 15.6. *There is a uniform bound,*

$$c_{\text{sc}}(C) \leq D,$$

where C ranges over all Teichmüller curves in $\Omega\mathcal{M}_g$, and D only depends on the genus.

This follows easily from the following results.

Theorem 15.7 ([Mas90], [EM01]). *Given any compact subset $K \subset \Omega\mathcal{M}_g(\mathbf{n})$, there is a constant $c(K)$ such that for any $(X, \omega) \in K$,*

$$N_{\text{sc}}((X, \omega), L) < c(K)L^2.$$

Theorem 15.8. *Given any connected component S of a stratum $\Omega_1\mathcal{M}_g(\mathbf{n})$, there is a compact subset $K \subset S$ which intersects the $\text{SL}_2\mathbb{R}$ orbit of each point in S .*

Theorem 15.8 is a corollary of the main result in Athreya's thesis [Ath06].

Proof of Theorem 15.6. For each connected component S of a stratum, choose a K as in Theorem 15.8. Then for every Teichmüller curve $C \subset S$, we have $c_{\text{sc}}(C) < c(K)$ by Theorem 15.7. Since there are only finitely many connected components of strata in each genus, we obtain a bound depending only on the genus. ■

Let $K_\epsilon \subset \Omega_1\mathcal{M}_g(\mathbf{n})$ be the locus of translation surfaces which have no saddle connection of length less than ϵ , and let $\tilde{K}_\epsilon$ be the complement of K_ϵ .

Corollary 15.9. *For any Teichmüller curve $C \subset \Omega_1\mathcal{M}_g(\mathbf{n})$, we have*

$$\frac{\text{vol}(C \cap \tilde{K}_\epsilon)}{\text{vol}(C)} < D\epsilon^2,$$

for some constant D depending only on the genus g .

Proof. Let χ_ϵ be the indicator function of the disk of radius ϵ . We have

$$\hat{\chi}_\epsilon \geq \chi_{\tilde{K}_\epsilon}.$$

Even though χ_ϵ is not continuous, we can apply (15.2) by the monotone convergence theorem. We have

$$\frac{\text{vol}(C \cap \tilde{K}_\epsilon)}{\text{vol}(C)} \leq \frac{1}{\text{vol}(C)} \int_C \hat{\chi}_\epsilon = c_{\text{sc}}(C) \pi \epsilon^2 \leq D \pi \epsilon^2,$$

where D is the constant of Theorem 15.6. ■

Extension of $\mathcal{F}_D$ to Y_D . We equipped the foliation $\mathcal{F}_D$ of X_D by $\text{SL}_2\mathbb{R}$ -orbits with a transverse invariant measure $\mathbb{P}\mu'_D$, so we can integrate 2-forms over $\mathcal{F}_D$ to obtain a closed current $\langle \mathcal{F}_D \rangle$ on X_D . The foliation $\mathcal{F}_D$ does not extend to a foliation of Y_D . In fact, $\mathcal{F}_D$ has isolated singularities at the points c_P as well as along the one-dimensional loci in the curves C_P consisting of forms whose horizontal foliation has two cylinders. Nevertheless, the current defined by $\mathcal{F}_D$ extends to a closed current on Y_D .

Theorem 15.10. *The foliation $\mathcal{F}_D$ defines a closed current $\langle \mathcal{F}_D \rangle$ on Y_D , defined by*

$$\langle \mathcal{F}_D \rangle(\omega) = \int_{\mathcal{F}_D} \omega$$

for each C^∞ 2-form ω on Y_D .

We will defer the proofs of this theorem and the next to a future paper. The idea of the proof of Theorem 15.10 is to compare $\int_{\mathcal{F}_D} \omega$ with $\text{vol}(\mu'_D)$ using the Schwartz lemma, and then to apply Theorem 4.8 that these volumes are finite.

Let $[\mathcal{F}_D]$ be the cohomology class in $H^2(Y_D; \mathbb{R})$ defined by $\langle \mathcal{F}_D \rangle$.

Theorem 15.11. *For any component C of $\overline{W}_D$, $\overline{P}_D$, or $\overline{S}_D^2$, we have*

$$[C] \cdot [\mathcal{F}_D] = 0.$$

The conclusion of this theorem is true for any closed leaves of a measured foliation which are not atoms of the transverse measure. The curves W_D , P_D , and S_D^2 are leaves of $\mathcal{F}_D$, but their cusps pass through singular points of the extension of $\mathcal{F}_D$ to Y_D . The proof of Theorem 15.11 amounts to estimating integrals of smooth forms around these singularities.

Proof of Theorem 15.1. Consider the pullbacks of the bundles E and L over $\mathbb{P}\Omega\mathcal{M}_2$ by $\pi: X_D \rightarrow \mathbb{P}\Omega\mathcal{M}_2$. We have

$$\begin{aligned} \pi^* E &= \Omega^1 X_D \oplus \Omega^2 X_D, \quad \text{and} \\ \pi^* L &= \Omega^1 X_D. \end{aligned}$$

Proposition 2.8 implies

$$\begin{aligned}\pi^*\gamma_1 &= \omega_1 + \omega_2, \quad \text{and} \\ \pi^*\gamma_2 &= \omega_1.\end{aligned}$$

Let μ be an ergodic $\mathrm{SL}_2\mathbb{R}$ -invariant measure on $\Omega_1\mathcal{M}_2$ whose support is not one of the strata $\Omega_1\mathcal{M}_2(2)$ or $\Omega_1\mathcal{M}_2(1,1)$. By McMullen's classification of ergodic, invariant measures, each such μ is supported on one of the eigenform loci $\Omega_1 E_D$. Then $\mathbb{P}\mu$ is a transverse, invariant measure to the foliation $\mathcal{F}_D$ of X_D , and (15.1) becomes

$$\lambda_2(\mu) = \frac{\int_{\mathbb{P}\mu} \omega_2}{\int_{\mathbb{P}\mu} \omega_1}. \quad (15.3)$$

If the ergodic, invariant measure μ is supported on a connected component C of W_D , then $\lambda_2(\mu) = 1/3$ by (15.3) and Corollary 12.3.

Now suppose μ is supported on a Teichmüller curve $C \subset \mathbb{P}\Omega\mathcal{M}_2(1,1)$ (so C is either the decagon curve D_{10} or a component of $W_{d^2}[n]$). By Corollary 10.3, we have

$$[\overline{W}_D] - [\overline{P}_D] + [\overline{S}_D^2] = -[\omega_1] + 2[\omega_2]. \quad (15.4)$$

Since $\overline{C}$ is disjoint from each of the three curves on the left, pairing $[\overline{C}]$ with both sides of (15.4) yields $\lambda_2(\mu) = 1/2$. If μ is the period measure μ'_D on E_D , then

$$\int_{\mathbb{P}\mu'_D} \omega_i = [\mathcal{F}_D](\omega_i).$$

Then we obtain $\lambda_2(\mu'_D) = 1/2$ by pairing $[\mathcal{F}_D]$ with both sides of (15.4) and applying Theorem 15.11.

It remains to deal with the measures $\mu'(\mathbf{n})$ on the strata $\Omega_1\mathcal{M}_2(\mathbf{n})$. Given a point $p \in \mathbb{P}\Omega\mathcal{M}_2$, let L be the leaf of $\mathcal{F}$ through p . Define

$$f(p) = \frac{\gamma_1|_L}{\gamma_2|_L}.$$

This is a well-defined real number because it is the ratio of top-degree forms.

Lemma 15.12. *As a function on $\mathbb{P}\Omega\mathcal{M}_2$, we have*

$$1 \leq f \leq 2.$$

Proof. Since f is continuous, it suffices to prove this bound for every Teichmüller curve $C \subset \mathbb{P}\Omega\mathcal{M}_2$. The curve C lies on some Hilbert modular surface X_D . Let

$$\tilde{C} \subset \mathbb{H} \times \mathbb{H}$$

be a connected component of the inverse image of C in the universal cover of X_D , and let $\pi_i: \tilde{C} \rightarrow \mathbb{H}$ be the two projections. Since C is transverse to the vertical foliation $\mathcal{A}_D$ of X_D , the first projection π_1 is a conformal isomorphism

of $\tilde{C}$ with $\mathbb{H}$. By the Schwartz lemma, the projection π_2 is a contraction. Since ω_i is the pullback of the hyperbolic area form on $\mathbb{H}$ by ω_i , we have

$$0 \leq \frac{\omega_2|_C}{\omega_1|_C} \leq 1.$$

The claim follows, since

$$f|_C = \frac{\omega_1|_C + \omega_2|_C}{\omega_1|_C}.$$

■

We need to show that

$$\frac{1}{\text{vol } \mu'(\mathbf{n})} \int_{\Omega_1 \mathcal{M}_2(\mathbf{n})} f d\mu'(\mathbf{n}) = \begin{cases} 1/3; & \text{if } \mathbf{n} = (2); \\ 1/2; & \text{if } \mathbf{n} = (1, 1), \end{cases}$$

where we regard f as a function on $\Omega_1 \mathcal{M}_2(\mathbf{n})$ by pulling it back from the projectivization. Let $\{\nu_m\}$ be a sequence of measures supported on Teichmüller curves in $\Omega_1 \mathcal{M}_2(\mathbf{n})$ and converging to $\mu'(\mathbf{n})$. Since the average of f over any Teichmüller curve C is either $1/2$ or $1/3$, depending on the stratum in which C lies, it is enough to show that

$$\frac{1}{\text{vol } \nu_m} \int f d\nu_m \rightarrow \frac{1}{\text{vol } \mu'(\mathbf{n})} \int f d\mu'(\mathbf{n}).$$

Recall that we defined K_ϵ to be the locus of translation surfaces with no saddle connection shorter than ϵ . Since K_ϵ is compact (as is shown in [KMS86]), $\text{vol } \Omega \mathcal{M}_2(\mathbf{n}) < \infty$, and the K_ϵ exhaust $\Omega_1 \mathcal{M}_2(\mathbf{n})$ as $\epsilon \rightarrow 0$, we can choose for any $\delta > 0$ an ϵ small enough that

$$\frac{\mu'(\mathbf{n})(\tilde{K}_\epsilon)}{\text{vol } \mu'(\mathbf{n})} < \frac{\delta}{2}. \quad (15.5)$$

By Corollary 15.9, we can also choose ϵ small enough that

$$\frac{\nu_m(\mathbf{n})(\tilde{K}_\epsilon)}{\text{vol } \nu_m(\mathbf{n})} < \frac{\delta}{2}. \quad (15.6)$$

Let g_ϵ be a continuous, compactly supported function on $\Omega_1 \mathcal{M}_2(\mathbf{n})$ such that $g \equiv 1$ on K_ϵ , and $0 \leq g \leq 1$. We have,

$$\begin{aligned} & \lim_{m \rightarrow \infty} \left| \frac{1}{\text{vol } \nu_m} \int f d\nu_m - \frac{1}{\text{vol } \mu'(\mathbf{n})} \int f d\mu'(\mathbf{n}) \right| \\ & \leq \lim_{m \rightarrow \infty} \left| \frac{1}{\text{vol } \nu_m} \int g_\epsilon f d\nu_m - \frac{1}{\text{vol } \mu'(\mathbf{n})} \int g_\epsilon f d\mu'(\mathbf{n}) \right| \\ & \quad + \frac{\mu'(\mathbf{n})(\tilde{K}_\epsilon)}{\text{vol } \mu'(\mathbf{n})} + \frac{\nu_m(\mathbf{n})(\tilde{K}_\epsilon)}{\text{vol } \nu_\epsilon(\mathbf{n})} \\ & < \delta, \end{aligned}$$

since the limit of the first term is zero by Theorem 15.3, and the other two terms are bounded by (15.5) and (15.6).

This completes the proof of Theorem 15.1. Note that the last part of the proof applies to the $\Omega_1 E_{d^2}$, so we can prove Theorem 15.1 for these spaces without appealing to Theorems 15.10 and 15.11.

A Normal varieties

In this section, we record standard facts about normality for algebraic varieties and analytic spaces which we use in this paper. We will consider all of our algebraic varieties to be over $\mathbb{C}$.

Normal algebraic varieties. A point p on an algebraic variety X is said to be *normal* if the local ring $\mathcal{O}_p$ of X at p is an integral domain which is integrally closed in its field of fractions. A variety X is *normal* if it is normal at each of its points.

A *normalization* of a variety X is a normal variety Y together with a finite surjective morphism $\pi: Y \rightarrow X$. More generally (following [Mum99]), if X is an irreducible variety, and if L is a finite algebraic extension of $K(X)$, then a *normalization of X in L* is a normal variety Y with function field $K(Y) = L$, together with a finite surjective morphism $\pi: Y \rightarrow X$ such that $\pi^*: K(X) \rightarrow K(Y) = L$ is the given inclusion of $K(X)$ in L . If $L = K(X)$, this is just the usual normalization of X .

Theorem A.1 ([Mum99, Theorem III:8.3]). *For any irreducible variety X and finite algebraic extension L of $K(X)$, there is a normalization of X in L , and any two such normalizations are equivalent.*

Normalization is also characterized by a universal property. In the case, where $L = K(X)$, this is an exercise in [Har77, p.91].

Theorem A.2. *The normalization $\pi: Y \rightarrow X$ of X in $L = K(X)$ has the following universal property. Let Z be a normal variety with a dominant morphism $q: Z \rightarrow X$, and let $r: K(X) \rightarrow K(Z)$ be an isomorphism such that $r \circ \pi^* = q^*$ as inclusions $K(X) \rightarrow K(Z)$. Then there is a unique morphism $s: Z \rightarrow Y$ such that $s^* = r$ and $\pi \circ s = q$.*

We will also have use for the following theorems:

Theorem A.3 ([Mum99, III:8.4]). *The normalization of a projective variety in a finite algebraic extension is projective.*

Theorem A.4 (Zariski's Main Theorem, [Har77]). *Let $f: X \rightarrow Y$ be a birational morphism of projective varieties, and assume that Y is normal. Then for every $y \in Y$, the fiber $f^{-1}(y)$ is connected.*

Theorem A.5 (Zariski, [ZS60, Theorem VIII.32]). *If X is a normal variety, $p \in X$, and $\widehat{\mathcal{O}}_p$ is the completion of $\mathcal{O}_p$, then $\widehat{\mathcal{O}}_p$ is an integrally closed integral domain.*

Normal analytic spaces. A complex analytic space is a ringed space which is locally modeled on analytic subvarieties of $\mathbb{C}^n$. We start by recalling some basic notions about complex analytic spaces. See [GR65] and [Gun70] for details.

Consider the set of pairs (U, X) , where $U \subset \mathbb{C}^n$ is a neighborhood of some point p , and $X \subset U$ is an analytic subvariety of U . We consider two such pairs

to be equivalent if the varieties agree on some common neighborhood of p . An equivalence class of such pairs is called a *germ on an analytic variety at p* . A germ is considered to be irreducible if it contains no proper subgerms.

Theorem A.6 ([Gun70, p.11]). *Any germ of an analytic subvariety at p can be written uniquely as the union of finitely many irreducible germs of analytic varieties at p .*

Given a point $p \in X$, an analytic subvariety of an open subset of $\mathbb{C}^n$, a *branch* of X through p is an irreducible subgerm of the germ of X at p . Informally, a branch of X through p is a connected component of $U \cap (X \setminus X_{\text{sing}})$, where U is a small and sufficiently regular neighborhood of p , and X_{sing} is the singular set of X . Given a point p in an analytic space X , let $\mathcal{O}_p$ be the local ring of germs of holomorphic functions on X at p . Just as for algebraic varieties, we say that p is a *normal point* of X if $\mathcal{O}_p$ is an integrally closed integral domain. We say that X is *normal* if each of its points is normal.

There is also a geometric characterization of normality. Following [Whi72], we say that a function f on an open subset V of an analytic variety X is *weakly holomorphic* if there is a subvariety $V' \subset V$ with the following properties:

- V' is nowhere dense in V , and $V_{\text{sing}} \subset V'$.
- f is holomorphic in $V \setminus V'$.
- f is locally bounded in V .

It follows from [Whi72, Theorem 4.10I] that an analytic space X is weakly holomorphic at p if and only if every germ of a weakly holomorphic function at p is in fact holomorphic.

A *normalization* of an analytic space X is a normal analytic variety Y together with a holomorphic map $p: Y \rightarrow X$ such that

- $p: Y \rightarrow X$ is proper and has finite fibers.
- If X_{sing} is the singular set of X , and $A = p^{-1}(X_{\text{sing}})$, then $Y \setminus A$ is dense in Y , and $p: Y \setminus A \rightarrow X \setminus X_{\text{sing}}$ is biholomorphic.

Theorem A.7 ([Lau71, p.38]). *Every analytic space has a unique normalization.*

Theorem A.8 ([Lau71, p.37]). *If $\pi: Y \rightarrow X$ is the normalization of X , then for each $p \in X$, the map π induces a bijective correspondence between the points of $\pi^{-1}(p)$ and the branches of X through P .*

Theorem A.9. *If X is an algebraic variety, and $\pi: Y \rightarrow X$ is the normalization of X as an algebraic variety then it is also the normalization of X as an analytic space.*

Proof. By the definition of the normalization of an analytic space, we need only to show that Y is normal as an analytic space. For this proof, write $\mathcal{O}_p^{\text{an}}$ for the local ring of germs of holomorphic functions on X at p , to differentiate from the local ring of $\mathcal{O}_p$ of algebraic functions at p . Let $\widehat{\mathcal{O}}_p$ be the completion of $\mathcal{O}_p$.

Let f in the quotient field $K(\mathcal{O}_p^{\text{an}})$ be integral over $\mathcal{O}_p^{\text{an}}$. Since $\mathcal{O}_p^{\text{an}} \subset \widehat{\mathcal{O}}_p$, it follows that f is integral over $\widehat{\mathcal{O}}_p$. Since $\widehat{\mathcal{O}}_p$ is integrally closed by Theorem A.5, it follows that $f \in \widehat{\mathcal{O}}_p$. Then $f \in \mathcal{O}_p^{\text{an}}$ because $K(\mathcal{O}_p^{\text{an}}) \cap \widehat{\mathcal{O}}_p = \mathcal{O}_p^{\text{an}}$ in $K(\widehat{\mathcal{O}}_p)$. ■

A *resolution* of an analytic space X is a complex manifold Y together with a proper analytic map $\pi: Y \rightarrow X$ such that $\pi: Y \setminus \pi^{-1}(X_{\text{sing}}) \rightarrow X \setminus X_{\text{sing}}$ is biholomorphic and such that $\pi^{-1}(X \setminus X_{\text{sing}})$ is dense in Y . The following was proved by Mumford in [Mum61], by Grauert in [Gra62], and also by Laufer in [Lau71, Theorem 4.4].

Theorem A.10. *Let $\pi: Y \rightarrow X$ be a resolution of X , and let $p \in X$ be an isolated normal point. Suppose that $A = \pi^{-1}(p)$ is the union of irreducible curves A_i which are nonsingular and intersect transversely, with at most two branches of $\bigcup_i A_i$ passing through any point. Then the intersection matrix*

$$(A_i \cdot A_j)$$

is negative definite.

References

- [Abi77] W. Abikoff. Degenerating families of Riemann surfaces. *Ann. of Math. (2)*, 105(1):29–44, 1977.
- [Ahl60] L. Ahlfors. The complex analytic structure of the space of closed Riemann surfaces. In *Analytic functions*, pages 45–66. Princeton Univ. Press, Princeton, N.J., 1960.
- [Ath06] J. Athreya. Quantitative recurrence and large deviations for Teichmüller geodesic flow. *Geom. Dedicata*, 119:121–140, 2006.
- [Bai58] W. Baily, Jr. Satake’s compactification of V_n . *Amer. J. Math.*, 80:348–364, 1958.
- [Bai60] W. Baily, Jr. On the moduli of Jacobian varieties. *Ann. of Math. (2)*, 71:303–314, 1960.
- [Bai62] W. Baily, Jr. On the theory of θ -functions, the moduli of abelian varieties, and the moduli of curves. *Ann. of Math. (2)*, 75:342–381, 1962.
- [Ber61a] L. Bers. Correction to “Spaces of Riemann surfaces as bounded domains”. *Bull. Amer. Math. Soc.*, 67:465–466, 1961.
- [Ber61b] L. Bers. Holomorphic differentials as functions of moduli. *Bull. Amer. Math. Soc.*, 67:206–210, 1961.
- [Ber73] L. Bers. Fiber spaces over Teichmüller spaces. *Acta. Math.*, 130:89–126, 1973.
- [Ber74] L. Bers. On spaces of Riemann surfaces with nodes. *Bull. Amer. Math. Soc.*, 80:1219–1222, 1974.
- [Ber81] L. Bers. Finite-dimensional Teichmüller spaces and generalizations. *Bull. Amer. Math. Soc. (N.S.)*, 5(2):131–172, 1981.
- [BL04] C. Birkenhake and H. Lange. *Complex Abelian Varieties*, volume 302 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2004.
- [BM] I. Bouw and M. Möller. Teichmüller curves, triangle groups, and Lyapunov exponents, arXiv:math.AG/0511738.
- [Bou69] N. Bourbaki. *Éléments de mathématique. Fasc. XXXV. Livre VI: Intégration. Chapitre IX: Intégration sur les espaces topologiques séparés*. Actualités Scientifiques et Industrielles, No. 1343. Hermann, Paris, 1969.

- [BS66] A. Borevich and I. Shafarevich. *Number Theory*. Translated from the Russian by Newcomb Greenleaf. Pure and Applied Mathematics, Vol. 20. Academic Press, New York, 1966.
- [Cal04] K. Calta. Veech surfaces and complete periodicity in genus two. *J. Amer. Math. Soc.*, 17(4):871–908, 2004.
- [Coh75] H. Cohen. Sums involving the values at negative integers of L -functions of quadratic characters. *Math. Ann.*, 217(3):271–285, 1975.
- [CR02] W. Chen and Y. Ruan. Orbifold Gromov-Witten theory. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 25–85. Amer. Math. Soc., Providence, RI, 2002.
- [DM69] P. Deligne and D. Mumford. The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.*, (36):75–109, 1969.
- [EM01] A. Eskin and H. Masur. Asymptotic formulas on flat surfaces. *Ergodic Theory Dynam. Systems*, 21(2):443–478, 2001.
- [EMS03] A. Eskin, H. Masur, and M. Schmoll. Billiards in rectangles with barriers. *Duke Math. J.*, 118(3):427–463, 2003.
- [EMZ03] A. Eskin, H. Masur, and A. Zorich. Moduli spaces of abelian differentials: the principal boundary, counting problems, and the Siegel-Veech constants. *Publ. Math. Inst. Hautes Études Sci.*, (97):61–179, 2003.
- [EO01] A. Eskin and A. Okounkov. Asymptotics of numbers of branched coverings of a torus and volumes of moduli spaces of holomorphic differentials. *Invent. Math.*, 145(1):59–103, 2001.
- [EOP] A. Eskin, A. Okounkov, and R. Pandharipande. The theta characteristic of a branched covering, arXiv:math.AG/0312186.
- [GH78] P. Griffiths and J. Harris. *Principles of Algebraic Geometry*. Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics.
- [GJ00] E. Gutkin and C. Judge. Affine mappings of translation surfaces: geometry and arithmetic. *Duke Math. J.*, 103(2):191–213, 2000.
- [GR65] R. Gunning and H. Rossi. *Analytic Functions of Several Complex Variables*. Prentice-Hall Inc., Englewood Cliffs, N.J., 1965.
- [Gra62] H. Grauert. Über Modifikationen und exzeptionelle analytische Mengen. *Math. Ann.*, 146:331–368, 1962.

- [Gun70] R. Gunning. *Lectures on Complex Analytic Varieties: The Local parametrization theorem*. Mathematical Notes. Princeton University Press, Princeton, N.J., 1970.
- [Har66] R. Hartshorne. *Residues and Duality*. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20. Springer-Verlag, Berlin, 1966.
- [Har77] R. Hartshorne. *Algebraic Geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [Hir73] F. Hirzebruch. Hilbert modular surfaces. *Enseignement Math. (2)*, 19:183–281, 1973.
- [HL06] P. Hubert and S. Lelièvre. Prime arithmetic Teichmüller discs in $\mathcal{H}(2)$. *Israel J. Math.*, 151:281–321, 2006.
- [HM98] J. Harris and I. Morrison. *Moduli of Curves*, volume 187 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1998.
- [IR90] K. Ireland and M. Rosen. *A Classical Introduction to Modern Number Theory*, volume 84 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1990.
- [KM76] F. Knudsen and D. Mumford. The projectivity of the moduli space of stable curves. I. Preliminaries on “det” and “Div”. *Math. Scand.*, 39(1):19–55, 1976.
- [KMS86] S. Kerckhoff, H. Masur, and J. Smillie. Ergodicity of billiard flows and quadratic differentials. *Ann. of Math. (2)*, 124(2):293–311, 1986.
- [Kon97] M. Kontsevich. Lyapunov exponents and Hodge theory. In *The mathematical beauty of physics (Saclay, 1996)*, volume 24 of *Adv. Ser. Math. Phys.*, pages 318–332. World Sci. Publishing, River Edge, NJ, 1997.
- [Kra90] I. Kra. Horocyclic coordinates for Riemann surfaces and moduli spaces. I. Teichmüller and Riemann spaces of Kleinian groups. *J. Amer. Math. Soc.*, 3(3):499–578, 1990.
- [KS00] R. Kenyon and J. Smillie. Billiards on rational-angled triangles. *Comment. Math. Helv.*, 75(1):65–108, 2000.
- [KZ] M. Kontsevich. and A. Zorich. Lyapunov exponents and Hodge theory, arXiv:hep-th/9701164.
- [KZ03] M. Kontsevich and A. Zorich. Connected components of the moduli spaces of Abelian differentials with prescribed singularities. *Invent. Math.*, 153(3):631–678, 2003.

- [Lau71] H. Laufer. *Normal Two-Dimensional Singularities*. Princeton University Press, Princeton, N.J., 1971. Annals of Mathematics Studies, No. 71.
- [Lel06] S. Lelièvre. Siegel-Veech constants in $\mathcal{H}(2)$. *Geom. Topol.*, 10:1157–1172 (electronic), 2006.
- [LR06] S. Lelièvre and E. Royer. Orbitwise countings in $\mathcal{H}(2)$ and quasimodular forms. *International Mathematics Research Notices*, 2006:Article ID 42151, 30 pages, 2006.
- [Mar87] A. Marden. Geometric complex coordinates for Teichmüller space. In *Mathematical aspects of string theory (San Diego, Calif., 1986)*, volume 1 of *Adv. Ser. Math. Phys.*, pages 341–354. World Sci. Publishing, Singapore, 1987.
- [Mas82] H. Masur. Interval exchange transformations and measured foliations. *Ann. of Math. (2)*, 115(1):169–200, 1982.
- [Mas90] H. Masur. The growth rate of trajectories of a quadratic differential. *Ergodic Theory Dynam. Systems*, 10(1):151–176, 1990.
- [McMa] C. McMullen. Dynamics of $\mathrm{SL}_2\mathbb{R}$ over moduli space in genus two. *Preprint*, 10/2003.
- [McMb] C. McMullen. Foliations of Hilbert modular surfaces. *Preprint*, 2/2005.
- [McM03a] C. McMullen. Billiards and Teichmüller curves on Hilbert modular surfaces. *J. Amer. Math. Soc.*, 16(4):857–885, 2003.
- [McM03b] C. McMullen. Teichmüller geodesics of infinite complexity. *Acta Math.*, 191(2):191–223, 2003.
- [McM05a] C. McMullen. Teichmüller curves in genus two: Discriminant and spin. *Math. Ann.*, 333(1):87–130, 2005.
- [McM05b] C. McMullen. Teichmüller curves in genus two: the decagon and beyond. *J. Reine Angew. Math.*, 582:173–199, 2005.
- [McM06] C. McMullen. Teichmüller curves in genus two: torsion divisors and ratios of sines. *Invent. Math.*, 165(3):651–672, 2006.
- [Miy89] T. Miyake. *Modular Forms*. Springer-Verlag, Berlin, 1989. Translated from the Japanese by Yoshitaka Maeda.
- [Möl06] M. Möller. Periodic points on Veech surfaces and the Mordell-Weil group over a Teichmüller curve. *Invent. Math.*, 165(3):633–649, 2006.
- [MT02] H. Masur and S. Tabachnikov. Rational billiards and flat structures. In *Handbook of dynamical systems, Vol. 1A*, pages 1015–1089. North-Holland, Amsterdam, 2002.

- [Mum61] D. Mumford. The topology of normal singularities of an algebraic surface and a criterion for simplicity. *Inst. Hautes Études Sci. Publ. Math.*, (9):5–22, 1961.
- [Mum71] D. Mumford. The structure of the moduli spaces of curves and Abelian varieties. In *Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 1*, pages 457–465. Gauthier-Villars, Paris, 1971.
- [Mum77] D. Mumford. Hirzebruch’s proportionality theorem in the noncompact case. *Invent. Math.*, 42:239–272, 1977.
- [Mum83] D. Mumford. Towards an enumerative geometry of the moduli space of curves. In *Arithmetic and geometry, Vol. II*, volume 36 of *Progr. Math.*, pages 271–328. Birkhäuser Boston, Boston, MA, 1983.
- [Mum99] D. Mumford. *The Red Book of Varieties and Schemes*, volume 1358 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, expanded edition, 1999. Includes the Michigan lectures (1974) on curves and their Jacobians, With contributions by Enrico Arbarello.
- [Nam73] Y. Namikawa. On the canonical holomorphic map from the moduli space of stable curves to the Igusa monoidal transform. *Nagoya Math. J.*, 52:197–259, 1973.
- [Rat94] M. Ratner. Invariant measures and orbit closures for unipotent actions on homogeneous spaces. *Geom. Funct. Anal.*, 4(2):236–257, 1994.
- [Sat56a] I. Satake. On a generalization of the notion of manifold. *Proc. Nat. Acad. Sci. U.S.A.*, 42:359–363, 1956.
- [Sat56b] I. Satake. On the compactification of the Siegel space. *J. Indian Math. Soc. (N.S.)*, 20:259–281, 1956.
- [Sch05] M. Schmoll. Spaces of elliptic differentials. In *Algebraic and topological dynamics*, volume 385 of *Contemp. Math.*, pages 303–320. Amer. Math. Soc., Providence, RI, 2005.
- [Ser56] J. P. Serre. Géométrie algébrique et géométrie analytique. *Ann. Inst. Fourier, Grenoble*, 6:1–42, 1955–1956.
- [Ser59] J. P. Serre. *Groupes algébriques et corps de classes*. Publications de l’institut de mathématique de l’université de Nancago, VII. Hermann, Paris, 1959.
- [Sie36] C. Siegel. The volume of the fundamental domain for some infinite groups. *Trans. Amer. Math. Soc.*, 39(2):209–218, 1936.

- [Sie69] C. Siegel. Berechnung von Zetafunktionen an ganzzahligen Stellen. *Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II*, 1969:87–102, 1969.
- [vdG88] G. van der Geer. *Hilbert Modular Surfaces*, volume 16 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1988.
- [Vee86] W. Veech. The Teichmüller geodesic flow. *Ann. of Math. (2)*, 124(3):441–530, 1986.
- [Vee89] W. Veech. Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards. *Invent. Math.*, 97(3):553–583, 1989.
- [Vee90] W. Veech. Moduli spaces of quadratic differentials. *J. Analyse Math.*, 55:117–171, 1990.
- [Vee92] W. Veech. The billiard in a regular polygon. *Geom. Funct. Anal.*, 2(3):341–379, 1992.
- [Vee98] W. Veech. Siegel measures. *Ann. of Math. (2)*, 148(3):895–944, 1998.
- [Whi72] H. Whitney. *Complex Analytic Varieties*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1972.
- [Wol85] S. Wolpert. On obtaining a positive line bundle from the Weil-Petersson class. *Amer. J. Math.*, 107(6):1485–1507 (1986), 1985.
- [Zor99] A. Zorich. How do the leaves of a closed 1-form wind around a surface? In *Pseudoperiodic Topology*, volume 197 of *Amer. Math. Soc. Transl. Ser. 2*, pages 135–178. Amer. Math. Soc., Providence, RI, 1999.
- [Zor06] A. Zorich. Flat surfaces. In *Frontiers in Number Theory, Physics and Geometry. Volume 1: On random matrices, zeta functions and dynamical systems*, pages 439–586. Springer-Verlag, Berlin, 2006.
- [ZS60] O. Zariski and P. Samuel. *Commutative Algebra. Vol. II*. The University Series in Higher Mathematics. D. Van Nostrand Co., Inc., Princeton, N. J.-Toronto-London-New York, 1960.